ORIGINAL ARTICLES
(CC BY-SA)
UDC: 616.132.2-085.8-036
DOI: https://doi.org/10.2298/VSP200810126S

The effects of cardiac rehabilitation on haemodynamic parameters measured by impedance cardiography in patients with coronary artery disease

Uticaj kardiovaskularne rehabilitacije na hemodinamski status bolesnika sa koronarnom bolešću srca procenjen korišćenjem kardiografske impedance

Milovan Stojanović*, Marina Deljanin-Ilić*†, Stevan Ilić‡, Dejan Petrović*†, Bojan Ilić*

***Institute for Treatment and Rehabilitation "Niška Banja", Niš, Serbia; †University of Niš, Faculty of Medicine, Niš, Serbia; ‡Clinic for Internal Diseases CardioPoint, Niš, Serbia**

Abstract

Background/Aim. Well-organized cardiovascular rehabilitation (CVR) reduces cardiovascular burden by influencing cardiovascular risk factors, improving the quality of life, and reducing mortality and hospital readmission. However, its effects on hemodynamic status are largely unknown. The aim of our study was to evaluate the influence of a three-week CVR program on hemodynamic status and to investigate if there is a correlation between physical strain tolerance and hemodynamic parameters measured by impedance cardiography (ICG) before and after the CVR program in patients with coronary artery disease (CAD). **Methods.** Fifty-two patients attended a three-week CVR program. At the beginning and the end of the rehabilitation program, laboratory tests, exercise stress tests (EST), and ICG measurements were taken. **Results.** Patients showed better strain tolerance on the second exercise stress test (EST2) by achieving a higher strain level ($Z = 2.315$; $p = 0.021$) and a longer duration of the test ($Z = 2.305$; $p = 0.021$). There was a strong positive correlation between the level of EST2 and cardiac output (CO) ($r = 0.538$; $p < 0.001$) and stroke volume (SV) ($r = 0.380$; $p = 0.017$) on the second ICG (ICG2). Moreover, there was a strong negative correlation between EST2 level and systemic vascular resistance (SVR) ($r = -0.472$; $p = 0.002$) and SVR index (SSVRI) ($r = -0,407$; $p = 0.010$) on ICG2. There was a strong positive correlation between EST2 duration and CO ($r = 0.517$; $p = 0.001$) as well as between EST2 duration and SV ($r = 0.340$; $p = 0.034$), and a strong negative correlation between EST2 duration and SVR ($r = -0.504$; $p = 0.001$) as well as between EST2 duration and SVRI ($r = -0.448$; $p = 0.004$), according to ICG2. **Conclusion.** Our study showed that a well-designed CVR program can lead to better physical strain tolerance in patients with CAD. Furthermore, CVR led to a significant positive correlation between EST and CO as well as between EST and SV measured by ICG. On the other hand, there was a significant negative correlation between EST and vascular-related parameters according to ICG at the end of the CVR program.

Key words:
cardiography, impedance; coronary artery disease; myocardial infarction; physical exertion; plethysmography, impedance; rehabilitation.

Apstrakt

Uvod/Cilj. Dobro organizovana kardiovaskularna rehabilitacija (KVR) smanjuje kardiovaskularno opterećenje uticajem na faktore rizika od kardiovaskularnih bolesti, poboljšavajući kvalitet života i smanjujući smrtnost i ponovni kardiovaskularni događaj. Međutim, njeni efekti na hemodinamski status bolesnika uglavnom su nepoznati. Cilj rada bio je da se proceni uticaj tronedeljnog programa KVR na hemodinamski status i da se istraži postojanje povezanosti između tolerancije fizičkog napora i hemodinamskih parametara merenih kardiografskom impedancom (*impedance cardiography* – ICG) pre i posle programa KVR kod bolesnika sa oboljenjem koronarnih arterija. **Metode.** Ukupno 52 bolesnika učestvovala su u tronedeljnom programu KVR. Na početku i na kraju programa rehabilitacije vršeni su laboratorijski testovi, testovi fizičkim opterećenjem (TFO) i ICG merenja. **Rezultati.** Bolesnici su pokazali bolju toleranciju fizičkog napora na drugom TFO (TFO2) dostižući viši nivo opterećenja ($Z = 2,315$; $p = 0,021$) i duže trajanje testa ($Z = 2,305$; $p = 0,021$). Postojala je jaka korelacija između nivoa drugog TFO i minutnog volumena (MV) ($r = 0,538$; $p < 0,001$) i udarnog volumena (UV) ($r = 0,380$; $p = 0,017$) na drugoj ICG (ICG2). Takođe, postojala je jaka negativna korelacija između

Correspondence to: Milovan Stojanović, Institute for Treatment and Rehabilitation "Niška Banja", Srpskih junaka 2, 18 205 Niš, Serbia. E-mail: milovanstojanovic1987@gmail.com

nivoa TFO2 i sistemskog vaskularnog otpora (SVO) (r = -0,472; p = 0,002) i indeksa SVO (SVOI) (r = -0,407; p = 0,010) na ICG2. Postojala je jaka pozitivna korelacija između dužine trajanja TFO2 i MV (r = 0,517; p = 0,001), kao i između dužine trajanja TFO2 i UV (r = 0,340; p = 0,034), i jaka negativna koralcija između trajanja TFO2 i SVO (r = -0,504; p = 0,001), kao i između trajanja TFO2 i SVOI (r = -0,448; p = 0,004) na ICG2. **Zaključak.** Naša studija je pokazala da dobro dizajniran program KVR može dovesti do poboljšanja tolerancije fizičkog napora kod bolesnika sa oboljenjem koronarnih arterija. Takođe, KVR je dovela do pozitivne korelacije između TFO i MV kao i TFO i UV procenjenih primenom ICG. Sa druge strane, postojala je značajna negativna korelacija između TFO i vaskularne rezistencije procenjene korišćenjem ICG na kraju programa KVR.

Ključne reči:
kardiografija, impedansna; koronarna bolest; infarkt miokarda; napor, fizički; pletizmografija, impedansna; rehabilitacija.

Introduction

Cardiovascular rehabilitation (CVR) is of great importance in the secondary prevention of cardiovascular diseases (CVD). On the one hand, it promotes recovery from CVD, and on the other, it prevents future cardiovascular events [1]. Well-organized CVR, which includes supervised physical training, education, social support, and lifestyle changes, reduces cardiovascular burden by influencing cardiovascular risk factors, improving the quality of life, and reducing mortality and hospital readmission [2–4].

The influence of CVR on hemodynamic status has been a subject of numerous studies. It can reduce heart rate, blood pressure, and rate-blood pressure product values [5, 6]. On the other hand, it can increase arterial compliance, especially in patients with diabetes mellitus [7]. This is why all relevant guidelines consider CVR as an indispensable part of secondary prevention in patients with CVD [8, 9].

Thoracic impedance or impedance cardiography (ICG) is a simple, non-invasive diagnostic method used for assessing hemodynamic status by monitoring electrical conductivity changes of the thorax [10]. It is usually used in patients with heart failure where it can accurately assess stroke volume (SV) and cardiac output (CO) [11–13]. Other indications include pulmonary hypertension, congenital heart diseases, sleep apnea, arterial hypertension, etc. [14–17].

Using ICG, we can assess CO, SV, systemic vascular resistance (SVR), thoracic fluid content, and many other parameters that can help us both with the CVD diagnostic algorithm and the therapeutic approach.

The aim of our study was to evaluate the influence of a three-week CVR program on hemodynamic status in patients with coronary artery disease (CAD). Furthermore, we tried to investigate whether there was a correlation between physical strain tolerance and ICG parameters before and after the CVR program. To our knowledge, this was the first study that has provided this kind of investigation so far.

Methods

Recruitment and patients

The study involved 52 patients referred to the Institute for Treatment and Rehabilitation “Niška Banja” for a three-week CVR program after surviving myocardial infarction (MI), percutaneous coronary intervention (PCI), or coronary artery bypass grafting (CABG). Patients with an artificial valve, implemented permanent pacemaker, reduced ejection fraction (EF < 50%), significant unsolved stenotic lesions on coronary arteries (stenosis ≥ 70%), atrial fibrillation, ventricular arrhythmia, severe anemia, impaired physical or mental conditions were excluded from the study. Data about ejection fraction, valve disease, and other echocardiographic parameters were extracted from previous medical examinations done after MI/PCI/CABG and before study enrollment.

Study design

The three-week CVR program included supervised and personalized physical training that involved cardiovascular exercise training, bicycle riding, and walking. At the beginning and the end of the CVR program, laboratory tests, exercise stress tests (EST), and ICG measurements were taken.

The tests were done on the Treadmill track using the Bruce protocol. After the initial EST, all patients underwent regular aerobic physical training, which included aerobic exercises, bicycle riding, and walking for 45 min *per* session, two sessions on a daily basis. All patients attended physical training five times a week, three weeks in a row. The intensity of physical activity was limited to the submaximal physical capacity determined at the level when 70–80% of the maximum heart rate was reached on the initial EST.

Exercise stress test

ESTs were done on the treadmill (3017 Full Vision Drive, Newton, Kansas, USA) according to the Bruce protocol. Tests were limited by submaximal heart rate (calculated as 85% from the 220-age equation), symptoms like chest pain, dyspnea, fatigue, dizziness, etc., and electropathological changes on electrocardiogram (ECG), which included horizontal or down-sloping ST-segment depression ≥ 1 mm, complexed ventricular arrhythmia that involved couplets of ventricular premature beats or ventricular tachycardia, patient's request to stop the test, and hypertensive reaction which was defined as a sudden increase of systolic BP to values ≥ 220 mmHg, or decreased in systolic BP > 10 mmHg.

Impedance cardiography

Hemodynamic data were obtained noninvasively using CardioScreen® 2000 (Medis GmbH, Germany). All patients underwent ICG monitoring for 15 min. Measurements were done at the same time, approximately half an hour after morning medications and before breakfast. All measurements were done in a supine position after 15 min of rest. Blood pressure was measured every three minutes, and ICG analyzes were taken every five min. The last recorded values were taken for statistical analysis.

Statistical analysis

Statistical analysis was performed using SPSS for Windows (Version 20; SPSS, Chicago, IL, USA). Frequencies and percentages were used for the description of the patient's characteristics. Numerical data were expressed as mean ± standard deviation (SD). The Kolmogorov-Smirnov test was used to test the normality of data. Statistical significance for nominal data was tested with the χ^2 test and, where appropriate, the Fisher exact test. The Student's *t*-test was used to assess the statistical significance of parametric continuous data, and Mann–Whitney *U*-test and Wilcoxon Signed Ranks Test were for nonparametric continuous variables. The Pearson's correlation was used to determine the correlation between variables. Statistical significance was set to a level of $p < 0.05$.

Results

Baseline characteristics

The study included 52 patients with CAD, 42 (80.8%) men and 10 (19.2%) women. The age structure was similar in both genders (male 56.62 ± 9.80 *vs* female 62.30 ± 9.08; $p = 0.101$). Thirty-nine (75%) patients suffered from arterial hypertension, 47 (90.4%) had lipid disorder or took lipid-lowering drugs, and 21 (40.4%) patients suffered from diabetes mellitus. Thirty-nine (75%) patients were smokers or ex-smokers, and 21 (40.4%) had a positive family anamnesis for CVD. Forty-four (84.6) patients survived MI, 35 (67.3%) underwent PCI, and 13 (25%) patients had CABG (Table 1).

Laboratory measurements

At the beginning and the end of CVR, laboratory parameters were measured (Table 2). Cholesterol, low-density lipoprotein (LDL), glycosylated hemoglobin (HbA1c), and creatinine kinase (CK) were lower at the end of CVR compared to the values at the beginning of CVR but without statistical significance. On the other hand, the values of erythrocytes ($t = 3.859$; $p < 0.001$), hemoglobin ($t = 3.245$; $p = 0.002$), and hematocrit ($t = 3.551$; $p = 0.001$) were significantly higher during the second measurement.

Table 1

Baseline characteristics of patients

Parameters	No	Yes
	n (%)	n (%)
Arterial hypertension	13 (25.0)	39 (75.0)
Hyperlipidemia	5 (9.6)	47 (90.4)
Diabetes mellitus	31 (59.6)	21 (40.4)
Smoking habit	13 (25.0)	39 (75.0)
Heredity	21(40.4)	31 (59.6)
Myocardial infarction	8 (15.4)	44 (84.6)
Coronary artery by-pass surgery	39 (75.0)	13 (25.0)
Percutaneous coronary intervention	17 (32.7)	35 (67.3)

Table 2

Laboratory measurements at the beginning and the end of rehabilitation

Parameters	At the beginning (mean ± SD)	At the end (mean ± SD)	t/*Z	*p*
Cholesterol	4.08 ± 0.93	4.01 ± 0.86	0.683	0.498
LDL	2.42 ± 0.74	2.39 ± 0.77	0.250	0.804
HDL	0.99 ± 0.20	0.94 ± 0.18	2.067	0.044
Triglyceride	1.59 ± 0.72	1.59 ± 0.64	0.770	0.939
Glucose	5.97 ± 1.17	6.22 ± 1.67	0.376*	0.707
HbA1c	5.42 ± 1.42	5.28 ± 1.47	1.374*	0.078
CK	157.60 ± 109.27	142.60 ± 78.77	1.167	0.250
CKMB	17.08 ± 8.45	17.87 ± 9.08	0.596*	0.551
Hematocrit	0.41 ± 0.03	0.42 ± 0.04	3.551	0.001
Erythrocytes	4.74 ± 0.44	4.85 ± 0.45	3.859	< 0.001
Hemoglobin	139.02 ± 12.24	141.35 ± 12.36	3.245	0.002
Creatinine	98.22 ± 19.37	98.44 ± 15.81	0.102	0.919
AUR	341.48 ± 106.03	331.14 ± 104.53	1.065	0.294

***Z* – Wilcoxon Signed Ranks Test; LDL – low-density lipoproteins; HDL – high-density lipoproteins; HbA1c – glycosylated hemoglobin; CK – creatinine kinase; MB – myocardial fraction; AUR – acidum uricum; SD – standard deviation.**

Exercise stress test and impedance cardiography

Additionally, all patients underwent EST and ICG measurements at the beginning and the end of the CVR program (Table 3). Patients showed better strain tolerance on the second EST (EST2) by achieving a higher strain level (Z = 2.315; $p = 0.021$) and longer duration of the test (Z = 2.305; $p = 0.021$) compared to the values of the first EST (EST1). Furthermore, a higher number of patients achieved submaximal heart rate values on the EST2 (Z = 3.153; $p = 0.002$).

According to impedance cardiography, CO and SV were higher at the end of CVR, but without any statistical significance ($t = 0.512$; $p = 0.611$ and Z = 1.349; $p = 0.184$). On the other hand, heart rate, systemic vascular resistance index (SVRI), and stroke systemic vascular resistance index (SSVRI) were lower during the second measurement, but also without statistical significance (Z = 1.068; $p = 0.292$, Z = 0.510; $p = 0.613$ and Z = 0.950; $p = 0.348$).

Correlation between exercise stress test and impedance cardiography

There was no significant correlation between the levels of the EST1 and the parameters of the first ICG measurements (ICG1). On the other hand, there was a strong positive correlation between the levels of EST2 and CO ($r = 0.538$; $p < 0.001$) as well as between the levels of EST2 and SV ($r = 0.380$; $p = 0.017$), according to the second ICG measurements (ICG2). Moreover, there was a strong negative correlation between EST2 levels and SVR ($r = -0.472$; $p = 0.002$) as well as between EST2 levels and SVRI ($r = -0.407$; $p = 0.010$), measured by ICG2 (Table 4).

CO significantly positively ($r = 0.357$; $p = 0.026$) and SVR ($r = -0.409$; $p = 0.010$) and SVRI ($r = -0.356$; $p = 0.026$) significantly negatively correlated with EST1 duration when ICG1 parameters were compared. There was a strong positive correlation between EST2 duration and CO ($r = 0.517$; $p = 0.001$) as well as between EST2 duration and

Table 3

Exercise stress test and impedance cardiography

Parameters	At the beginning (mean ± SD)	At the end (mean ± SD)	*t*/*Z	*p*
Exercise stress test				
level	2.93 ± 1.10	3.24 ± 0.87	2.315*	0.021
duration (min)	7.13 ± 3.30	7.79 ± 2.63	2.305*	0.021
SHR (bpm)	0.43 ± 0.50	0.70 ± 0.46	3.153*	0.002
Impedance cardiography				
CO (L/min)	5.85 ± 1.19	5.91 ± 1.32	0.512	0.611
SV (mL)	90.91 ± 22.91	93.63 ± 22.69	1.349	0.184
ACI (1/100/s²)	77.09 ± 26.53	79.21 ± 27.39	0.679	0.490
VI (1/1000/s)	47.07 ± 15.04	47.84 ± 14.19	0.499	0.620
BP systolic (mmHg)	127.77 ± 9.54	128.77 ± 12.94	0.453	0.666
BP diastolic (mmHg)	75.84 ± 9.34	85.79 ± 8.02	0.041	0.968
HR (bpm)	65.14 ± 9.73	63.84 ± 9.27	1.068	0.292
SVR (dyne·s/cm^5)	1237.93 ± 275.98	1238.40 ± 311.27	0.004	0.997
SVRI (dyne·s/cm^5/m²)	2397.40 ± 450.33	2368.81 ± 502.97	0.510	0.613
SSVRI (dyne·s/cm^5/m²)	156.95 ± 34.58	152.91 ± 36.40	0.950	0.348

***Z* – Wilcoxon Signed Ranks Test; SHR – submaximal heart rate; CO – cardiac output; SV – stroke volume; ACI – acceleration index; VI – velocity index; BP – blood pressure; HR – heart rate; SVR – systemic vascular resistance; SVRI – SVR index; SSVRI – stroke SVRI; SD – standard deviation.**

Table 4

Correlation between the exercise stress test (EST) level and impedance cardiography (ICG) parameters

ICG parameters	EST1 level		EST2 level	
	r	*p*	*r*	*p*
CO (L/min)	0.266	0.102	0.538	< 0.001
SV (mL)	0.177	0.280	0.380	0.017
ACI (1/100/s²)	0.075	0.652	0.265	0.104
VI (1/1000/s)	0.172	0.295	0.284	0.079
BP systolic (mmHg)	0.078	0.635	0.128	0.438
BP diastolic (mmHg)	0.082	0.619	0.008	0.964
HR (bpm)	0.155	0.345	0.187	0.255
SVR (dyne·s/cm^5)	-0.209	0.201	-0.472	0.002
SVRI (dyne·s/cm^5/m²)	-0.163	0.322	-0.407	0.010
SSVRI (dyne·s/cm^5/m²)	-0.031	0.850	-0.236	0.148

For abbreviations see under Table 3.

SV (r = 0.340; p = 0.034), and a strong negative correlation between EST2 duration and SVR (r = -0.504; p = 0.001) as well as between EST2 duration and SVRI (r = -0.448; p = 0.004) measured by ICG2 (Table 5).

Table 5

Correlation between exercise stress test (EST) duration and impedance cardiography (ICG) parameters

ICG parameters	EST1 duration (min)		EST2 duration (min)	
	r	*p*	*r*	*p*
CO (L/min)	0.357	0.026	0.517	0.001
SV (mL)	0.246	0.131	0.340	0.034
ACI (1/100/s²)	0170	0.302	0.226	0.167
VI (1/1000/s)	0.257	0.115	0.264	0.104
BP systolic (mmHg)	-0.075	0.649	0.061	0.714
BP diastolic (mmHg)	-0.040	0.807	-0.056	0.737
HR (bpm)	0.161	0.328	0.225	0.169
SVR (dyne·s/cm^5)	-0.409	0.010	-0.504	0.001
SVRI (dyne·s/cm^5/m²)	-0.356	0.026	-0.448	0.004
SSVRI (dyne·s/cm^5/m²)	-0.182	0.267	-0.246	0.131

For abbreviations see under Table 3.

Discussion

The beneficial effects of CVR are proven in patients with coronary artery disease, severe arterial hypertension, heart failure, heart valve repair, and heart transplantation [18]. A well-designed CVR program reduces cardiovascular mortality and hospital readmission and increases the quality of life in CVD patients [19–21]. This is why relevant Guidelines give recommendations for CVR in patients with CVD [8, 22–27]. Three main postulates of CVR are exercise training, lifestyle changes, and psychological intervention [18]. The main goal of CVR is "to provide the best possible physical, mental, and social conditions, so that the patients may, by their efforts, preserve or resume optimal functioning in their community and through improved health behavior slow or reverse the progression of the disease" [28].

Physical activity as a cornerstone of CVR has a positive effect on CVD regardless of patients' reduced or preserved ejection fraction [8, 29]. In addition, the beneficial effects of exercise-based CVR do not depend on gender or age structure [30, 31]. It increases physical strain tolerance [30] and arterial compliance [7], improves lipid profile [32, 33], reduces the symptoms of depression [34–36], and leads to weight and blood pressure reduction [37, 38]. It can even induce gray matter volume recovery [39]. Our study once again showed that personalized and supervised physical training improves physical strain tolerance. The average strain level was higher, and the duration of the test was significantly longer in the second EST. Additionally, a higher number of patients achieved submaximal heart rate on EST2 compared to EST1.

On the other hand, CVR did not lead to significant changes in lipid profile or any other laboratory parameters except red blood count. It did reduce total cholesterol, LDL, and HbA1c, but without statistical significance. This is probably due to the shorter duration of the CVR program, as our study included patients with three-week CVR. For significant laboratory changes, CVR should last longer, at least eight weeks [31, 32, 40]. For example, in a study with 547 coronary heart disaese (CHD) patients, CVR led to a better lipid profile (total cholesterol, triglycerides, LDL, and HDL). However, the CVR program in this study included 24 exercise sessions scheduled over eight weeks [41]. Furthermore, Vergès et al. [42] showed in their research that CVR could improve response to the hypolipidemic therapy in patients with CHD. However, the average duration of CVR in this study was two months. On the other hand, we noted an improvement in red blood count which can have a positive impact on functional capacity [43].

By using ICG, we tried to investigate the effects of CVR on hemodynamic status in CHD patients. Besides CO and SV, we assessed contractility-related (VI, ACI) and vascular system-related parameters (SVR, SVRI, SSVRI). It is well known that physical activity can decrease blood pressure values through the reduction of vascular resistance [44]. The exact mechanism of how physical activity leads to vascular resistance reduction is unknown, but it probably includes the sympathetic nervous system and renin-angiotensin-aldosterone system [45]. Furthermore, a combination of endurance and resistance training can improve endothelial function and arterial compliance [46, 47]. In our study, exercise-based CVR did lead to SVRI and SSVRI reduction, but without statistical significance (Z = 0.510; p = 0.613 and Z = 0.950; p = 0.348). However, we emphasize the fact that the beneficial effects of exercise training are dose-dependent. It would be interesting to see whether the reduction of vascular-related parameters would be statistically significant if the CVR program lasted eight or twelve weeks.

Beneficial but temporary results of CVR on hemodynamic status are found in patients with stable chronic heart failure [48]. Using ICG, Gielerak et al. [49] showed that CVR could reduce fluid retention in patients with HF. Moreover, exercise training can improve SV and CO [50]. However, in our study, CO and SV were higher at the end of CVR, but without any statistical significance (t = 0.512; p = 0.611 and Z = 1.349; p = 0.184). Other studies that investigated the influence of physical training on SV or CO included patients with reduced ejection fraction or patients with heart failure with preserved EF (HFpEF). Our study included stable patients with preserved EF and without any symptoms and signs of HF.

The next step in our investigation was to assess whether there was a correlation between the parameters of the exercise stress test and ICG. There was no significant correlation between the average level of EST1 and ICG1 parameters. On the other hand, there was a strong positive correlation between the level of EST2 and CO ($r = 0.538$; $p < 0.001$) and between the level of EST2 and SV ($r = 0.380$; $p = 0.017$) according to ICG2. Moreover, there was a strong negative correlation between the EST2 level and SVR ($r = -0.472$; $p = 0.002$) as well as between the EST2 level and SVRI ($r = -0.407$; $p = 0.010$) according to ICG2. CO significantly positively ($r = 0.357$; $p = 0.026$) and SVR ($r = -0.409$; $p = 0.010$) and SVRI ($r = -0.356$; $p = 0.026$) significantly negatively correlated with EST1 duration when ICG1 parameters were compared. There was a strong positive correlation between EST2 duration and CO ($r = 0.517$; $p = 0.001$) as well as between EST2 duration and SV ($r = 0.340$; $p = 0.034$), and a strong negative correlation between EST2 duration and SVR ($r = -0.504$; $p = 0.001$) as well as between EST2 duration and SVRI ($r = -0.448$; $p = 0.004$) according to ICG2.

These results show that CVR has a great influence on hemodynamic parameters. Namely, CVR led to a positive correlation between physical strain tolerance and cardiac and stroke volume and a better negative correlation between physical strain tolerance and peripheral vascular resistance. As far as we know, this is the first study that investigated the effects of CVR on hemodynamic parameters and its correlation with physical strain tolerance in patients with CAD and preserved ejection fraction. An implementation of ICG as a valuable tool in assessing the effects of CVR on the hemodynamic status of CAD patients should be considered.

Study limitations

There are several limitations to this study: 1) the study sample was relatively small. However, almost all studies that involved ICG as a diagnostic tool included 50 patients or less; 2) we did not perform echocardiographic examinations. It would be interesting to see how some ICG parameters (*i.e.*, SV and CO) correlate with some echocardiographic parameters (*i.e.*, left ventricular mass index and cardiac index); 3) our study included the CVR program that lasted for three weeks. We consider that a longer CVR duration (*i.e.*, eight weeks) could have greater beneficial effects on the hemodynamic status of CHD patients.

Conclusion

Our study once again showed that a well-designed CVR program could lead to better physical strain tolerance in patients with CAD. Furthermore, CVR led to a significant positive correlation between EST and cardiac output as well as between EST and stroke volume measured by ICG. On the other hand, there was a significant negative correlation between EST and vascular-related parameters according to ICG at the end of the CVR program.

Conflict of interests

The authors have no conflict of interest to declare regarding the present paper.

R E F E R E N C E S

1. *Kachur S, Chongthammakun V, Lavie CJ, De Schutter A, Arena R, Milani RV,* et al. Impact of cardiac rehabilitation and exercise training programs in coronary heart disease. Prog Cardiovasc Dis 2017; 60(1): 103–14.
2. *Choo CC, Chew PK, Lai SM, Soo SC, Ho CS, Ho RC*, et al. Effect of cardiac rehabilitation on quality of life, depression and anxiety in Asian patients. Int J Environ Res Public Health 2018; 15(6): 1095.
3. *Lavie CJ, Menezes AR, De Schutter A, Milani RV, Blumenthal JA.* Impact of cardiac rehabilitation and exercise training on psychological risk factors and subsequent prognosis in patients with cardiovascular disease. Can J Cardiol 2016; 32(10 Suppl): S365–73.
4. *Smart NA.* Benefit of in-hospital cardiac rehabilitation on mortality and readmissions in heart failure. Eur J Prev Cardiol 2019; 26(8): 806–7.
5. *Ghashghaei FE, Sadeghi M, Marandi SM, Ghashghaei SE.* Exercise-based cardiac rehabilitation improves hemodynamic responses after coronary artery bypass graft surgery. ARYA Atheroscler 2012; 7(4): 151–6.
6. *Neves A, Alves AJ, Ribeiro F, Gomes JL, Oliveira J.* The effect of cardiac rehabilitation with relaxation therapy on psychological, hemodynamic, and hospital admission outcome variables. J Cardiopulm Rehabil Prev 2009; 29(5): 304–9.
7. *Mourot L, Boussuges A, Campo P, Maunier S, Debussche X, Blanc P.* Cardiovascular rehabilitation increase arterial compliance in type 2 diabetic patients with coronary artery disease. Diabetes Res Clin Pract 2006; 84(2): 138–44.
8. *Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL,* et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). Eur Heart J 2016; 37(29): 2315–81.
9. *Herdy AH, López-Jiménez F, Terzic CP, Milani M, Stein R, Carvalho T,* et al. South American guidelines for cardiovascular disease prevention and rehabilitation. Arq Bras Cardiol 2014; 103(2 Suppl 1): 1–31. (England, Portuguese)
10. *Chabchoub S, Mansouri S, Salah RB.* Determination of Arterial Compliance Using Thoracic Bioimpedance to Predict the Presence of Heart Diseases. Int J Cardiovasc Pract 2018; 3(2): 25–9.
11. *Sadauskas S, Naudžiūnas A, Unikauskas A, Mašanauskienė E, Bakšytė G, Macas A.* Applicability of Impedance Cardiography During Heart Failure Flare-Ups. Med Sci Monit 2016; 22: 3614–22.
12. *Facchini C, Malfatto G, Giglio A, Facchini M, Parati G, Branzi G.* Lung ultrasound and transthoracic impedance for noninvasive evaluation of pulmonary congestion in heart failure. J Cardiovasc Med (Hagerstown) 2016; 17(7): 510–7.

13. *Yancy C, Abraham WT*. Noninvasive hemodynamic monitoring in heart failure: utilization of impedance cardiography. Congest Heart Fail 2003; 9(5): 241–50.
14. *Dupuis M, Noel-Savina E, Prévot G, Tétu L, Pillard F, Rivière D,* et al. Determination of cardiac output in pulmonary hypertension using impedance cardiography. Respiration 2018; 96(6): 500–6.
15. *Ebrahim M, Hegde S, Printz B, Abcede M, Proudfoot JA, Davis C*. Evaluation of impedance cardiography for measurement of stroke volume in congenital heart disease. Pediatr Cardiol 2016; 37(8): 1453–7.
16. *Yasuda Y, Umezu A, Horihata S, Yamamoto K, Miki R, Koike S*. Modified thoracic impedance plethysmography to monitor sleep apnea syndromes. Sleep Med 2005; 6(3): 215–24.
17. *Taler SJ, Textor SC, Augustine JE*. Resistant hypertension: comparing hemodynamic management to specialist care. Hypertension 2002; 39(5): 982–8.
18. *Bellmann, B, Lin T, Greissinger K, Rottner L, Rillig A, Zimmerling S*. The Beneficial Effects of Cardiac Rehabilitation. Cardiol Ther 2020; 9(1): 35–44.
19. *Anderson L, Oldridge N, Thompson DR, Zwisler AD, Rees K, Martin N*, et al. Exercise-Based Cardiac Rehabilitation for Coronary Heart Disease: Cochrane Systematic Review and Meta-Analysis. J Am Coll Cardiol 2016; 67(1): 1–12.
20. *Goel K, Pack QR, Lahr B, Greason KL, Lopez-Jimenez F, Squires RW,* et al. Cardiac rehabilitation is associated with reduced long-term mortality in patients undergoing combined heart valve and CABG surgery. Eur J Prev Cardiol 2015; 22(2): 159–68.
21. *Heran BS, Chen JM, Ebrahim S, Moxham T, Oldridge N, Rees K,* et al. Exercise-based cardiac rehabilitation for coronary heart disease. Cochrane Database Syst Rev 2011; (7): CD001800.
22. Surveillance report 2017 – Myocardial infarction: cardiac rehabilitation and prevention of further cardiovascular disease (2013) NICE guideline CG172 [Internet]. London: National Institute for Health and Care Excellence (UK); 2017.
23. *Gimigliano F, Negrini S*. The World Health Organization "Rehabilitation 2030: a call for action". Eur J Phys Rehabil Med 2017; 53(2): 155–68.
24. *Smith SC Jr, Benjamin EJ, Bonow RO, Braun LT, Creager MA, Franklin BA,* et al. AHA/ACCF secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation endorsed by the World Heart Federation and the Preventive Cardiovascular Nurses Association. J Am Coll Cardiol 2011; 58(23): 2432–46.
25. *Chew DP, Scott IA, Cullen L, French JK, Briffa TG, Tideman PA,* et al. NHFA/CSANZ ACS Guideline 2016 Executive Working Group: National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand: Australian Clinical Guidelines for the Management of Acute Coronary Syndromes 2016. Heart Lung Circ 2016; 25(9): 895–951.
26. *Amsterdam EA, Wenger NK, Brindis RG, Casey DE Jr, Ganiats TG, Holmes DR Jr,* et al. 2014 AHA/ACC Guideline for the Management of Patients with Non-ST-Elevation Acute Coronary Syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2014; 64(24): e139–e228.
27. *Price KJ, Gordon BA, Bird SR, Benson AC*. A review of guidelines for cardiac rehabilitation exercise programs: is there an international consensus? Eur J Prevent Cardiol 2016; 23(16): 1715–33.
28. British Association for Cardiovascular Prevention and Rehabilitation. The BACPR standards and core components for cardiovascular disease prevention and rehabilitation 2017. 3rd ed. London: BACPR, 2017.
29. *Taylor RS, Long L, Mordi IR, Madsen MT, Davies EJ, Dalal H,* et al. Exercise-based rehabilitation for heart failure: Cochrane systematic review, meta-analysis, and trial sequential analysis. JACC Heart Fail 2019; 7(8): 691–705.
30. *Deljanin-Ilić M, Stojanović M, Ilić S*. The effect of cardiovascular rehabilitation on physical strain tolerance-does gender really matter? Vojnosanit Pregl 2019; DOI: https://doi.org/10.2298/VSP190727146D.
31. *Lavie CJ, Milani RV, Littman AB*. Benefits of cardiac rehabilitation and exercise training in secondary coronary prevention in the elderly. J Am Coll Cardiol 1993; 22(3): 678–83.
32. *Mouine N, Amah G, Guiti C, Gagey S, Duval M, Abdennbi K*. Impact of cardiac rehabilitation in LDL cholesterol level. Arch Cardiovasc Dis Suppl 2020; 12(1): 152.
33. *Vasilcu TF, Statescu C, Sascau R, Roca M, Costea CF, Zota M,* et al. Cardiopulmonary testing and biochemical profile of coronary patients subject to cardiovascular recovery programs. Parameters 2018; 10: 11.
34. *Blumenthal JA, Babyak MA, O'Connor C, Keteyian S, Landzberg J, Howlett J,* et al. Effects of exercise training on depressive symptoms in patients with chronic heart failure: the HF-ACTION randomized trial. JAMA 2012; 308(5): 465–74.
35. *Sharif F, Shoul A, Janati M, Kojuri J, Zare N*. The effect of cardiac rehabilitation on anxiety and depression in patients undergoing cardiac bypass graft surgery in Iran. BMC Cardiovasc Disord 2012; 12: 40.
36. *Ernstsen L, Rangul V, Nauman J, Nes BM, Dalen H, Krokstad S,* et al. Protective Effect of Regular Physical Activity on Depression After Myocardial Infarction: The HUNT Study. Am J Med 2016; 129(1): 82–8.e1.
37. *AHA Writing Group: Balady GJ, Fletcher BJ, Froelicher ES, Hartley LH, Krauss RM, Oberman A*, et al. Cardiac rehabilitation programs: a statement for health care professionals from the American Heart Association. Circulation 1994; 90(3): 1602–10.
38. *Taylor C, Houston Miller N, Ahn D, Haskell W, DeBusk RF*. The effect of exercise training programs on psychological improvement in uncomplicated postmyocardial infarction patients. J Psychosom Res 1986; 30(5): 581–7.
39. *Anazodo UC, Shoemaker JK, Suskin N, St Lawrence KS*. An investigation of changes in regional gray matter volume in cardiovascular disease patients, pre and post cardiovascular rehabilitation. Neuroimage Clin 2013; 3: 388–95.
40. *Jørgensen MM, Petersen AK, Nielsen MM, Petersen MS, Møller BK, Møller K,* et al. Exercise-based cardiac rehabilitation reduces key inflammatory biomarkers in atherosclerosis: a dose response study. Ann Atheroscler Res 2018; 1(3): 1012.
41. *Sarrafzadegan N, Rabiei K, Kabir A, Asgary S, Tavassoli A, Khosravi A*, et al. Changes in lipid profile of patients referred to a cardiac rehabilitation program. Eur J Cardiovasc Prev Rehabil 2008; 15(4): 467–72.
42. *Vergès BL, Patois-Vergès B, Cohen M, Casillas JM*. Comprehensive cardiac rehabilitation improves the control of dyslipidemia in secondary prevention. J Cardiopulm Rehabil 1998; 18(6): 408–15.
43. *Church TS, Lavie CJ, Milani RV, Kirby GS*. Improvements in blood rheology after cardiac rehabilitation and exercise training in patients with coronary heart disease. Am Heart J 2002; 143(2): 349–55.
44. *Cornelissen VA, Fagard RH*. Effects of endurance training on blood pressure, blood pressure–regulating mechanisms, and cardiovascular risk factors. Hypertension 2005; 46(4): 667–75.
45. *Fagard RH*. Exercise is good for your blood pressure: effects of endurance training and resistance training. Clin Exp Pharmacol Physiol 2006; 33(9): 853–6.
46. *Umpierre D, Stein R*. Hemodynamic and vascular effects of resistance training: implications for cardiovascular disease. Arq Bras Cardiol 2007; 89(4): 256–62.

47. *Antunes BM, Rossi FE, Cholewa JM, Lira FS*. Regular Physical Activity and Vascular Aging. Curr Pharm Des 2016; 22(24): 3715–29.
48. *Kayanakis JG, Page E, Aros F, Borau F*. Rehabilitation of patients with chronic cardiac insufficiency. Immediate and midterm effects. Presse Med 1994; 23(3): 121–6. (French)
49. *Gielerak G, Piotrowicz E, Krzesiński P, Kowal J, Grzęda M, Piotrowicz R*. The effects of cardiac rehabilitation on haemodynamic parameters measured by impedance cardiography in patients with heart failure. Kardiol Pol 2011; 69(4): 309–17.
50. *Motohiro M, Yuasa F, Hattori T, Sumimoto T, Takeuchi M, Kaida M,* et al. Cardiovascular adaptations to exercise training after uncomplicated acute myocardial infarction. Am J Phys Med Rehabil 2006; 84(9): 684–91.

Received on August 10, 2020
Revised on November 9, 2020
Accepted on November 19, 2020
Online First December 2020