UDK 612.33:579.8]:612.11/.17
COBISS.SR-ID 38711049

GASTROENTEROKARDIOLOGIJA - ILI ŠTA IMAJU ZAJEDNIČKO CREVO I SRCE?

Zoran Joksimović (1), Dušan Bastać (2), Snežana Pavlović (3)

(1) INTERNISTIČKA ORDINACIJA „JOKSIMOVIĆ“ BOR; (2) INTERNISTIČKA ORDINACIJA „DR BASTAĆ“ ZAJEČAR; (3) SPECIJALISTIČKA ORDINACIJA ZA INTERNU MEDICINU „DR PAVLOVIĆ KARDIOLOGIJA“ BEOGRAD

Sažetak: Crevna mikrobiota našeg organizma je zajednica bakterija, arheja, gljivica, virusa i parazita koji čine jedinstveni ekosistem u digestivnom traktu koji se sastoji od oko 10^{14} mikroorganizama. Raznolikost ove zajednice kod pojedinih osoba nastaje zbog razlika u genomu domaćina i uticaja faktora životne sredine, uključujući higijenu, ishranu, način života i upotrebu različitih lekova. Značajan broj dokaza sugerše da bi promene u mikrobioti mogle da igraju ulogu u kardiovaskularnim bolestima. Rezultati istraživačkih radova tokom poslednje dve decenije potvrdili su da izmenjeni sastav mikrobiote (disbioza) doprinosi razvoju različitih bolesti, uključujući kardiovaskularne bolesti, dijabetes tipa 2, hronične bolesti bubrega, nealkoholnu masnu jetru, hronične upalne bolesti creva i čak određene vrste karcinoma. Sve je više dokaza da ubuduće, pored dosadašnjih faktora predispozicije za kardiovaskularne i metaboličke bolesti, uključujući genetske faktore, faktore životne sredine i životnog stila, treba računati i na nove faktore rizika poput nutritivne disproporcije i disbioze creva. Tako „novim očima“ gledamo na vezu između gastrointestinalnog trakta i kardiovaskularnog sistema odnosno na „osovinu crevo-srce“.

Ključne reči: crevna mikrobiota, nutritivna disproporcija, disbioza, kardiovaskularne bolesti, metabolički poremećaji

Uvod

Proučavanje crevne mikrobiote čoveka i njegove uloge u raznim bolestima značajno je napredovalo u poslednjoj deceniji. Mikrobiotu čoveka čine svi mikroorganizmi koji žive u simbiozi s ljudskim telom, a mikrobiom predstavlja skup svih gena mikrobiote.

Humani mikrobiom sadrži i do stotinu puta veći broj gena negoli ljudski genom. Mikroorganizme možemo naći u raznim delovima ljudskog tela, ali najbrojnija je mikrobiota organa za varenje. Mikrobiota je varijabilna među zdravim ljudima i jedinstvena za pojedinca - ne postoje dva ista mikrobioma kod ljudi, kao što ne postoje ni isti otisci prstiju. Iako ne postoje dva čoveka sa istim sastavom mikrobiote, ipak, postoje veće sličnosti u sastavu mikrobiote među pripadnicima istih rasa, etničkih grupa i osoba u krvnom srodstvu. Crevna mikrobiota je skup od približno 10^{14} mikroorganizama. Broj bakterija u svakom čoveku je veći od broja ljudi koji su ikada živeli na zemlji. To je zajednica desetostruko brojnija od svih ćelija našeg organizma. Sastoji se od bakterija (oko 1000 različitih vrsta), arheja, gljivica, virusa i parazita koji čine jedinstveni ekosistem. Mikrobiota ima veoma važnu ulogu

za čovekovo zdravlje. Od ključne je važnosti za održavanje homeostatskih funkcija gastrointestinalnog trakta, jer učestvuje u procesima varenja domaćina, metabolizmu i regulaciji crevnog imunog sistema [1,2]. Po rođenju, digestivni trakt novorođenčeta nije naseljen mikroorganizmima. U prvim satima života ga kolonizuju majčini mikroorganizmi, u početku koliformne bakterije i streptokoke, kasnije laktobacili i enterokoke, a broj mikroorganizama u crevnom traktu počinje da se povećava, postepeno formirajući dinamičku ravnotežu crevne mikrobiote. Naravno, porast ovih bakterija zavisi i od načina porođaja - prirodni ili carskim rezom [3]. U odrasloj dobi, većina crevne mikrobiote sastoji se od pet grupa mikroba, a to su: *Bacteroidetes*, *Firmicutes*, *Actinobacteria*, *Proteobacteria* i *Cerrucomicrobia*. Približno je proporcionalni broj Gram-pozitivnih *Firmicutes* i Gram-negativnih *Bacteroidetes* koje čine većinu vrsta u zdravom odraslom crevu > 90% [4-5]. Odnos *Firmicutes* i *Bacteroidetes* ostaje relativno konstantna kod zdrave jedinice, iako nije jednak kod svih pojedinaca. Razlike nastaju zbog razlika u genomima domaćina, faktorima okoline kao što su higijena, prehrana, način života i upotreba antibiotika [4]. Zbog kiselog okruženja i intenzivne peristaltike, u

GASTROENTEROCARDIOLOGY-OR WHAT DO THE GUT AND THE HEART HAVE IN COMMON?

Zoran Joksimović (1), Dušan Bastać (2), Snežana Pavlović (3)

(1) INTERNISTIČKA ORDINACIJA „JOKSIMOVIĆ“ BOR; (2) INTERNISTIČKA ORDINACIJA „DR BASTAĆ“ ZAJEČAR; (3) SPECIJALISTIČKA ORDINACIJA ZA INTERNU MEDICINU „DR PAVLOVIĆ KARDIOLOGIJA“ BEOGRAD

Summary: The gut microbiota of our organism is a community of bacteria, archaea, fungi, viruses and parasites that make up a unique ecosystem in the digestive tract, which consists of about 1014 microorganisms. The diversity of this community between individuals occurs because of the differences in the host genome and the impact of environmental factors, including hygiene, diet, lifestyle and the use of different drugs. Significant evidence suggests that changes in the microbiota could play a role in cardiovascular diseases. The results of research papers for the last two decades have confirmed that altered gut microbiota composition (dysbiosis) contributes to the development of various diseases, including cardiovascular diseases, type 2 diabetes, chronic kidney disease, nonalcoholic fatty liver disease, chronic inflammatory bowel disease and even certain types of cancer. There is growing evidence that in the future, apart from current predisposing factors for cardiovascular and metabolic diseases, including genetic, environmental and lifestyle factors, one should count on new risk factors such as nutritional disproportion and gut dysbiosis. Thus, we look upon the relationship between the gastrointestinal tract and cardiovascular system, i.e. the "gut-heart axis" in a new way.

Key words: intestinal microbiota, nutritional disproportion, dysbiosis, cardiovascular diseases, metabolic disorders

Introduction

The study of the human gut microbiota and its role in various diseases has advanced significantly in the last decade. The human microbiota consists of all microorganisms that live in a symbiosis with the human body, while the microbiome represents the sum of all genes of the microbiota.

The human microbiome contains up to 100 times more genes than the human genome. Microorganisms can be found in various parts of the human body, but the most numerous is the microbiota of the digestive organs. The microbiota is variable among healthy people but it is also unique for an individual – no two microbiomes are the same in humans, just as there are no the same fingerprints. Although there are no two people with the same composition of the microbiota, however, there are larger similarities in the composition of microbiota among the members of the same races, ethnic groups and blood relatives. Gut microbiota is a collection of approximately 1014 microorganisms. The number of bacteria in every human being is larger than the number of

people who have ever lived on earth. It is a community ten times more numerous than all the cells of our organism. It consists of bacteria (about 1000 different species), archaea, fungi, viruses and parasites that make up a unique ecosystem. The microbiota plays a very important role in human health. It is of utmost importance for maintaining the homeostatic functions of the gastrointestinal tract, as it participates in digestion processes of the host, metabolism and regulation of the gut immune system. [1,2]. After birth, the digestive tract of the newborn is not inhabited by microorganisms. In the first hours of life, it is colonized by maternal microorganisms, initially coliform bacteria and streptococci, later lactobacilli and enterococci, and the number of microorganisms in the gut tract begins to increase, gradually forming a dynamic balance of the gut microbiota. Of course, the growth of these bacteria also depends on the method of birth - natural or by caesarean section [3]. In adulthood, most of the gut microbiota consists of five groups of microbes, namely: Bacteroidetes, Firmicutes, Actinobacteria, Proteobacteria, and Verrucomicrobia. The number of Gram-

želucu i dvanaestopalačnom crevu je prisutno manje mikroorganizama (10–1000/ml), od kojih su većina gram-pozitivne bakterije. Enterokoke i Laktobacili su prisutni u dvanaestopalačnom crevu, a broj bakterija u ovom području je obično 10^4 /ml. Brojem i raznolikošću vrsta najbogatije je debelo crevo (10^{12} /ml), koje je pretežno naseljeno gram-negativnim i anaerobnim bakterijama [5].

Šta je tačno zdrava mikrobiota nije do kraja razjašnjeno, ali je pokazano da u slučaju narušene ravnoteže - disbioze, može doći do razvoja bolesti. Kada prehrambene navike, faktori okoline, crevna infekcija, neki lekovi ili drugi faktori dovedu do promena u vrsti i količini crevnih mikroorganizama, dolazi do

disbioze creva, koja uzrokuje zapaljenjske i metaboličke poremećaje. Homeostaza crevne mikrobiote presudna je za održavanje ljudskog zdravlja, dok disbioza doprinosi razvoju raznih bolesti, uključujući kardiovaskularne, hronične bolesti bubrega, dijabetes tipa 2, nealkoholnu masnu jetru, pa i neke vrste raka [1,6,7]. Disbioza creva može objasniti zašto su neki pojedinci skloniji razvoju određenih bolesti. Promene u sastavu mikrobiote nedavno su identifikovane kao važan faktor narušavanja „osovine crevo-srce“ što doprinosi razvoju ateroskleroze i hipertenzije - dva glavna faktora rizika za razvoj kardiovaskularnih bolesti [1,7,8].

slika 1. Uticaj Disbioze creva na kardiovaskularne bolesti

(preuzeta sa sajta [https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964\(20\)30024-4/fulltext](https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964(20)30024-4/fulltext) u naučne svrhe a ne koristi se u komercijalne svrhe)

Mikrobiota creva se može smatrati endokrinim organom, a svaki mikrob ima sposobnost da proizvede stotine različitih poznatih i nepoznatih metabolita koji deluju dalje od samog creva. Kako većinu bakterija koje nastanjuju organe za varenje za sada nije moguće kolonizovati u laboratoriji, za utvrđivanje sastava mikrobiote koriste se metoda sekvenciranja nove generacije i bioinformatičke analize ekstrahovane mikrobne dezoksiribonukleinske kiseline (DNK).

Poslednjih godina proučavan je uticaj sastava mikrobiote na različite hronične i autoimune bolesti, posebno u ispitivanjima na životinjama [3,8]. Ova istraživanja pokazuju značaj mikrobiote u odnosu na zdravlje i imunitet i nude nove, još neotkrivene mogućnosti za upotrebu ovog znanja u lečenju nekih drugih bolesti, poput metaboličkog sindroma, insulinske rezistencije, hroničnih inflamatornih bolesti creva, karcinoma [1,6,9].

positiveFirmicutesand Gram-negative Bacteroidetes that make up the most species in a healthy adult gut> 90% (4-5) is approximately proportional.The ratio between Firmicutes and Bacteroidetes remains relatively constant in a healthy individual although it is not the same in all individuals. The differences occur because of the difference in the host genomes, environmental factors as hygiene, diet, lifestyle and use of antibiotics [4]. Due to the acidic environment and intense peristalsis, fewer microorganisms (10–1000 / ml),most of which are Gram-positive bacteria, are present in the stomach and duodenum. Enterococci and Lactobacilli are present in the duodenum, and the number of bacteria in this area is usually 104 / ml. The colon, which is predominantly inhabited by gram-negative and anaerobic bacteria, is the richest in the number and variety of species (1012/ ml) [5].

It hasn't been fully clarified what a healthy microbiota actually is, but it has been

shown that in the case of disturbed balance - dysbiosis, a disease can develop. When eating habits, environmental factors, gut infection, some drugs or other factors lead to changes in the type and amount of gut microorganisms, there occurs gut dysbiosis, which causes inflammatory and metabolic disorders. Homeostasis of the gut microbiota is crucial for maintaining health in human beings, while dysbiosis contributes to the development of various diseases, including cardiovascular, chronic kidney disease, type 2 diabetes, nonalcoholic fatty liver, and even some types of cancer [1,6,7]. Gut dysbiosis may explain why some individuals are more prone to developing certain diseases. Changes in the composition of the microbiota have recently been identified as an important factor in the dysfunction of the "gut-heart axis", which contributes to the development of atherosclerosis and hypertension - two main risk factors for the development of cardiovascular diseases [1,7,8].

Picture 1. Influence of Intestinal Dysbiosis on Cardiovascular Diseases (taken from [https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964\(20\)30024-4/fulltext](https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964(20)30024-4/fulltext) for scientific purposes and not used for commercial purposes)

The gut microbiota can be considered an endocrine organ, and each microbe has the ability to produce hundreds of different known and unknown metabolites that act beyond the gut itself. As most of the bacteria that inhabit the digestive organs cannot be colonized in the

laboratory at the moment, for the purpose of determining the composition of the microbiota next-generation sequencing method and bioinformatics analysis of extracted microbial deoxyribonucleic acid (DNA) are used. In recent years, the impact of the

Disbioza creva i ateroskleroza

Ateroskleroza je glavni faktor rizika za kardiovaskularne bolesti. Ovaj proces karakteriše akumulacija holesterola i makrofaga (inflamatornih ćelija) u vaskularnim zidovima, što doprinosi stvaranju aterosklerotičkih plakova. Nedavne studije pokazale su da crevna disbioza može da doprinese razvoju ateroskleroze modulirajući zapaljenske procese i formirajući određene mikrobne metabolite [10-15].

Integritet crevne sluznice je prva barijera koja štiti domaćina od upada patogena, prolaska crevnog sadržaja i bakterijskih komponenata u krvne sudove. Smanjena koncentracija proteina koji osiguravaju bliske ćelijske kontakte i njihovu nepropusnost, uključujući zonula occludens-1 (ZO-1) poznat i kao Tight junction protein-1 (TJP1), kladin 1 i okcludin, omogućava povećanu propustljivost crevnog zida remeteći ravnotežu između odumiranja i regeneracije mukoznih ćelija [1,13,14]. Ako je sluzna barijera oštećena, prodor mikroba i njihovih produkata u krvne sudove pokreće imuni odgovor, tkivno i sistemsko zapaljenje. Oštećenje integriteta crevne barijere izazvano crevnom disbiozom, dakle, deluje kao faktor rizika koji pokreće hronično zapaljenje koje leži u osnovi različitih bolesti uključujući i aterosklerozu. Glavni molekuli-produkti bakterija koji su i pokretači imunog i zapaljenjskog odgovora su „patogeni povezani sa molekularnim obrascima“ (engl. Pathogen-associated molecular pattern- PAMP). PAMP-ovi aktiviraju urođeni imunološki odgovor, štiteći domaćina od infekcije. Širok spektar različitih vrsta molekula može da služi kao PAMP, uključujući glikane i glikokonjugate. Bakterijski lipopolisaharidi (LPS), endotoksini koji se nalaze na ćelijskim membranama gram negativnih bakterija, smatraju se prototipskom klasom PAMP-a. Povezanost nivoa LPS u plazmi i kardiovaskularnog rizika prvi put su proučavali 1999. godine Niebauer i saradnici [15]. Rezultati studije potvrdili su da je nivo endotoksemije bio najveći kod pacijenata sa najtežom kardiovaskularnom bolešću. Čani i saradnici potvrdili su u svojoj studiji da je crevna disbioza sprečila stvaranje „proteina bliskog kontakta“, što je rezultiralo povećanjem propustljivosti sluzokože creva, a samim tim i prolazom LPS u krv [16]. LPS, koji se proizvode u povećanim količinama u crevnoj disbiozi, mogu igrati važnu ulogu u modulaciji „receptora sličnih putarini“ (Toll-like receptors-TLR) koji prepoznaju

bakterijske proizvode i regulišu imunološki sistem domaćina. TLR su klasa proteina koji igraju ključnu ulogu u urođenom imunološkom sistemu. Oni su jednokratno prolazni membranski receptori koji se obično nalaze na sentinel ćelijama (ćelije prve linije odbrane) poput makrofaga i dendritičnih ćelija koje prepoznaju strukturno očuvane molekule izvedene iz mikroba. Jednom kada mikrobi probiju fizičke barijere poput kože ili sluznice crevnog trakta, prepoznaju ih TLR-ovi, koji aktiviraju reakcije imunih ćelija. Kliničke studije su pokazale da je porast TLR povezan sa antiinflamatornom aktivnošću i promoviše razvoj ateroskleroze kod ljudi. Rezultati ovih istraživanja poslednjih godina time potvrđuju ulogu i značaj crevne mikrobiote i disbioze u riziku od ateroskleroze [8,9,10,17].

U metabolizmu crevnih bakterija nastaju različiti metaboliti koji učestvuju u razvoju ateroskleroze. Među najvažnijim su različiti amini, metilamini, poliamini, masne kiseline kratkog lanca, trimetilamin i sekundarne žučne kiseline. Konkretno, masne kiseline sa kratkim lancem (SCFA - Short-chain fatty acid) su grupa crevnih mikrobni metabolita koji su važni za metaboličke bolesti. Studije su pokazale da je crevna mikrobiota uključena u stvaranje trimetilamin N-oksida (TMAO) [8,14]. Trimetilamin (TMA) je nusprodukt bakterijskog metabolizma koji se apsorbuje u krvotok i pretvara u TMAO u jetri pomoću specifičnih jetrenih enzima, monooksigenaza koje sadrže flavin. Različiti bakterijski sastavi prirodno imaju različite sposobnosti da formiraju TMAO. Studije na miševima su potvrdile da TMAO ubrzava razvoj ateroskleroze stimulisanjem priliva holesterola, inhibicijom izlučivanja holesterola, inhibicijom metabolizma sekundarnih žučnih kiselina i/ ili prekomernom aktivacijom trombocita [3,8,10]. Prema istraživačima, pored uloge biološkog markera za aterosklerozu i kardiovaskularne bolesti, TMAO bi mogao da predstavlja i mogući terapijski cilj u budućnosti. Zanimljivo je da su razvijeni inhibitori proizvodnje TMAO koji ciljaju različite mikrobne TMA liaze. Ovi lekovi smanjuju nivo TMAO i preokreću aterosklerozu na životinjskim modelima. TMA liaza je postala trenutni potencijalni terapijski cilj TMAO modulacije [18].

Mikrobiota creva i hipertenzija

Osim dislipidemije i ateroskleroze, hipertenzija je još jedan od najvažnijih faktora

composition of microbiota on various chronic and autoimmune diseases has been studied, especially in animal trials [3,8]. These trials demonstrate the importance of the microbiota in relation to health and immunity and offer new, as yet undiscovered possibilities for the use of this knowledge in the treatment of some other diseases, such as metabolic syndrome, insulin resistance, chronic inflammatory bowel disease, cancer [1,6,9].

Gut dysbiosis and atherosclerosis

Atherosclerosis is the main risk factor for cardiovascular diseases. This process is characterized by the accumulation of cholesterol and macrophages (inflammatory cells) in the vascular walls, which contributes to the formation of atherosclerotic plaques. Recent studies have shown that gut dysbiosis can contribute to the development of atherosclerosis by modulating inflammatory processes and by forming certain microbial metabolites [10-15]. Integrity of the gut mucosa is the first barrier that protects the host from the intrusion of pathogens, the passage of intestinal contents and bacterial components into the blood vessels. Reduced concentrations of proteins that ensure close cell contacts and their impermeability, including zonula occludens-1 (ZO-1) also known as Tight junction protein-1 (TJP1), claudin 1, and occludin, allow increased permeability of the gut wall by disrupting the balance between mucosal cell death and regeneration of the mucosal cells [1,13,14]. If the mucous barrier is damaged, the penetration of microbes and their products into the blood vessels triggers an immune response, tissue and systemic inflammation. Impairment of the gut barrier integrity caused by gut dysbiosis, therefore, acts as a risk factor that triggers a chronic inflammation that underlies various diseases, including atherosclerosis. The main molecules-products of bacteria that are the drivers of the immune and inflammatory response are "Pathogen-associated molecular pattern"- PAMP. PAMPs activate the innate immune response, protecting the host from infection. A wide range of different types of molecules can serve as PAMP, including glycans and glycoconjugates. Bacterial lipopolysaccharides (LPS), endotoxins found on cell membranes of Gram-negative bacteria, are considered a prototype class of PAMP. The relationship between plasma LPS levels and cardiovascular risk was first studied in 1999 by Niebauer et al. [15]. The results of the study

confirmed that the level of endotoxemia was the highest in patients with the most severe cardiovascular disease. Cani et al. confirmed in their study that gut dysbiosis prevented the formation of "close contact proteins", which resulted in increased permeability of gut mucosa, and thus the passage of LPS into the blood [16]. LPS, which are produced in increased amounts in intestinal dysbiosis, can play an important role in modulating "toll-like receptors (TLRs)" which recognize bacterial products and regulate the host's immune system. TLRs are a class of proteins that play a key role in the innate immune system. They are single-pass transmembrane receptors commonly found on sentinel cells (first-line defense cells) such as macrophages and dendritic cells, which recognize structurally conserved molecules derived from microbes. Once microbes break through physical barriers such as the skin or lining of the gut tract, they are recognized by TLRs that activate immune cell responses. Clinical studies have shown that an increase in TLR is associated with anti-inflammatory activity and promotes the development of atherosclerosis in humans. The results of these studies in recent years thus confirm the role and importance of gut microbiota and dysbiosis as risk factors in the development of atherosclerosis [8,9,10,17].

In the metabolism of gut bacteria, various metabolites that participate in the development of atherosclerosis are formed. Among the most important are various amines, methylamines, polyamines, short-chain fatty acids, trimethylamine and secondary bile acids. In particular, short-chain fatty acids (SCFA) are a group of intestinal microbial metabolites that are important for metabolic diseases. Studies have shown that the intestinal microbiota is involved in the formation of trimethylamine N-oxide (TMAO) [8,14]. Trimethylamine (TMA) is a by-product of bacterial metabolism that is absorbed into the bloodstream and converted to TMAO in the liver by specific liver enzymes, flavin-containing monooxygenases. Different bacterial compositions naturally have different abilities to form TMAO. Studies in mice have confirmed that TMAO accelerates the development of atherosclerosis by stimulating cholesterol influx, inhibiting cholesterol excretion, inhibiting secondary bile acid metabolism, and / or by excessive platelet activation [3,8,10]. According to researchers,

rizika za KVB koji je genetski osjetljiv i pod uticajem faktora okoline [19]. Još 1982. godine pokazano je da lečenje antibioticima može prouzrokovati viši krvni pritisak [15]. Sa druge strane, jedan niz studija pokazao je povoljan efekat korišćenja antibiotika na krvni pritisak. Ovi podaci, kao i uočena veza između disbioze i cerebrovaskularnih događaja je indirektno upućivalo na vezu između mikrobiote creva i hipertenzije koja je procenjivana u novijim studijama [20,21]. Dalje, kod spontano hipertenzivnih pacova dokumentovano je da dolazi do značajnog smanjenja broja i raznolikosti mikroba u crevima i smanjenja broja cecalnih „dobrih bakterija“ iz vrste *Bacteroidetes*, što je praćeno proporcionalnim povećanjem broja „loših bakterija“ iz vrste *Firmicutes*. Studije su, takođe, pokazale da transplantacija cecalnog mikrobiološkog sadržaja od hipertenzivnih životinja davaoca može reprodukovati hipertenziju kod prethodno normotenzivnih životinja [22]. U studiji na miševima pokazano je da disbioza crevne mikrobiote može da izazove angiotenzin-II-indukovanu vaskularnu disfunkciju i hipertenziju. Kao što je u drugom istraživanju utvrđeno da odsustvo crevne mikrobiote štiti miševima od arterijske hipertenzije indukovane angiotenzinom II, vaskularne disfunkcije i oštećenja krajnjih organa izazvanih hipertenzijom [23,24].

Dakle, očigledno je da je mikrobiota creva uključena u razvoj ili pogoršanje hipertenzije. Iako tačni osnovni mehanizmi i međusobna veza između mikrobiote creva i hipertenzije nisu utvrđeni, postojeći dokazi iz istraživanja na životinjama i kliničkih studija ističu ulogu masnih kiselina kratkog lanca -SCFA i oksidisanog lipoproteina male gustine (ox-LDL) u nastanku hipertenzije. Masne kiseline kratkog lanca poput acetata, proprionata i butirata, nastaju uglavnom iz polisaharida rastvorljivih dijetalnih vlakana [23]. Grupe mikroba u crevima koje metabolizuju polisaharide u različite vrste SCFA su specifične. Glavne bakterije koje proizvode acetat su *Streptococcus*, *Prevotella*, *Bifidobacterium*, *Clostridium* i *A. Muciniphila* [25]. Propionate generišu *Bacteroides*, *Salmonella*, *Dialister*, *Veillonella*, *Roseburia*, *Coprococcus*, *Blautia*, i dr. [26]. Butirate stvaraju *Lachnospiraceae*, *Ruminococcac*, i *Acid amino coccaceae* [27]. Suplementacija vlakana i acetata dovodi do povećanja broja *Bacteroides acidifaciens* i bila je povezana sa poboljšanjem

crevne disbioze, hipertenzije i srčane insuficijencije kod hipertenzivnih miševa [28]. Preveliki broj bakterija koje proizvode butirat povezan je sa povišenim sistolnim i dijastolnim krvnim pritiskom kod trudnica [29].

G protein vezani receptori (GPCR) su receptori na ćelijskoj površini koji otkrivaju molekule SCFA izvan ćelije i aktiviraju ćelijske odgovore. Tri GPCR-a koja su regulisani SCFA-om su: GPR41, GPR43 i GPR109A [30]. SCFA stimulišu puteve regulisane GPCR-om da utiču na sistem renin-angiotenzin da modulira krvni pritisak.

Olfaktorni receptor 78 (Olf78) je još jedan tip GPCR-a eksprimiran u bubregu koji detektuje SCFA [31]. I Olf78 i GPR41 su izraženi u ćelijama glatkih mišića krvnih sudova malog prečnika. U drugoj studiji, stimulacija GPR41 rezultirala je smanjenjem hipotenzivnog odgovora [32]. SCFA, propionat indukuje vazodilataciju i proizvodi akutni hipotenzivni odgovor kod miševa modulacijom aktivnosti Olf78 i GPR41 [33]. Svi ovi nalazi otkrivaju da mikrobiota creva igra važnu ulogu u modulaciji krvnog pritiska preko SCFA i upućuje da je hipertenzija povezana sa disbiozom.

Pored izmenjene regulacije različitih receptora preko SCFA, crevna disbioza, takođe, doprinosi hipertenziji vazokonstrikcijom posredovanom oksidacijom LDL [34]. Mikrobna disbioza promoviše ekspresiju pro-inflamatornih citokina i indukuje oksidativni stres koji stimuliše oksidaciju LDL [35]. Viši nivoi oksidisanog LDL (Ox-LDL) smanjuju proizvodnju NO, smanjuju stepen vazodilatacije i stimulišu stvaranje vazokonstriktornih supstanci među kojima je i endotelin-1 koji igra presudnu ulogu u održavanju vaskularnog tonusa i kardiovaskularne homeostaze. Narušena ravnoteža dovodi do hipertenzije [35]. Ipak, uzročni odnos između disbioze creva i hipertenzije je složen i nije u potpunosti procenjen. Tačna uloga crevne mikrobiote u posredovanju hipertenzije, uključeni putevi i mehanizmi zahtevaju dalja detaljna istraživanja.

Crevna mikrobiota i srčana insuficijencija

Sve je više dokaza o povezanosti mikrobiote creva i patogeneze srčane insuficijencije. U engleskoj literaturi se za definisanje ove veze koristi izraz „crevna hipoteza srčane insuficijencije“ - “gut hypothesis of heart failure” [36-39]. Ova hipoteza objašnjava da smanjeni minutni volumen srca

apart from the role of a biological marker for atherosclerosis and cardiovascular diseases, TMAO could also represent a possible therapeutic goal in the future. Interestingly, inhibitors of TMAO production have been developed that target various microbial TMA lyases. These drugs reduce TMAO levels and reverse atherosclerosis in animal models. TMA lyase has become the current potential therapeutic target of TMAO modulation [18].

Gut microbiota and hypertension

Apart from dyslipidemia and atherosclerosis, hypertension is another major risk factor for CVD that is genetically sensitive and influenced by environmental factors [19]. As early as 1982, it was shown that antibiotic treatment could cause higher blood pressure [15]. On the other hand, a number of studies showed that antibiotic use had a beneficial effect on blood pressure. These data, as well as the observed relationship between dysbiosis and cerebrovascular events, indirectly suggested a relationship between the gut microbiota and hypertension as assessed in recent studies. [20,21]. Furthermore, in spontaneously hypertensive rats, a significant decrease in the number and diversity of microbes in the gut and a decrease in the number of cecal "good bacteria" from the species *Bacteroidetes* was documented, which was accompanied by a proportional increase in the number of "bad bacteria" from the species *Firmicutes*. The studies have also shown that transplantation of cecal microbiological content from hypertensive animal donors can reproduce hypertension in previously normotensive animals [22]. In a study on mice, it was shown that dysbiosis of the gut microbiota can cause angiotensin-II-induced vascular dysfunction and hypertension. As another study found, the absence of gut microbiota protects mice from angiotensin II-induced arterial hypertension, vascular dysfunction, and end-organ damage caused by hypertension [23,24].

Thus, it is obvious that the gut microbiota is involved in the development or worsening of hypertension. Although the exact underlying mechanisms and the relationship between the gut microbiota and hypertension have not been established, existing evidence from animal trials and clinical studies highlights the role of short-chain fatty acids - SCFA and oxidized low-density lipoprotein (ox-LDL) in the

development of hypertension. Short-chain fatty acids, such as acetate, propionate, and butyrate, are formed mainly from soluble dietary fiber polysaccharides [23]. The groups of microbes in the gut that metabolize polysaccharides to different types of SCFA are specific. The main acetate-producing bacteria are *Streptococcus*, *Prevotella*, *Bifidobacterium*, *Clostridium*, and *A. Muciniphila* [25]. Propionates are produced by *Bacteroides*, *Salmonella*, *Dialister*, *Veillonella*, *Roseburia*, *Coprococcus*, *Blautia*, and others. [26]. Butyrates are produced by *Lachnospiraceae*, *Ruminococcaceae* Acid amino coccaceae [27]. Fiber and acetate supplementation led to an increase in the number of *Bacteroides acidifaciens* and was associated with improved gut dysbiosis, hypertension, and heart failure in hypertensive mice [28]. Too many butyrate-producing bacteria have been associated with elevated systolic and diastolic blood pressure in pregnant women [29]. G protein-coupled receptors (GPCRs) are receptors on the cell surface that detect SCFA molecules outside the cell and activate cellular responses. The three GPCRs regulated by SCFA are: GPR41, GPR43, and GPR109A [30]. SCFAs stimulate GPCR-regulated pathways to affect the renin-angiotensin system to modulate blood pressure. Olfactory receptor 78 (Olf78) is another type of GPCR expressed in the kidney that detects SCFA [31]. Both Olf78 and GPR41 are expressed in smooth muscle cells of small-diameter blood vessels. In another study, stimulation of GPR41 resulted in a reduction in the hypotensive response [32]. SCFA, propionate induces vasodilation and produces an acute hypotensive response in mice by modulating the activity of Olf78 and GPR41 [33]. All these findings reveal that the gut microbiota plays an important role in modulating blood pressure via SCFA and suggests that hypertension is associated with dysbiosis.

Apart from altered regulation of various receptors via SCFA, gut dysbiosis also contributes to hypertension by Oxidized LDL-mediated vasoconstriction [34]. Microbial dysbiosis promotes the expression of pro-inflammatory cytokines and induces oxidative stress that stimulates LDL oxidation [35]. Higher levels of oxidized LDL (Ox-LDL) reduce NO production, reduce the degree of vasodilation and stimulate the production of vasoconstrictor substances, including endothelin-1, which plays

(MVS) i povećana sistemski zastoj mogu prouzrokovati crevnu ishemiju i/ili edem crevnog zida, što dovodi do povećanog prodiranja bakterija u krvne sudove, povećavajući tako koncentraciju endotoksina u cirkulaciji. To može pokrenuti upalu kod pacijenata sa srčanom insuficijencijom. Egzogeni faktori kao što su dijeta, izloženost bakterijskim infekcijama ili uzimanje lekova mogu smanjiti raznolikost crevne flore. Endogeni faktori kao što su akutna humoralna neravnoteža, hronična crevna kongestija odnosno ishemija-hipoksija, kiselinsko-bazni disbalans, oslabljena gastrointestinalna pokretljivost i nutritivni nedostatak mogu potencijalno promeniti crevnu floru [40]. Sa razvojem srčane insuficijencije, menjaju se i karakteristike zajednice bakterija. Studije su pokazale da se brojnost crevne flore kod pacijenata sa hroničnom srčanom insuficijencijom smanjila, a broj patogenih bakterija značajno se povećavao sa napredovanjem bolesti, uključujući *Campylobacter*, *Shigella*, *Salmonella*, *Iersinia enterocolitica* i *Candida* vrste [41,42,43]. Analiza 16SrRNA kod pacijenata sa srčanom insuficijencijom izvestila je o smanjenju bakterija koje proizvode SCFA, kao što su *Eubacterium rectale* i *Dorea longicatena* [43]. Druga studija je pokazala da se sastav mikrobiote creva kod hronične srčane insuficijencije odlikuje smanjivanjem broja bakterija sa potencijalom za proizvodnju butirata [44]. Butirat vrši lokalne antiinflamatorne efekte u sluznici creva i stimuliše regulatorne T-ćelije [45]. Primećeno je da je brojnost mikrobioloških gena za LPS biosintezu i stvaranje TMAO povišena, a gena za butirat acetoacetat koenzim A transferaze (ključni enzim za proizvodnju butirata) smanjena u hroničnoj srčanoj insuficijenciji [41].

Primećeno je, takođe, da su pacijenti sa srčanom insuficijencijom i sa perifernim edemima imali viši nivo endotoksina i inflamatornih citokina u plazmi u poređenju sa pacijentima bez edema. Nakon kratkotrajne terapije diureticima, koncentracije endotoksina u serumu, ali ne i citokina, su se smanjile [46]. U drugoj studiji istraživači su potvrdili da su pacijenti sa srčanom insuficijencijom i smanjenim crevnim protokom krvi imali veće serumske koncentracije imunoglobulina A - anti-lipopolisaharida. U poređenju sa kontrolnom grupom, pacijenti su imali drugačiji sastav mikrobiote, broj korisnih bakterija se smanjio, a

patogenih povećao [24]. Pojava i razvoj srčane insuficijencije mogu biti povezani sa smanjenjem bakterija koje proizvode SCFA i porastom bakterija koje proizvode TMAO, što može postati nova meta za lečenje srčane insuficijencije. Nedavno su prikazani efekti inhibitora enzima trimetilamin liaze koji u studijama na miševima pokazuju učinak sličan dejstvu anti-aterotrombotičkih agenasa. [47,48].

Mikrobiota creva i infarkt miokarda

Aterosklerotični plakovi sadrže bakterijsku DNK. Vrste bakterija koje se nalaze u aterosklerotskim plakovima, međutim, prisutne su i u crevima istih pojedinaca [18,19,36]. Studija iz Južne Koreje pokazuje da je tokom akutne faze STEMI u koronarnom trombu detektovano prisustvo bakterija (mikrobna rRNA). Mikrobiološki potpis u koronarnom trombu pokazao je korelaciju sa oralnim i crevnim mikrobiomom [20]. Iz ovoga se može zaključiti da crevne mikrobne zajednice mogu biti izvor bakterija u plaku, što može uticati na stabilnost plaka i razvoj kardiovaskularnih bolesti. Nedavno istraživanje na pacovima izvestilo je o povezanosti između crevne mikrobiote i obima infarkta miokarda [38,39]. Studija je proučavala Dahl S pacove (Dahl Salt Sensitive Rat- pacovi hranjeni hranom sa visokim sadržajem soli - 8% NaCl) koji su pili vodu za piće uz dodatak antibiotika vankomicina, koji je smanjio nivo cirkulišućeg leptina za 38%, prouzrokovao po obimu manji infarkt miokarda (smanjenje površine od 27%) i poboljšao obnavljanje postishemijske kontraktilnosti miokarda u poređenju sa kontrolnim životinjama koje nisu dobijale antibiotik. Vankomicin je izmenio obilnost crevnih bakterija i gljivica mereno količinom 16S i 18S rRNA [39].

U studijama na glodarima, primena *Lactobacillus plantarum* kao probiotika rezultirala je smanjenjem cirkulišućeg leptina za 41%, infarktom miokarda za 29% i boljim oporavkom kontraktilne funkcije miokarda za 23%. Međutim, ako su glodari pre studije primili leptin u dozi od 0,12 µg / kg i.v. zaštitni efekat probiotika na srce je bio poništen. Ova studija je prva koja je potvrdila direktnu vezu između promena crevne mikrobiote i infarkta miokarda. To pokazuje da dodatak probiotika može smanjiti stepen infarkta miokarda [46]. Druga studija na životinjskom modelu koja je koristila *Lactobacillus rhamnosus* pokazala je povoljan

a crucial role in maintaining vascular tone and cardiovascular homeostasis. Disturbed balance leads to hypertension. However, the causal relationship between gut dysbiosis and hypertension is complex and has not been fully assessed. The exact role of the gut microbiota in mediating hypertension, the pathways and mechanisms involved require further detailed research.

Gut microbiota and heart failure

There is growing evidence of a connection between the gut microbiota and the pathogenesis of heart failure. In the English literature, the term "gut hypothesis of heart failure" [36-39] is used to define this connection. This hypothesis explains that decreased cardiac output (DCO) and increased systemic arrest can cause gut ischemia and / or edema of the gut wall, leading to increased bacterial penetration into blood vessels, thus increasing the concentration of endotoxins in the circulation. This can trigger inflammation in patients with heart failure. Exogenous factors such as diet, exposure to bacterial infections, or medication may reduce the diversity of the gut flora. Endogenous factors such as acute humoral imbalance, chronic gut congestion or ischemia-hypoxia, acid-base imbalance, impaired gastrointestinal motility, and nutritional deficiency can potentially alter the gut flora [40]. With the development of heart failure, the characteristics of the bacterial community change. Studies have shown that the number of gut flora in patients with chronic heart failure decreased, and the number of pathogenic bacteria increased significantly with the progression of the disease, including *Campylobacter*, *Shigella*, *Salmonella*, *Yersinia enterocolitica* and *Candida* species [41,42,43]. 16SrRNA analysis in patients with heart failure has reported a reduction in SCFA-producing bacteria, such as *Eubacterium rectale* and *Dorea longicatena* [43]. Another study showed that the composition of the gut microbiota in chronic heart failure is characterized by a decrease in the number of bacteria with the potential to produce butyrate [44]. Butyrate exerts local anti-inflammatory effects in the gut mucosa and stimulates regulatory T cells [45]. It has been observed that the number of microbiological genes for LPS biosynthesis and TMAO production is increased, while the abundance of genes for butyrate acetoacetate coenzyme A

transferase (key enzyme for butyrate production) is reduced in chronic heart failure [41]. It was also observed that patients with heart failure and peripheral edema had higher plasma endotoxin and inflammatory cytokine levels compared with patients without edema. After short-term diuretic therapy, serum endotoxin but not cytokine concentrations [46] decreased. In another study, the researchers confirmed that patients with heart failure and reduced gut blood flow had higher serum concentrations of immunoglobulin A - anti-lipopolysaccharide. Compared with the control group, patients had a different composition of the microbiota, the number of beneficial bacteria decreased, and the number of pathogens increased [24]. The onset and development of heart failure may be associated with a decrease in SCFA-producing bacteria and an increase in TMAO-producing bacteria, which may become a new target for the treatment of heart failure. Recently, in studies in mice, the effects of trimethylamine-lyase enzyme inhibitors have been shown to have an effect similar to that of anti-atherothrombotic agents. [47,48].

Gut microbiota and myocardial infarction

Atherosclerotic plaques contain bacterial DNA. However, the types of bacteria found in atherosclerotic plaques are also present in the gut of the same individuals [18,19,36]. A study from South Korea shows that the presence of bacteria (microbial rRNA) was detected in the coronary thrombus during the acute phase of STEMI. The microbiological signature in the coronary thrombus showed a correlation with oral and intestinal microbiome [20]. From this it can be concluded that gut microbial communities can be a source of bacteria in plaque, which can affect plaque stability and the development of cardiovascular diseases. A recent study in rats reported a connection between the gut microbiota and the extent of myocardial infarction [38,39].

The study looked into Dahl Salt-Sensitive Rats - rats fed with foods high in salt - 8% NaCl) that drank drinking water with antibiotic vancomycin, which reduced the level of circulating leptin by 38%, causing a smaller myocardial infarction (area reduction of 27%) and improved restoration of postischemic myocardial contractility compared to control animals that did not receive the antibiotic.

efekat na srčanu funkciju nakon veštački izazvanog infarkta miokarda [49].

Mikrobiota creva i metaboličke bolesti

Mnoga istraživanja pokazuju vezu između sastava crevne mikrobiote i metaboličkih poremećaja u organizmu [50,51,52]. Uloga mikrobiote creva u razvoju gojaznosti dokazana je studijama sprovedenim na obeskličenim miševima (germfree-GF-miševi) u poređenju sa konvencionalno gajenim miševima (CONV-R). Sterilni miševi bez klica se uzgajaju u izolatorima koji u potpunosti blokiraju izloženost mikroorganizmima, sa namerom da ih sačuvaju od otkrivenih bakterija, virusa i eukariotskih mikroba. CONV-R imaju 40% veći sadržaj telesne masti od GF-miševa, što je fenomen nezavisan od unosa hrane. Međutim, nakon kolonizacije GF-miševa crevnom florom koja dolazi od CONV-R miševa, uočen je značajan porast telesne težine i ~ 60% porast telesne masti, zajedno sa povećanom sintezom jetrenih triglicerida u primaocima fekalnog transplantata (GF miševi), nezavisno od unosa hrane i ukupne potrošnje energije [53]. Čini se da je mehanizam putem kojeg crevni mikrobi doprinose povećanoj apsorpciji energije stvaranjem kratkolančanih masnih kiselina (SCFA), što je rezultat hidrolize i fermentacije dijetalnih polisaharida. SCFA, kao što su propionat, butirat i acetat, vrše složene metaboličke akcije koje utiču na apetit domaćina, vreme crevnog tranzita, apsorpciju i taloženje masti [52]. SCFA, takođe, povećavaju unutrašnju apsorpciju monosaharida stimulišući ekspresiju transportera natrijuma/glukoze 1. SCFA doprinose modulaciji apetita domaćina i unosu hrane u interakciji sa G-vezanim proteinima eksprimiranim enteroendokrinim ćelijama i promovišu oslobađanje glukagona sličnog peptida-1 (GLP-1) koji utiče na sitost. Pored toga, SCFA utiču na metabolizam lipida povećavajući lipogenezu i inhibirajući oksidaciju masnih kiselina [53]. Studije su dokazale specifične promene u sastavu mikrobiote creva kod genetski gojaznih miševa u poređenju sa genetski mršavim miševima, pokazujući smanjenje obilja *Bacteroidetes*-a za 50% i proporcionalno povećanje *Firmicutes*-a. Čini se da ove specifične promene doprinose povećanoj proizvodnji SCFA i nagomilavanju masti kod gojaznih miševa i kod GF miševa kolonizovanih mikrobiotom gojaznih miševa [54]. Postoje i drugi mogući mehanizmi. Dijeta sa visokim

sadržajem masti pokazala je da povećava udeo gram-negativnih vrsta u mikrobioti creva, što doprinosi povećanoj apsorpciji crevnih fragmenata bakterija, poput lipopolisaharida (LPS) u crevima. Povećani nivoi LPS u cirkulaciji dovode do „metaboličke endotoksemije“ koja se manifestuje kao povećanje telesne težine, hiperglikemija natašte i hiperinsulinemija [55]. Sve je više dokaza koji ukazuju na to da dijeta sa visokim sadržajem masti promoviše promene u sastavu mikrobiote creva, ali je kasniji razvoj gojaznog fenotipa povezan sa metaboličkom endotoksemijom [56].

Poslednjih godina istraživači su, takođe, proučavali veze između disbioze i gojaznosti, dijabetesa tipa 2, dislipidemije i nealkoholne masne jetre (NASH) [50,51]. Početna ispitivanja na životinjama i ljudima podržala su povezanost između gojaznosti i obilja grupe bakterija *Firmicutes* u poređenju sa grupom *Bacteroidetes*; dijabetes tipa 2, međutim, povezan je sa smanjenim obiljem bakterija koje formiraju butirat i povećanim obiljem *Lactobacillus spp* [1,3,8,10]. Crevna mikrobiota je uključena u razvoj dislipidemija putem sekundarnih žučnih kiselina [52,53]. U istraživanju NASH-a utvrđeno je da neke bakterije (*Clostridium coccoides*, *Lactobacillus reuteri*, *Parabacteroides*) utiču na metabolizam masti, integritet crevnog zida i proces fibroze i tako utiču i na dislipidemiju [52].

Primena u praksi

Primeri kliničke korisnosti dugotrajne promene mikrobiote su: dijetetske mere, pre- i probiotička terapija, antibiotska terapija, unos ciljanih inhibitora enzima, fekalna mikrobna transplantacija itd. [57,58].

Istraživanja su pokazala da čak i petodnevna promena u ishrani dovodi do kratkotrajnog preuređivanja broja i vrsta crevnih mikroba [4]. Primer za to je prehrambeni pristup za zaustavljanje hipertenzije (DASH dijeta -Dietary Approaches to Stop Hypertension) koji se sastoji od obroka sa voćem, povrćem, integralnim žitaricama itd. [59]. Pacijenti u studiji na ovoj dijeti su pokazali poboljšanje kvaliteta života i bolju elastičnost arterijskih krvnih sudova nakon tri meseca pridržavanja mera [60]. Pored toga, opisano je da osobe koje se ne pridržavaju propisane dijeta i imaju „zapadnjačku dijetu“ sa dosta masti i crvenog mesa imaju povišeni nivo TMAO u urinu u poređenju sa pacijentima koji se

Vancomycin changed the abundance of gut bacteria and fungi measured by the amount of 16S and 18S rRNA [39].

In rodent-based studies the use of *Lactobacillus plantarum* as a probiotic resulted in the reduction of circulating leptin by 41%, myocardial infarction by 29% and better recovery of myocardial contractile function by 23%. However, if rodents received leptin at a dose of 0.12 µg / kg i.v. the protective effect of probiotics on the heart was reversed. This study is the first to confirm a direct link between changes in the gut microbiota and myocardial infarction. This shows that the addition of probiotics can reduce the degree of myocardial infarction [46]. Another animal study using *Lactobacillus rhamnosus* showed a beneficial effect on cardiac function after an artificially induced myocardial infarction [49].

Gut microbiota and metabolic diseases

Many studies show a link between the composition of the gut microbiota and metabolic disorders in the body [50,51,52]. The role of the gut microbiota in the development of obesity has been proven by studies conducted on dehydrated mice (germfree-GF-mice) compared with conventionally bred mice (CONV-R). Sterile, germ-free mice are bred in isolators that completely block exposure to microorganisms, with the intention of protecting them from detected bacteria, viruses, and eukaryotic microbes. CONV-R mice have a 40% higher body fat content than GF-mice, which is a phenomenon independent of food intake. However, after colonization of GF-mice by gut flora coming from CONV-R mice, a significant increase in body weight and ~ 60% increase in body fat was observed, together with increased synthesis of liver triglycerides in faecal transplant recipients (GF mice), independent of food intake and total energy consumption [53]. It appears to be the mechanism by which gut microbes contribute to increased energy absorption by the formation of short-chain fatty acids (SCFA), which is the result of hydrolysis and fermentation of dietary polysaccharides. SCFAs, such as propionate, butyrate, and acetate, perform complex metabolic actions that affect host appetite, gut transit time, and fat absorption and deposition [52]. SCFAs also increase the internal absorption of monosaccharides by stimulating the expression of the sodium-

glucose transporter 1. SCFAs also contribute to the modulation of host appetite and food intake in interaction with G-linked proteins expressed by enteroendocrine cells and promote the release of glucagon-like peptide-1 that affects satiety. Apart from that, SCFAs affect lipid metabolism by increasing lipogenesis and inhibiting fatty acid oxidation [53]. Studies have shown specific changes in the composition of the gut microbiota in genetically obese mice compared to genetically lean mice, showing a 50% decrease in abundance of Bacteroidetes and a proportional increase in Firmicutes. These specific changes appear to contribute to increased SCFA production and fat accumulation in obese mice and in GF mice colonized by obese microbiota [54]. There are other possible mechanisms. A high-fat diet has been shown to increase the proportion of Gram-negative species in the gut microbiota, which contributes to the increased absorption of gut fragments of bacteria, such as lipopolysaccharides (LPS) in the gut. Increased circulating LPS levels lead to "metabolic endotoxemia" which manifests as weight gain, fasting hyperglycemia, and hyperinsulinemia [55]. There is growing evidence to suggest that a high-fat diet promotes changes in the composition of the gut microbiota, but the later development of the obesity phenotype is associated with metabolic endotoxemia [56]. In recent years, researchers have also studied the links between dysbiosis and obesity, type 2 diabetes, dyslipidemia, and nonalcoholic fatty liver disease (NAFLD) [50,51]. Initial studies in animals and humans supported a correlation between obesity and the abundance of the Firmicutes group of bacteria compared to the Bacteroidetes group; type 2 diabetes, however, is associated with a reduced abundance of butyrate-forming bacteria and an increased abundance of *Lactobacillus* spp [1,3,8,10]. The gut microbiota is involved in the development of dyslipidemias via secondary bile acids [52,53]. In the research of NAFLD, it was determined that some bacteria (*Clostridium* spp, *Lactobacillus reuteri*, *Parabacteroides*) affect fat metabolism, integrity of the gut wall and the process of fibrosis, thus affecting dyslipidemia [52].

Application in practice

Examples of clinical benefit of long-term microbiota change are: dietary measures, pre and probiotic therapy, antibiotic therapy, intake

pridržavaju propisanog DASH režima [61,62]. Smanjen unos dijetalnih vlakna povezan je sa smanjenom bakterijskom proizvodnjom butirata masnih kiselina kratkog lanca, koji ima imuno-modulacione efekte u sluznici creva, a takođe služi i kao glavni energetski supstrat za kolonocite. Smanjenje nivoa butirata u crevima moglo bi da podstakne lokalnu upalu, pogorša disbiozu i doprinese oštećenju funkcije crevne barijere, što posledično dovodi do „curenja“ bakterijskih toksina kao što je LPS, što dodatno podstiče lokalnu i sistemsku upalu. Dijeta bogata vlaknima može poboljšati rast bakterija koje proizvode acetat, smanjiti visok krvni pritisak i sprečiti srčanu fibrozu i hipertrofiju [63].

Probiotici i prebiotici

Probiotici su živi mikroorganizmi koji, kada se daju u odgovarajućim količinama, domaćinu donose zdravstvenu korist [64]. Probiotici u kliničkoj upotrebi sadrže bakterijske i gljivične mikroorganizme, uključujući rodove *Lactobacillus* i *Bifidobacterium* i gljivicu *Saccharomyces boulardii* [65]. Rezultati životinjskih modela sugerišu da bi određeni sojevi laktobacila mogli imati kardioprotektivne efekte. Pacovi lečeni suplementom koji sadrži *Lactobacillus plantarum* 299v pre podvezivanja koronarne arterije smanjili su veličinu infarkta i poboljšali funkciju leve komore [66]. Druga studija pokazala je slične kardioprotektivne rezultate na pacovskom modelu ishemije miokarda nakon suplementa sa *Lactobacillus rhamnosus* GR-1 [67]. Pilot studija na ljudima izvestila je ne samo o smanjenom sistemskom zapaljenju, već i o poboljšanju eejekcione frakcije leve komore nakon intervencije sa probiotičkim kvasnicama *Saccharomyces Boulardii* kod pacijenata sa hroničnom srčanom insuficijencijom [68]. S obzirom na potencijalni klinički uticaj modulacije mikrobiota, kao i visok morbiditet i mortalitet od srčane insuficijencije, modulacija mikrobiote nije potpuno bez rizika [69]. Preporučuje se brižno kliničko praćenje i unapred definisane mere bezbednosti koje treba da slede iste standarde kao u drugim kliničkim ispitivanjima [70]. Ovo stoga što su nedavno prijavljeni genomske i epidemiološki dokazi o bakterijemiji probiotičkih bakterija, odnosno o prenosu bakterija sa probiotičkih kapsula u krv kod pacijenata u jedinicama intenzivne nege [71].

Prebiotici su supstrati koje mikroorganizmi domaćina selektivno koriste i

daju potencijalnu korist za zdravlje. Kao prebiotici najčešće se koriste dijetalna vlakna i oligosaharidi [72]. Većina savremenih studija obrade mikrobiote kod pacijenata sa kardiovaskularnim bolestima izveštava o iscrpljivanju mikroba sa kapacitetom stvaranja SCFA kao što je butirat. Prebiotici koji promovišu mikrobiološku fermentaciju dijetalnih vlakana u SCFA mogu stoga biti od potencijalne koristi u poboljšanju metaboličke regulacije [73]. Neki prebiotici, poput inulina, imaju potencijal da se suprotstave štetnim efektima antibiotika promovišući raznolikost i funkcionalni kapacitet crevne mikrobiote [74]. Randomizovana studija sa suplementom hrane inulinom ili inulin-propionatnim estrom pokazala je smanjenje markera sistemske upale sa povećanim generisanjem SCFA proprionata u debelom crevu [75]. Stoga su ciljanje proizvodnje mikrobioloških SCFA dodacima inulina ili drugih prebiotika atraktivna strategija za buduća ispitivanja kardiovaskularnih bolesti, mada trenutni naučni dokazi ne daju potvrđene preporuke za upotrebu probiotika ili prebiotika kao dopunske terapije kod pacijenata sa srčanom insuficijencijom ili koronarnom bolešću.

Antibiotici. Upotreba antibiotika utiče na sastav, raznovrsnost i funkciju normalne flore [76]. Antibiotici su uspešno korišćeni na životinjskim modelima za smanjenje stepena oštećenja srčanih ćelija nakon infarkta miokarda [77,78]. Ranije studije kod pacijenata sa srčanom insuficijencijom, fokusirale su se na dekontaminaciju creva antibioticima širokog spektra radi smanjenja translokacije biotoma i bakterijske upale. Iako je ovaj pristup uspeo u smanjenju markera sistemske upale, klinički efekat nije dokazan [79,80]. U nedavnoj studiji je pokazano da je koktel oralnog antibiotika širokog spektra znatno povećao postinfarktnu rupturu i smrt u mišjem modelu ligacije koronarne arterije [81] što bi moglo da sugeriše da je za pravilan oporavak miokarda potrebna netaknuta mikrobna zajednica u vreme povrede miokarda. Ova studija je u suprotnosti sa prethodnim životinjskim eksperimentalnim modelom koji je pokazao da je oralni vankomicin smanjio veličinu infarkta i poboljšao postinfarktnu srčanu funkciju kod pacova [82], kao i sa studijom koja izveštava da je mešavina streptomcina, neomicina, polimiksina B i bacitracina smanjila veličinu infarkta zajedno sa promenama u metabolitima povezanim sa

of targeted enzyme inhibitors, fecal microbial transplantation, etc. [57,58]. Studies have shown that even a five-day change in diet leads to a short-term rearrangement of the number and types of gut microbes [4]. An example of this is the dietary approach aimed at stopping hypertension (DASH diet - Dietary Approaches to Stop Hypertension), which consists of meals with fruits, vegetables, whole grains, etc. [59]. Patients in the study who were on this diet showed an improvement in quality of life and better elasticity of arterial blood vessels after three months of adherence to the measures [60]. Besides, it has been described that individuals who do not follow a prescribed diet and have a "Western diet" high in fat and red meat have elevated levels of TMAO in their urine compared to patients who follow a prescribed DASH regimen [61,62]. Reduced dietary fiber intake is associated with reduced bacterial production of short-chain fatty acid butyrate, which has immuno-modulatory effects in the gut mucosa and also serves as a major energy substrate for colonocytes. Decreased levels of butyrate in the gut could induce local inflammation, worsen dysbiosis and contribute to impaired gut barrier function, resulting in "leakage" of bacterial toxins such as LPS, which further induces local and systemic inflammation. A high-fiber diet can improve the growth of acetate-producing bacteria, reduce high blood pressure, and prevent heart fibrosis and hypertrophy [63].

Probiotics and prebiotics

Probiotics are living microorganisms which, when given in appropriate amounts, bring health benefits to the host [64]. Probiotics in clinical use contain bacterial and fungal microorganisms, including the genera *Lactobacillus* and *Bifidobacterium* and the fungus *Saccharomyces boulardii* [65]. The results of animal models suggest that certain strains of lactobacilli could have cardioprotective effects. Rats treated with a supplement containing *Lactobacillus plantarum* 299v prior to coronary artery ligation reduced infarct size and improved left ventricular function [66]. Another study showed similar cardioprotective results in a rat model of myocardial ischemia after supplementation with *Lactobacillus rhamnosus* GR-1 [67]. A pilot study in humans reported not only reduced systemic inflammation, but also improved ejection fraction after intervention with probiotic yeast *Saccharomyces Boulardii* in patients with

chronic heart failure [68]. Given the potential clinical impact of microbiota modulation, as well as high morbidity and mortality from heart failure, microbiota modulation is not completely risk-free [69]. Careful clinical monitoring and pre-defined safety measures which should follow the same standards as in other clinical trials are recommended, [70] because recent genomic and epidemiological evidence of probiotic-related bacteremia or transfer of bacteria from probiotic capsules into the blood of patients in intensive care units has been reported [71]. Prebiotics are substrates that microorganisms of the host selectively use and provide potential health benefits. Dietary fiber and oligosaccharides are most commonly used as prebiotics [72]. Most modern studies that deal with microbiota processing in patients with cardiovascular disease report microbial depletion with SCFA-forming capacity such as butyrate. Prebiotics that promote microbiological fermentation of dietary fiber in SCFA may, therefore, be of potential benefit in improving metabolic regulation [73]. Some prebiotics, such as inulin, have the potential to counteract the harmful effects of antibiotics by promoting the diversity and functional capacity of the gut microbiota [74]. A randomized study with inulin food supplement or inulin-propionate ester showed a decrease in markers of systemic inflammation with increased generation of SCFA propionate in the colon [75]. Therefore, targeting the production of microbial SCFAs with inulin supplements or other prebiotics is an attractive strategy for future cardiovascular disease testing, although current scientific evidence does not provide validated recommendations for the use of probiotics or prebiotics as adjunctive therapy in patients with heart failure or coronary heart disease.

Antibiotics

The use of antibiotics affects the composition, diversity and function of the normal flora [76]. Antibiotics have been used successfully in animal models to reduce the degree of damage to heart cells after myocardial infarction [77,78]. Previous studies in patients with heart failure focused on gut decontamination with broad-spectrum antibiotics to reduce biotome translocation and bacterial inflammation. Although this approach succeeded in reducing markers of systemic

mikrobiotom [83]. Bez obzira na razlike, ove studije na životinjama snažno ukazuju na ulogu sastava mikrobiote creva u akutnom infarktu miokarda, ali smer promena mikrobiote i potencijalni metabolički ili inflamatorni putevi još nisu dobro poznati. Modifikovanje kardiovaskularnih bolesti antibiotskom terapijom nije nova ideja. Između 1995. i 2005. godine > 19 000 pacijenata bilo je uključeno u studiju lečenja pneumonije kod pacijenata sa koronarnom bolešću. Studija kod ovih bolesnika nije pokazala kliničku korist od antibiotske terapije u vezi sa koronarnom ishemijom [84]. Pored očiglednog rizika od antimikrobne rezistencije, nedavno su se pojavili i drugi sigurnosni problemi sa potencijalnim značajem za buduća ispitivanja. Nedavno je desetogodišnje praćenje pokazalo povećanu kardiovaskularnu smrt kod pacijenata sa stabilnom koronarnom bolešću koji su lečeni klaritromicinom [85] što je dovelo do upozorenja FDA 2018. godine o upotrebi klaritromicina kod pacijenata sa koronarnom bolešću. FDA savetuje oprez pre propisivanja antibiotika klaritromicina pacijentima sa srčanim oboljenjima zbog potencijalnog povećanog rizika od srčanih problema ili smrti koji mogu nastupiti godinama kasnije. (<https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-review-finds-additional-data-supports-potential-increased-long>).

U decembru 2018. godine, objavljeno je upozorenje FDA o upotrebi fluorohinolona ukazujući na mogućnost ruptura aorte i disekcije aorte kod pacijenata sa povećanim rizikom, kao što su stariji pacijenti sa hipertenzijom ili perifernom aterosklerotskom vaskularnom bolešću

(<https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-review-finds-additional-data-supports-potential-increased-long>).

Još jedna studija izvestila je o povećanom riziku od kardiovaskularnih događaja kod starijih žena sa povećanom kumulativnom izloženošću antibioticima u odrasloj dobi [86]. Objašnjenje za ovaj povećani rizik nije do kraja kod svih pacijenata poznato, ali bi moglo uključivati produženje QT-intervalu, aritmiju ili pro-inflamatorne aktivnosti posredovane translokacijom mikrobiote creva, ili druge efekte posredovane crevnom mikrobiotom. S obzirom na ove sigurnosne probleme i nedostatak do kraja jasnog kliničkog efekta o učinku antibiotika na mikrobiotu trebalo bi pojačati oprez u budućim studijama usmerenim na

upotrebu antibiotika kod kardiovaskularnih bolesnika.

Ciljani inhibitori enzima. Pored napred navedene upotrebe TMA liaze [18] treba pomenuti rezultate studije u kojoj su miševi koristili analoge holina koji inhibiraju dejstvo enzima u metabolizmu TMA i time smanjuju koncentraciju TMAO u plazmi. Upotreba analoga holina bi, prema tome, mogla da omogući novi pristup smanjenju rizika za trombozu [69]. Još jedan zanimljiv aktivni sastojak koji deluje kao zaštitni faktor za crevnu sluznicu je Urolitin A (UroA) i njegov sintetički analog UAS03 koji poboljšavaju bliski ćelijski kontakt i funkciju gastrointestinalne barijere [87].

Fekalna mikrobna transplantacija (FMT) je poslednjih godina među najpominjanijim intervencijama koje se koriste za saniranje disbioze creva. Uvođenje „dobrih“ bakterija uzetih od zdravih ispitanika u gastrointestinalni trakt pacijenata koji pate od crevne disbioze i njenih posledica je nova i efikasna terapijska strategija. U kliničkoj studiji koja je ispitivala osobe sa metaboličkim sindromom došlo je do značajno poboljšane osetljivost na insulin posle 6 nedelja FMT u kojoj su donori bili zdrave osobe normalne telesne težine. Istovremeno FMT je povećala količinu bakterija koje proizvode butirat [88]. Iako je prihvatanje terapijske upotrebe FMT sve prisutnije, kako zbog percepcije ove metode kao „prirodnog“ lečenja i relativno jeftine primene, odnos rizika i koristi posebno u bolestima KVS ostaje nedovoljno jasno definisan jer je objavljeno iskustvo sa FMT ograničeno, a ni pravna normativa ove terapije još nije precizno uređena. Dalje, postoji strah od zaraznog potencijala terapije, što je navelo istraživače da istraže upotrebu proizvoda „sintetičke stolice“ sa definisanom populacijom bakterija da bi ublažili takve probleme, a razmatra se i upotreba „smrznutog donorskog materijala nešto poput koncepta banke matičnih ćelija [89].

Zaključak

Nova saznanja i tehnologije značajno menjaju medicinsku doktrinu i omogućavaju novi, drugačiji pogled na telo, organe i zdravlje, kao i na uzročne faktore bolesti. Istraživanja u nedavnoj prošlosti i ponekad iznenađujući nalazi potvrdili su da crevna mikrobiota može uticati na zdravlje domaćina i pokrenuti bolest različitim patofiziološkim mehanizmima. Crevna

inflammation, a clinical effect was not demonstrated [79,80].

A recent study showed that a broad-spectrum oral antibiotic cocktail significantly increased postinfarction rupture and death in a mouse model of coronary artery ligation [81], which could suggest that an intact microbial community is needed for proper myocardial recovery at the time of myocardial injury. This study is in contrast to a previous animal experimental model that showed that oral vancomycin reduced infarct size and improved postinfarction cardiac function in rats [82], and to a study reporting that a combination of streptomycin, neomycin, polymyxin B, and bacitracin reduced infarct size along with changes in metabolites associated with the microbiota [83].

Regardless of the differences, these animal studies strongly indicate the role of gut microbiota composition in acute myocardial infarction, but the direction of microbiota changes and potential metabolic or inflammatory pathways are not yet well known. Modifying cardiovascular diseases with antibiotic therapy is not a new idea.

Between 1995 and 2005 > 19.000 patients were included in a study to treat pneumonia in patients with coronary heart disease. The study in these patients showed no clinical benefit from antibiotic therapy in relation to coronary ischemia [84]. Apart from the apparent risk of antimicrobial resistance, other safety concerns have recently emerged with potential significance for future testing. A recent ten-year follow-up showed increased cardiovascular death in patients with stable coronary heart disease treated with clarithromycin [85], leading to a 2018 FDA warning on the use of clarithromycin in patients with coronary heart disease. The FDA advises caution before prescribing clarithromycin antibiotic to patients with heart disease because of a potential increased risk of heart problems or death that may occur years later. (<https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-review-finds-additional-data-supports-potential-increased-long>). In December 2018, the FDA issued a warning on the use of fluoroquinolones indicating the possibility of aortic rupture and aortic dissection in high-risk patients, such as elderly patients with hypertension or peripheral atherosclerotic vascular disease (<https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-review-finds-additional-data-supports-potential-increased-long>).

review-finds-additional-data-supports-potential-increased-long). Another study reported increased risk of cardiovascular events in older women with increased cumulative antibiotic exposure in adulthood [86]. The explanation for this increased risk is not fully known in all patients, but it could include QT prolongation, arrhythmias or pro-inflammatory activities mediated by gut microbiota translocation, or other effects mediated by the gut microbiota. Considering these safety concerns and the lack of a completely clear clinical effect antibiotics have on the microbiota, caution should be exercised in future studies targeting the use of antibiotics in cardiovascular patients.

Targeted enzyme inhibitors

Apart from the above-mentioned use of TMA lyase [18], mention should be made of the results of a study in which mice used choline analogs that inhibit the action of enzymes in TMA metabolism, thereby reducing plasma TMAO concentrations. The use of choline analogues could, therefore, provide a new approach to reducing the risk of thrombosis [69]. Another interesting active ingredient that acts as a protective factor for the gut mucosa is Urolitin A (UroA) and its synthetic analogue UAS03, which improve close cell contact and gastrointestinal barrier function [87].

In recent years, fecal microbial transplantation (FMT) has been among the most mentioned interventions used to treat intestinal dysbiosis. The introduction of "good bacteria" taken from healthy subjects into the gastrointestinal tract of patients suffering from gut dysbiosis and its consequences is a new and effective therapeutic strategy. In a clinical study examining people with metabolic syndrome, there was a significantly improved insulin sensitivity after 6 weeks of FMT in which the donors were healthy people of normal weight. At the same time, FMT increased the amount of butyrate-producing bacteria [88]. Although the acceptance of the therapeutic use of FMT is increasingly present, due to the perception of this method as a "natural" treatment and relatively cheap application, the risk-benefit ratio particularly in CVS diseases remains insufficiently clearly defined because the published experience with FMT is limited and the legal norm of this therapy has not yet been precisely regulated. Furthermore, there is a fear of the infectious potential of the therapy, which led researchers to investigate the use of "synthetic stool" products with a defined

mikrobiota i disbioza su područja istraživanja koja će u budućnosti verovatno promeniti neke danas uspostavljene metode prevencije i lečenja bolesti. Iako možemo da promenimo sastav mikrobiote pomoću prebiotika, probiotika, antibiotika, dijetete i „ciljanih inhibitora enzima“, zasad ne možemo predvideti i detaljno proceniti sve efekte ovih intervencija u prevenciji različitih bolesti. Uz sve podatke dobijene u biomedicini poslednjih decenija, čini se neobičnim da je trebalo toliko vremena pre nego što su naučnici i kardiolozi počeli sistematski da se bave uticajem 2kg mikroorganizama koji nas kolonizuju i žive sa nama „u dobru i zlu“. Iako su predstavljeni samo neki mehanizmi koji

povezuju crevnu mikrobiotu i pojedine kardiovaskularne bolesti, moramo biti svesni mogućnosti ovog istraživačkog područja u razvoju potencijalnih lekova u budućnosti. Novorazjašnjene veze između disbioze i patogeneze kardiovaskularnih bolesti nude nove mogućnosti za rano i ciljano delovanje.

P.S. Možda će nova istraživanja dovesti i do nove subspecializacije u internoj medicini koja će se (kako u svom predavanju na kongresu Internista 2019. godine anegdotski reče prof. Miodrag Krstić) zvati gastroenterokardiologija?

LITERATURA:

- Mengchao Jin Mengchao Jin, Zhiyuan Qian, Jiayu Yin, Weiting Xu, and Xiang Zhou The role of intestinal microbiota in cardiovascular disease *Journal of Cellular and Molecular Medicine* 2019; 23(4): 2343–2350.
- Sekirov I, Russell SL, Antunes LC, Finlay BB. Gut microbiota in health and disease. *Physiol Rev.* 2010;90(3):859-904. DOI: 10.1152/physrev.00045.2009 PMID: 20664075
- Tang WH, Kitai T, Hazen SL. Gut Microbiota in Cardiovascular Health and Disease. *Circ Res.* 2017;120(7):1183-96. DOI: 10.1161/CIRCRESAHA.117.309715 PMID: 28360349
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature.* 2014;505(7484):559-63. DOI: 10.1146/annurev-med-060513-093205 PMID: 25587655
- Tamburini S, Shen N, Wu HC, Clemente JC. The microbiome in early life: implications for health outcomes. *Nat Med.* 2016;22(7):713-22. DOI: 10.1038/nm.4142 PMID: 27387886
- Nallu A, Sharma S, Ramezani A, Muralidharan J, Raj D. Gut microbiome in chronic kidney disease: challenges and opportunities. *Transl Res.* 2017;179:24-37. DOI: 10.1016/j.trsl.2016.04.007 PMID: 27187743
- Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med.* 2013;368(17):1575-84. DOI: 10.1056/NEJMoa1109400 PMID: 23614584
- Ahmadmehrabi S, Tang WH. Gut microbiome and its role in cardiovascular diseases. *Curr Opin Cardiol.* 2017;32(6):761-6. DOI: 10.1097/HCO.0000000000000445 PMID: 29023288
- Yamashiro K, Tanaka R, Urabe T, Ueno Y, Yamashiro Y, Nomoto K, et al. Gut dysbiosis is associated with metabolism and systemic inflammation in patients with ischemic stroke. *PLoS One.* 2017;12(2):e0171521. DOI: 10.1371/journal.pone.0171521 PMID: 28166278
- Vinod N. The Novel Dimensions of Cardio-Metabolic Health: Gut Microbiota, Dysbiosis and its Fallouts. *Curre Res Diabetes & Obes J.* 2019;11(1):555805. DOI: 10.19080/CRDOJ.2019.11.555805
- ANTAL, I., JELIĆ, M., SILA, S., KOLAČEK, S. i TAMBIĆ ANDRAŠEVIĆ, A. LJUDSKA MIKROBIOTA I MIKROBIOM. *Acta medica Croatica*, 2019;73(1),3-11. Preuzeto sa <https://hrcak.srce.hr/>
- Lloyd-Price J, Abu-Ali G, Huttenhower C. The healthy human microbiome. *Genome Med* 2016; 8 (1):51.
- Qi Y, Aranda JM, Rodriguez V, Raizada MK, Pepine CJ. Impact of antibiotics on arterial blood pressure in a patient with resistant hypertension - A case report. *Int J Cardiol.* 2015;201:157-8. DOI: 10.1016/j.ijcard.2015.07.078 PMID: 26301638
- Gomez-Arango LF, Barrett HL, McIntyre HD, Callaway LK, Morrison M, Dekker Nitert M; SPRING Trial Group. Increased Systolic and Diastolic Blood Pressure Is Associated With Altered Gut Microbiota Composition and Butyrate Production in Early Pregnancy. *Hypertension.* 2016;68(4):974-81. DOI: 10.1161/HYPERTENSIONAHA.116.07910 PMID: 27528065
- Niebauer J, Volk HD, Kemp M, Dominguez M, Schumann RR, Rauchhaus M, et al. Endotoxin and immune activation in chronic heart failure: a prospective cohort study. *Lancet.* 1999;353(9167):1838-42. DOI: 10.1016/S0140-6736(98)09286-1 PMID: 10359409
- Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al. Metabolic Endotoxemia Initiates Obesity and Insulin Resistance. *Diabetes.* 2007;56(7):1761-72. DOI: 10.2337/db06-1491 PMID: 17456850
- Karlsson FH, Fåk F, Nookaew I, Tremaroli V, Fagerberg B, Petranovic D, et al. Symptomatic atherosclerosis is associated with an altered gut metagenome. *Nat Commun.* 2012;3(1):1245. DOI: 10.1038/ncomms2266 PMID: 23212374
- Wang Z, Roberts A.B, Buffa J.A, Levison B.S, Zhu W, Org E. et al. Non-lethal inhibition of gut microbial trimethylamine production for the treatment of atherosclerosis. *Cell.* 2015;163:1585-1595.
- Townsend MK, Aschard H, De Vivo I, Michels KB, Kraft P. Genomics, telomere length, epigenetics, and metabolomics in the nurses' health studies. *Am J Public Health* 2016;106(9): 1663-1668.
- Ju Seung Kwun, Si-Hyuck Kang, Hyo-Jung Lee, Chang-Hwan Yoon, Jung-Won Suh, Young-Seok Cho, METAGENOMIC ANALYSIS OF MICROBIOTA IN PATIENTS WITH ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 2. ORAL, GUT, AND

population of bacteria to alleviate such problems, and the use of "frozen donor material" such as the concept of a stem cell bank is being considered [89].

Concluding remarks

New knowledge and technologies are significantly changing medical doctrine, enabling a new, different view of the body, organs and health, as well as the causal factors of diseases. Research in the recent past, and sometimes surprising findings, have confirmed that the gut microbiota can affect the health of the host and trigger the disease by means of various pathophysiological mechanisms. Gut microbiota and dysbiosis are areas of research which, in the future, will likely change the established methods of prevention and treatment of diseases.

Although we can change the composition of the microbiota with prebiotics, probiotics, antibiotics, diet and "targeted enzyme inhibitors", for the time being we cannot predict and provide a detailed assessment of all

the effects of these interventions in the prevention of various diseases. With all the data obtained in biomedicine in recent decades, it seems unusual that it took so long before scientists and cardiologists began to systematically deal with the impact of 2 kg of microorganisms that colonize us and live with us "for better or worse". Although only some of the mechanisms that link the gut microbiota and certain cardiovascular diseases are presented, we must be aware of the possibilities of this research area in the development of potential drugs in the future. The newly clarified connections between dysbiosis and the pathogenesis of cardiovascular diseases offer new possibilities for early and targeted action.

P.S. Perhaps the new research will lead to a new subspecialization in internal medicine, which, as prof. Miodrag Krstićanec totally mentioned in his lecture at the Congress of Internal Medicine in 2019, will be called gastroenterocardiology.

LITERATURE:

- Mengchao Jin Mengchao Jin, Zhiyuan Qian, Jiayu Yin, Weiting Xu, and Xiang Zhou The role of intestinal microbiota in cardiovascular disease *Journal of Cellular and Molecular Medicine* 2019; 23(4): 2343–2350.
- Sekirov I, Russell SL, Antunes LC, Finlay BB. Gut microbiota in health and disease. *Physiol Rev.* 2010;90(3):859-904. DOI: 10.1152/physrev.00045.2009 PMID: 20664075
- Tang WH, Kitai T, Hazen SL. Gut Microbiota in Cardiovascular Health and Disease. *Circ Res.* 2017;120(7):1183-96. DOI: 10.1161/CIRCRESAHA.117.309715 PMID: 28360349
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature.* 2014;505(7484):559-63. DOI: 10.1146/annurev-med-060513-093205 PMID: 25587655
- Tamburini S, Shen N, Wu HC, Clemente JC. The microbiome in early life: implications for health outcomes. *Nat Med.* 2016;22(7):713-22. DOI: 10.1038/nm.4142 PMID: 27387886
- Nallu A, Sharma S, Ramezani A, Muralidharan J, Raj D. Gut microbiome in chronic kidney disease: challenges and opportunities. *Transl Res.* 2017;179:24-37. DOI: 10.1016/j.trsl.2016.04.007 PMID: 27187743
- Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med.* 2013;368(17):1575-84. DOI: 10.1056/NEJMoa1109400 PMID: 23614584
- Ahmadmehrabi S, Tang WH. Gut microbiome and its role in cardiovascular diseases. *Curr Opin Cardiol.* 2017;32(6):761-6. DOI: 10.1097/HCO.0000000000000445 PMID: 29023288
- Yamashiro K, Tanaka R, Urabe T, Ueno Y, Yamashiro Y, Nomoto K, et al. Gut dysbiosis is associated with metabolism and systemic inflammation in patients with ischemic stroke. *PLoS One.* 2017;12(2):e0171521. DOI: 10.1371/journal.pone.0171521 PMID: 28166278
- Vinod N. The Novel Dimensions of Cardio-Metabolic Health: Gut Microbiota, Dysbiosis and its Fallouts. *Curre Res Diabetes & Obes J.* 2019;11(1):555805. DOI: 10.19080/CRDOJ.2019.11.555805
- ANTAL, I., JELIĆ, M., SILA, S., KOLAČEK, S. i TAMBIĆ ANDRAŠEVIĆ, A. LJUDSKA MIKROBIOTA I MIKROBIOM. *Acta medica Croatica*, 2019;73(1),3-11. Preuzeto sa <https://hrcak.srce.hr/>
- Lloyd-Price J, Abu-Ali G, Huttenhower C. The healthy human microbiome. *Genome Med* 2016; 8 (1):51.
- Qi Y, Aranda JM, Rodriguez V, Raizada MK, Pepine CJ. Impact of antibiotics on arterial blood pressure in a patient with resistant hypertension - A case report. *Int J Cardiol.* 2015;201:157-8. DOI: 10.1016/j.ijcard.2015.07.078 PMID: 26301638
- Gomez-Arango LF, Barrett HL, McIntyre HD, Callaway LK, Morrison M, Dekker Nitert M; SPRING Trial Group. Increased Systolic and Diastolic Blood Pressure Is Associated With Altered Gut Microbiota Composition and Butyrate Production in Early Pregnancy. *Hypertension.* 2016;68(4):974-81. DOI: 10.1161/HYPERTENSIONAHA.116.07910 PMID: 27528065
- Niebauer J, Volk HD, Kemp M, Dominguez M, Schumann RR, Rauchhaus M, et al. Endotoxin and immune activation in chronic heart failure: a prospective cohort study. *Lancet.* 1999;353(9167):1838-42. DOI: 10.1016/S0140-6736(98)09286-1 PMID: 10359409
- Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al. Metabolic Endotoxemia Initiates Obesity and Insulin Resistance. *Diabetes.* 2007;56(7):1761-72. DOI: 10.2337/db06-1491 PMID: 17456850

- THROMBUS MICROBIOME IN ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 112JACC, 2020;75(11). Preuzeto 18.11.2020 sa <https://www.jacc.org/doi/pdf/10.1016/S0735-1097%2820%2930739-7>
21. Yamashiro K, Tanaka R, Urabe T, Ueno Y, Yamashiro Y, et al. Gut dysbiosis is associated with metabolism and systemic inflammation in patients with ischemic stroke. *PLoS One* 2017;12(2): e0171521.
 22. Honour J The possible involvement of intestinal bacteria in steroidal hypertension. *Endocrinology* 1982;110(1): 285-287.
 23. Yang T, Santisteban MM, Rodriguez V, Li E, Ahmari N, et al. Gut dysbiosis is linked to hypertension. *Hypertension* 2015;65(6):1331-1340.
 24. Karbach SH, Schonfelder T, Brandao I, Wilms E, Hörmann N, et al. Gut microbiota promotes angiotensin II-induced arterial hypertension and vascular dysfunction. *J Am Heart Assoc* 2016;5(9): e003698.
 25. Koh A, De Vadder F, Kovatcheva-Datchary P, Backhed F From dietary fiber to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell* 2016;165(6): 1332-1345.
 26. Rey FE, Faith JJ, Bain J, Muehlbauer MJ, Stevens RD, et al. Dissecting the in vivo metabolic potential of two human gut acetogens. *J Biol Chem* 2010;285(29): 22082-22090.
 27. Louis P, Flint HJ Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol* 2017;19(1): 29-41.
 28. Duncan SH, Barcenilla A, Stewart CS, Pryde SE, Flint HJ Acetate utilization and butyryl coenzyme A (CoA): acetate-CoA transferase in butyrate-producing bacteria from the human large intestine. *Appl Environ Microbiol* 2002;68(10): 5186-5190.
 29. Gomez-Arango LF, Barrett HL, McIntyre HD, Callaway LK, Morrison M, et al. Increased systolic and diastolic blood pressure is associated with altered gut microbiota composition and butyrate production in early pregnancy. *Hypertension* 2016;68(4): 974-981.
 30. Marques FZ, Nelson E, Chu PY, Horlock D, Fiedler A, et al. High-fiber diet and acetate supplementation change the gut microbiota and prevent the development of hypertension and heart failure in hypertensive mice. *Circulation* 2017;135(10): 964-977.
 31. Tan JK, McKenzie C, Marino E, Macia L, Mackay CR Metabolite-sensing G protein-coupled receptors-facilitators of diet-related immune regulation. *Annu Rev Immunol* 2017;35: 371-402.
 32. Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, et al. Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci U.S.A* 2013;110(11): 4410-4415.
 33. Natarajan N, Hori D, Flavahan S, Stepan J, Flavahan NA, et al. Microbial short chain fatty acid metabolites lower blood pressure via endothelial G protein-coupled receptor 41. *Physiol Genomics* 2016;48(11): 826-834.
 34. Miyamoto J, Kasubuchi M, Nakajima A, Irie J, Itoh H, et al. The role of short-chain fatty acid on blood pressure regulation. *Curr Opin Nephrol Hypertens* 2016;25(5): 379-383.
 35. Packer CS, Rice AE, Johnson TC Oxidized low density lipoprotein (OX-LDL) induced arterial muscle contraction signaling mechanisms. *Open Hyperten J* 2014;6: 20-26.
 36. Ma J, Li H. The Role of Gut Microbiota in Atherosclerosis and Hypertension. *Front Pharmacol*. 2018;9:1082. DOI: 10.3389/fphar.2018.01082 PMID: 30319417
 37. Ott SJ, El Mokhtari NE, Musfeldt M, Hellmig S, Freitag S, Rehman A, et al. Detection of diverse bacterial signatures in atherosclerotic lesions of patients with coronary heart disease. *Circulation*. 2006;113(7):929-37. DOI: 10.1161/CIRCULATIONAHA.105.579979 PMID: 16490835
 38. Lam V, Su J, Koprowski S, Hsu A, Tweddell JS, Rafiee P, et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J*. 2012;26(4):1727-35. DOI: 10.1096/fj.11-197921 PMID: 22247331
 39. Lam V, Su J, Hsu A, Gross GJ, Salzman NH, Baker JE. Intestinal Microbial Metabolites Are Linked to Severity of Myocardial Infarction in Rats. *PLoS One*. 2016;11(8):e0160840. DOI: 10.1371/journal.pone.0160840 PMID: 27505423
 40. Gan XT, Ettinger G, Huang CX, Burton JP, Haist JV, Rajapurohitam V, et al. Probiotic administration attenuates myocardial hypertrophy and heart failure after myocardial infarction in the rat. *Circ Heart Fail*. 2014;7(3):491-9. DOI: 10.1161/CIRCHEARTFAILURE.113.000978 PMID: 24625365
 41. Patterson E, Cryan J. F., Fitzgerald G. F., Ross R. P., Dinan T. G., Stanton C. Gut microbiota, the pharmabiotics they produce and host health. *Proceedings of the Nutrition Society*. 2014;73(4):477-489. doi: 10.1017/S0029665114001426.
 42. Pasini E, Aquilani R, Testa C, et al. Pathogenic gut flora in patients with chronic heart failure. *JACC: Heart Failure*. 2016;4(3):220-227. doi: 10.1016/j.jchf.2015.10.009
 43. Luedde M., Winkler T., Heinsen F. A., et al. Heart failure is associated with depletion of core intestinal microbiota. *ESC Heart Failure*. 2017;4(3):282-290. doi: 10.1002/ehf2.12155.
 44. Kamo T., Akazawa H., Suda W., et al. Dysbiosis and compositional alterations with aging in the gut microbiota of patients with heart failure. *PLoS One*. 2017;12(3):e0174099. doi: 10.1371/journal.pone.0174099.
 45. Kummén M., Mayerhofer C. C. K., Vestad B., et al. Gut microbiota signature in heart failure defined from profiling of 2 independent cohorts. *Journal of the American College of Cardiology*. 2018;71(10):1184-1186. doi: 10.1016/j.jacc.2017.12.057.
 46. Arpaia N., Campbell C., Fan X., et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature*. 2013;504(7480):451-455. doi: 10.1038/nature12726
 47. Kitai T, Kirsop J, Tang WH. Exploring the Microbiome in Heart Failure. *Curr Heart Fail Rep*. 2016;13(2):103-9. DOI: 10.1007/s11897-016-0285-9 PMID: 26886380
 48. Tang WHW, Wang Z, Fan Y, Levison B, Hazen JE, Donahue LM, et al. Prognostic Value of Elevated Levels of Intestinal Microbe-Generated Metabolite trimethylamine-N-oxide in Patients With Heart Failure: Refining the Gut Hypothesis. *J Am Coll Cardiol*. 2014;64(18):1908-14. DOI: 10.1016/j.jacc.2014.02.617 PMID: 25444145
 49. Skok P, Skok K. Prebavna cev in srčno-žilne bolezni – ali imajo kaj skupnega? *Zdrav Vestn*. 2020;89(9-10):528-38.

17. Karlsson FH, Fåk F, Nookaew I, Tremaroli V, Fagerberg B, Petranovic D, et al. Symptomatic atherosclerosis is associated with an altered gut metagenome. *Nat Commun.* 2012;3(1):1245. DOI: 10.1038/ncomms2266 PMID: 23212374
18. Wang Z, Roberts A.B, Buffa J.A, Levison B.S, Zhu W, Org E, et al. Non-lethal inhibition of gut microbial trimethylamine production for the treatment of atherosclerosis. *Cell.* 2015;163:1585-1595.
19. Townsend MK, Aschard H, De Vivo I, Michels KB, Kraft P. Genomics, telomere length, epigenetics, and metabolomics in the nurses' health studies. *Am J Public Health* 2016;106(9): 1663-1668.
20. Ju Seung Kwun, Si-Hyuck Kang, Hyo-Jung Lee, Chang-Hwan Yoon, Jung-Won Suh, Young-Seok Cho, METAGENOMIC ANALYSIS OF MICROBIOTA IN PATIENTS WITH ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 2. ORAL, GUT, AND THROMBUS MICROBIOME IN ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 112JACC, 2020;75(11). Preuzeto 18.11.2020 sa <https://www.jacc.org/doi/pdf/10.1016/S0735-1097%2820%2930739-7>
21. Yamashiro K, Tanaka R, Urabe T, Ueno Y, Yamashiro Y, et al. Gut dysbiosis is associated with metabolism and systemic inflammation in patients with ischemic stroke. *PLoS One* 2017;12(2): e0171521.
22. Honour J The possible involvement of intestinal bacteria in steroidal hypertension. *Endocrinology* 1982;110(1): 285-287.
23. Yang T, Santisteban MM, Rodriguez V, Li E, Ahmari N, et al. Gut dysbiosis is linked to hypertension. *Hypertension* 2015;65(6):1331-1340.
24. Karbach SH, Schonfelder T, Brandao I, Wilms E, Hörmann N, et al. Gut microbiota promotes angiotensin II-induced arterial hypertension and vascular dysfunction. *J Am Heart Assoc* 2016;5(9): e003698.
25. Koh A, De Vadder F, Kovatcheva-Datchary P, Backhed F From dietary fiber to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell* 2016;165(6): 1332-1345.
26. Rey FE, Faith JJ, Bain J, Muehlbauer MJ, Stevens RD, et al. Dissecting the in vivo metabolic potential of two human gut acetogens. *J Biol Chem* 2010;285(29): 22082-22090.
27. Louis P, Flint HJ Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol* 2017;19(1): 29-41.
28. Duncan SH, Barcenilla A, Stewart CS, Pryde SE, Flint HJ Acetate utilization and butyryl coenzyme A (CoA): acetate-CoA transferase in butyrate-producing bacteria from the human large intestine. *Appl Environ Microbiol* 2002;68(10): 5186-5190.
29. Gomez-Arango LF, Barrett HL, McIntyre HD, Callaway LK, Morrison M, et al. Increased systolic and diastolic blood pressure is associated with altered gut microbiota composition and butyrate production in early pregnancy. *Hypertension* 2016;68(4): 974-981.
30. Marques FZ, Nelson E, Chu PY, Horlock D, Fiedler A, et al. High-fiber diet and acetate supplementation change the gut microbiota and prevent the development of hypertension and heart failure in hypertensive mice. *Circulation* 2017;135(10): 964-977.
31. Tan JK, McKenzie C, Marino E, Macia L, Mackay CR Metabolite-sensing G protein-coupled receptors-facilitators of diet-related immune regulation. *Annu Rev Immunol* 2017;35: 371-402.
32. Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, et al. Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci U.S.A* 2013;110(11): 4410-4415.
33. Natarajan N, Hori D, Flavahan S, Stepan J, Flavahan NA, et al. Microbial short chain fatty acid metabolites lower blood pressure via endothelial G protein-coupled receptor 41. *Physiol Genomics* 2016;48(11): 826-834.
34. Miyamoto J, Kasubuchi M, Nakajima A, Irie J, Itoh H, et al. The role of short-chain fatty acid on blood pressure regulation. *Curr Opin Nephrol Hypertens* 2016;25(5): 379-383.
35. Packer CS, Rice AE, Johnson TC Oxidized low density lipoprotein (OX-LDL) induced arterial muscle contraction signaling mechanisms. *Open Hyperten J* 2014;6: 20-26.
36. Ma J, Li H. The Role of Gut Microbiota in Atherosclerosis and Hypertension. *Front Pharmacol.* 2018;9:1082. DOI: 10.3389/fphar.2018.01082 PMID: 30319417
37. Ott SJ, El Mokhtari NE, Musfeldt M, Hellmig S, Freitag S, Rehman A, et al. Detection of diverse bacterial signatures in atherosclerotic lesions of patients with coronary heart disease. *Circulation.* 2006;113(7):929-37. DOI: 10.1161/CIRCULATIONAHA.105.579979 PMID: 16490835
38. Lam V, Su J, Koprowski S, Hsu A, Tweddell JS, Rafiee P, et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J.* 2012;26(4):1727-35. DOI: 10.1096/fj.11-197921 PMID: 22247331
39. Lam V, Su J, Hsu A, Gross GJ, Salzman NH, Baker JE. Intestinal Microbial Metabolites Are Linked to Severity of Myocardial Infarction in Rats. *PLoS One.* 2016;11(8):e0160840. DOI: 10.1371/journal.pone.0160840 PMID: 27505423
40. Gan XT, Ettinger G, Huang CX, Burton JP, Haist JV, Rajapurohitam V, et al. Probiotic administration attenuates myocardial hypertrophy and heart failure after myocardial infarction in the rat. *Circ Heart Fail.* 2014;7(3):491-9. DOI: 10.1161/CIRCHEARTFAILURE.113.000978 PMID: 24625365
41. Patterson E, Cryan J. F., Fitzgerald G. F., Ross R. P., Dinan T. G., Stanton C. Gut microbiota, the pharmabiotics they produce and host health. *Proceedings of the Nutrition Society.* 2014;73(4):477-489. doi: 10.1017/S0029665114001426.
42. Pasini E., Aquilani R., Testa C., et al. Pathogenic gut flora in patients with chronic heart failure. *JACC: Heart Failure.* 2016;4(3):220-227. doi: 10.1016/j.jchf.2015.10.009
43. Luedde M., Winkler T., Heinsen F. A., et al. Heart failure is associated with depletion of core intestinal microbiota. *ESC Heart Failure.* 2017;4(3):282-290. doi: 10.1002/ehf2.12155.
44. Kamo T., Akazawa H., Suda W., et al. Dysbiosis and compositional alterations with aging in the gut microbiota of patients with heart failure. *PLoS One.* 2017;12(3):e0174099. doi: 10.1371/journal.pone.0174099.
45. Kummén M., Mayerhofer C. C. K., Vestad B., et al. Gut microbiota signature in heart failure defined from profiling of 2 independent cohorts. *Journal of the American College of Cardiology.* 2018;71(10):1184-1186. doi: 10.1016/j.jacc.2017.12.057.

50. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, et al. The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA* 2004;101(4):15718-15723.
51. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, et al. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 2006;444(7122):1027-1031.
52. Samuel BS, Shaito A, Motoike T, Rey FE, Backhed F, et al. Effects of the gut microbiota on host adiposity are modulated by the short-chain fatty-acid binding G protein-coupled receptor, Gpr41. *Proc Natl Acad Sci USA* 2008;105(43):16767-16772.
53. Bäckhed F, Manchester JK, Semenkovich CF, Gordon JI. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proc Natl Acad Sci USA* 2007;104(3):979-984.
54. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, et al. Obesity alters gut microbial ecology. *Proc Natl Acad Sci USA* 2005;102:11070-11075.
55. Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 2007;56(7):1761-1772.
56. de La Serre CB, Ellis CL, Lee J, Hartman AL, Rutledge JC, et al. Propensity to high-fat diet-induced obesity in rats is associated with changes in the gut microbiota and gut inflammation. *Am J Physiol Gastrointest Liver Physiol* 2010;299(4):G440-G448.
57. Harris K, Kassis A, Major G, Chou CJ. Is the gut microbiota a new factor contributing to obesity and its metabolic disorders? *J Obes* 2012;2012:879151. doi: 10.1155/2012/879151.
58. Jia Q, Li H, Zhou H, Zhang X, Zhang A, Xie Y, et al. Microbiota and irritable bowel syndrome: A critical inventory. *Cardiovasc Ther*. 2019;2019:5164298. DOI: 10.1155/2019/5164298 PMID: 31819762
59. Salehi-Abargouei A, Maghsoudi Z, Shirani F, Azadbakht L. Effects of Dietary Approaches to Stop Hypertension (DASH)-style diet on fatal or nonfatal cardiovascular diseases—incidence: a systematic review and meta-analysis on observational prospective studies. *Nutrition*. 2013;29(4):611-8. DOI: 10.1016/j.nut.2012.12.018 PMID: 23466047
60. Rifai L, Pisano C, Hayden J, Sulo S, Silver MA. Impact of the DASH diet on endothelial function, exercise capacity, and quality of life in patients with heart failure. *Proc Bayl Univ Med Cent*. 2015;28(2):151-6. DOI: 10.1080/08998280.2015.11929216 PMID: 25829641
61. Lopez-Garcia E, Rodriguez-Artalejo F, Li TY, Fung TT, Li S, Willett WC, et al. The Mediterranean-style dietary pattern and mortality among men and women with cardiovascular disease. *Am J Clin Nutr*. 2014;99(1):172-80. DOI: 10.3945/ajcn.113.068106 PMID: 24172306
62. De Filippis F, Pellegrini N, Vannini L, Jeffery IB, La Storia A, Laghi L, et al. High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut*. 2016;65(11):1812-21. DOI: 10.1136/gutjnl-2015-309957 PMID: 26416813
63. M. Trøseid et al. The gut microbiome in coronary artery disease and heart failure: Current knowledge and future directions [Review]. *2020;52:102649*. Published: February 18, 2020 DOI: <https://doi.org/10.1016/j.ebiom.2020.102649>
64. Hill C, Guarner F, Reid G, Gibson G.R., Merenstein D.J., Pot B, et al. Expert consensus document: the international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*. 2014;11:506-514.
65. Patel R, DuPont H.L. New approaches for bacteriotherapy: prebiotics, new-generation probiotics, and synbiotics. *Clin Infect Dis*. 2015; 60: S108-S121.
66. Lam V, Su J, Koprowski S, Hsu A, Tweddell J.S, Rafiee P, et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J*. 2012; 26: 1727-1735.
67. Gan X.T, Ettinger G, Huang C.X, Burton J.P, Haist J.V, Rajapurohitam V, et al. Probiotic administration attenuates myocardial hypertrophy and heart failure after myocardial infarction in the rat. *Circ Heart Fail*. 2014; 7: 491-499.
68. Costanza A.C, Moscovitch S.D, Faria Neto H.C, Mesquita E.T. Probiotic therapy with *saccharomyces boulardii* for heart failure patients: a randomized, double-blind, placebo-controlled pilot trial. *Int J Cardiol*. 2015; 179: 348-350.
69. He Z, Wang J, Chen Y, Cong X, Li N, Ding R, et al. Potential risk associated with direct modulation of the gut flora in patients with heart failure. *ESC Heart Fail*. 2019; 6: 555-556.
70. Mayerhofer C.C.K, Awoyemi A, Hov J.R, Troseld M, Broch K. Reply: potential risk associated with direct modulation of the gut flora in patients with heart failure. *ESC Heart Fail*. 2019; 6: 557-558.
71. Yelin I, Flett K.B, Merakou C, Mehrotra P, Stam J, Sniesrud E, et al. Genomic and epidemiological evidence of bacterial transmission from probiotic capsule to blood in ICU patients. *Nat Med*. 2019; 25: 1728-1732.
72. Gibson G.R, Hutkins R, Sanders M.E, Prescott S.L, Reimer R.A, Salminen S.J, et al. Expert consensus document: the international scientific association for probiotics and prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*. 2017;14:491-502.
73. Chambers E.S, Preston T, Frost G, Morrison D.J. Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Curr Nutr Rep*. 2018; 7: 198-206.
74. Johnson L.P, Walton G.E, Psichas A, Frost G.S, Gibson G.R, Barraclough T.G. Prebiotics modulate the effects of antibiotics on gut microbial diversity and functioning in vitro. *Nutrients*. 2015; 7: 4480-4497.
75. Chambers E.S, Byrne C.S, Morrison D.J, Murphy K.G, Preston T, Tedford C, et al. Dietary supplementation with inulin-propionate ester or inulin improves insulin sensitivity in adults with overweight and obesity with distinct effects on the gut microbiota, plasma metabolome and systemic inflammatory responses: a randomised cross-over trial. *Gut*. 2019; 68: 1430-1438.
76. Lin PP, Hsieh YM, Kuo WW, Lin YM, Yeh YL, Lin CC, et al. Probiotic-fermented purple sweet potato yogurt activates compensatory IGF1R/PI3K/Akt survival pathways and attenuates cardiac apoptosis in the hearts of spontaneously hypertensive rats. *Int J Mol Med*. 2013;32(6):1319-28. DOI: 10.3892/ijmm.2013.1524 PMID: 24127171
77. Tiihonen K, Tynkkynen S, Ouwehand A, Ahlroos T, Rautonen N. The effect of ageing with and without non-steroidal anti-inflammatory drugs on gastrointestinal microbiology and immunology. *Br J*

46. Arpaia N., Campbell C., Fan X., et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature*. 2013;504(7480):451–455. doi: 10.1038/nature12726
47. Kitai T, Kirsop J, Tang WH. Exploring the Microbiome in Heart Failure. *Curr Heart Fail Rep*. 2016;13(2):103–9. DOI: 10.1007/s11897-016-0285-9 PMID: 26886380
48. Tang WHW, Wang Z, Fan Y, Levison B, Hazen JE, Donahue LM, et al. Prognostic Value of Elevated Levels of Intestinal Microbe-Generated Metabolite trimethylamine-N-oxide in Patients With Heart Failure: Refining the Gut Hypothesis. *J Am Coll Cardiol*. 2014;64(18):1908–14. DOI: 10.1016/j.jacc.2014.02.617 PMID: 25444145
49. Skok P, Skok K. Prebavna cev in srčno-žilne bolezni – ali imajo kaj skupnega? *Zdrav Vestn*. 2020;89(9–10):528–38.
50. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, et al. The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA* 2004;101(4):15718–15723.
51. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, et al. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 2006;444(7122): 1027–1031.
52. Samuel BS, Shaito A, Motoike T, Rey FE, Backhed F, et al. Effects of the gut microbiota on host adiposity are modulated by the short-chain fatty-acid binding G protein-coupled receptor, Gpr41. *Proc Natl Acad Sci USA* 2008;105(43): 16767–16772.
53. Bäckhed F, Manchester JK, Semenkovich CF, Gordon JI. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proc Natl Acad Sci USA* 2007;104(3): 979–984.
54. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, et al. Obesity alters gut microbial ecology. *Proc Natl Acad Sci USA* 2005;102: 11070–11075.
55. Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 2007;56(7):1761–1772.
56. de La Serre CB, Ellis CL, Lee J, Hartman AL, Rutledge JC, et al. Propensity to high-fat diet-induced obesity in rats is associated with changes in the gut microbiota and gut inflammation. *Am J Physiol Gas- trointest Liver Physiol* 2010;299(4): G440–G448.
57. Harris K, Kassis A, Major G, Chou CJ. Is the gut microbiota a new factor contributing to obesity and its metabolic disorders?. *J Obes* 2012;2012: 879151. doi: 10.1155/2012/879151.
58. Jia Q, Li H, Zhou H, Zhang X, Zhang A, Xie Y, et al. Microbiota and irritable bowel syndrome: A critical inventory. *Cardiovasc Ther*. 2019;2019:5164298. DOI: 10.1155/2019/5164298 PMID: 31819762
59. Salehi-Abargouei A, Maghsoudi Z, Shirani F, Azadbakht L. Effects of Dietary Approaches to Stop Hypertension (DASH)-style diet on fatal or nonfatal cardiovascular diseases—incidence: a systematic review and meta-analysis on observational prospective studies. *Nutrition*. 2013;29(4):611–8. DOI: 10.1016/j.nut.2012.12.018 PMID: 23466047
60. Rifai L, Pisano C, Hayden J, Sulo S, Silver MA. Impact of the DASH diet on endothelial function, exercise capacity, and quality of life in patients with heart failure. *Proc Bayl Univ Med Cent*. 2015;28(2):151–6. DOI: 10.1080/08998280.2015.11929216 PMID: 25829641
61. Lopez-Garcia E, Rodriguez-Artalejo F, Li TY, Fung TT, Li S, Willett WC, et al. The Mediterranean-style dietary pattern and mortality among men and women with cardiovascular disease. *Am J Clin Nutr*. 2014;99(1):172–80. DOI: 10.3945/ajcn.113.068106 PMID: 24172306
62. De Filippis F, Pellegrini N, Vannini L, Jeffery IB, La Stora A, Laghi L, et al. High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut*. 2016;65(11):1812–21. DOI: 10.1136/gutjnl-2015-309957 PMID: 26416813
63. M. Trøseid et al. The gut microbiome in coronary artery disease and heart failure: Current knowledge and future directions Review| 2020;52:102649. Published: February 18, 2020 DOI:https://doi.org/10.1016/j.ebiom.2020.102649
64. Hill C, Guarner F, Reid G, Gibson G.R., Merenstein D.J., Pot B. et al. Expert consensus document. the international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*. 2014;11:506–514.
65. Patel R. DuPont H.L. New approaches for bacteriotherapy: prebiotics, new-generation probiotics, and synbiotics. *Clin Infect Dis*. 2015; 60: S108–S121.
66. Lam V, Su J, Koprowski S, Hsu A, Tweddell J.S. Rafiee P. et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J*. 2012; 26: 1727–1735.
67. Gan X.T. Ettinger G. Huang C.X. Burton J.P. Haist J.V. Rajapurohitam V. et al. Probiotic administration attenuates myocardial hypertrophy and heart failure after myocardial infarction in the rat. *Circ Heart Fail*. 2014; 7: 491–499.
68. Costanza A.C. Moscovitch S.D. Faria Neto H.C. Mesquita E.T. Probiotic therapy with *saccharomyces boulardii* for heart failure patients: a randomized, double-blind, placebo-controlled pilot trial. *Int J Cardiol*. 2015; 179: 348–350.
69. He Z. Wang J. Chen Y. Cong X. Li N. Ding R. et al. Potential risk associated with direct modulation of the gut flora in patients with heart failure. *ESC Heart Fail*. 2019; 6: 555–556.
70. Mayerhofer C.C.K. Awoyemi A. Hov J.R. Trøseid M. Broch K. Reply: potential risk associated with direct modulation of the gut flora in patients with heart failure. *ESC Heart Fail*. 2019; 6: 557–558.
71. Yelin I. Flett K.B. Merakou C. Mehrotra P. Stam J. Snesrud E. et al. Genomic and epidemiological evidence of bacterial transmission from probiotic capsule to blood in ICU patients. *Nat Med*. 2019; 25: 1728–1732.
72. Gibson G.R. Hutkins R. Sanders M.E. Prescott S.L. Reimer R.A. Salminen S.J. et al. Expert consensus document: the international scientific association for probiotics and prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*. 2017;14:491–502.
73. Chambers E.S. Preston T. Frost G. Morrison D.J. Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Curr Nutr Rep*. 2018; 7: 198–206.
74. Johnson L.P. Walton G.E. Psichas A. Frost G.S. Gibson G.R. Barraclough T.G. Prebiotics modulate the effects of antibiotics on gut microbial diversity and functioning in vitro. *Nutrients*. 2015; 7: 4480–4497.
75. Chambers E.S. Byrne C.S. Morrison D.J. Murphy K.G. Preston T. Tedford C. et al. Dietary supplementation

- Nutr. 2008;100(1):130-7. DOI: 10.1017/S000711450888871X PMID: 18279548
78. Zhou X, Li J, Guo J, Geng B, Ji W, Zhao Q, et al. Gut-dependent microbial translocation induces inflammation and cardiovascular events after ST-elevation myocardial infarction. *Microbiome*. 2018;6(1):66. DOI: 10.1186/s40168-018-0441-4 PMID: 29615110
 79. Conraads V.M. Jorens P.G. De Clerck L. Van Saene H.K. Ieven M.M. et al. Selective intestinal decontamination in advanced chronic heart failure: a pilot trial. *Eur J Heart Fail*. 2004; 6: 483-491.
 80. Fox M.A. Peterson S. Fabri B.M. van Saene H.K. Selective decontamination of the digestive tract in cardiac surgical patients. *Crit Care Med*. 1991; 19: 1486-1490.
 81. Tang T.W.H. Chen H.C. Chen C.Y. Yen C.Y.T. Lin C.J. Prajnamitra R.P. et al. Loss of gut microbiota alters immune system composition and cripples postinfarction cardiac repair. *Circulation*. 2019; 139: 647-659.
 82. Lam V. Su J. Koprowski S. Hsu A. Tweddell J.S. Rafiee P. et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J*. 2012; 26: 1727-1735.
 83. Lam V. Su J. Hsu A. Gross G.J. Salzman N.H. Baker J.E. Intestinal microbial metabolites are linked to severity of myocardial infarction in rats. *PLoS ONE*. 2016; 11:e0160840.
 84. Andraws R. Berger J.S. Brown D.L. Effects of antibiotic therapy on outcomes of patients with coronary artery disease: a meta-analysis of randomized controlled trials. *JAMA*. 2005; 293: 2641-2647.
 85. Winkel P. Hilden J. Hansen J.F. Kastrup J. Kolmos H.J. Kjoller E. et al. Clarithromycin for stable coronary heart disease increases all-cause and cardiovascular mortality and cerebrovascular morbidity over 10 years in the claricor randomised, blinded clinical trial. *Int J Cardiol*. 2015; 182: 459-465.
 86. Knoop K.A. McDonald K.G. Kulkarni D.H. Newberry R.D. Antibiotics promote inflammation through the translocation of native commensal colonic bacteria. *Gut*. 2016; 65: 1100-1109.
 87. Colman RJ, Rubin DT. Fecal microbiota transplantation as therapy for inflammatory bowel disease: a systematic review and meta-analysis. *J Crohns Colitis* 2014;8(12): 1569-1581.
 88. Vrieze A, Van Nood E, Holleman F, Salojärvi J, Kootte RS, et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 2012;143(4): 913-916.
 89. Petrof E, Gloor G, Vanner S, Weese SJ, Carter D, et al. Stool substitute transplant therapy for the eradication of *Clostridium difficile* infection: 'repopulating' the gut. *Microbiome* 2013;1(1): 3.

- with inulin-propionate ester or inulin improves insulin sensitivity in adults with overweight and obesity with distinct effects on the gut microbiota, plasma metabolome and systemic inflammatory responses: a randomised cross-over trial. *Gut*. 2019; 68: 1430-1438.
76. Lin PP, Hsieh YM, Kuo WW, Lin YM, Yeh YL, Lin CC, et al. Probiotic-fermented purple sweet potato yogurt activates compensatory IGFIR/PI3K/Akt survival pathways and attenuates cardiac apoptosis in the hearts of spontaneously hypertensive rats. *Int J Mol Med*. 2013;32(6):1319-28. DOI: 10.3892/ijmm.2013.1524 PMID: 24127171
 77. Tiihonen K, Tynkkynen S, Ouwehand A, Ahlroos T, Rautonen N. The effect of ageing with and without non-steroidal anti-inflammatory drugs on gastrointestinal microbiology and immunology. *Br J Nutr*. 2008;100(1):130-7. DOI: 10.1017/S000711450888871X PMID: 18279548
 78. Zhou X, Li J, Guo J, Geng B, Ji W, Zhao Q, et al. Gut-dependent microbial translocation induces inflammation and cardiovascular events after ST-elevation myocardial infarction. *Microbiome*. 2018;6(1):66. DOI: 10.1186/s40168-018-0441-4 PMID: 29615110
 79. Conraads V.M. Jorens P.G.De Clerck L.S.van Saene H.K. Ieven M.M. et al. Selective intestinal decontamination in advanced chronic heart failure: a pilot trial. *Eur J Heart Fail*. 2004; 6: 483-491.
 80. Fox M.A.Peterson S. Fabri B.M. van Saene H.K. Selective decontamination of the digestive tract in cardiac surgical patients. *Crit Care Med*. 1991; 19: 1486-1490.
 81. Tang T.W.H. Chen H.C. Chen C.Y. Yen C.Y.T. Lin C.J. Prajnamitra R.P. et al. Loss of gut microbiota alters immune system composition and cripples postinfarction cardiac repair. *Circulation*. 2019; 139: 647-659.
 82. Lam V. Su J. Koprowski S. Hsu A. Tweddell J.S. Rafiee P. et al. Intestinal microbiota determine severity of myocardial infarction in rats. *FASEB J*. 2012; 26: 1727-1735.
 83. Lam V. Su J. Hsu A. Gross G.J. Salzman N.H. Baker J.E. Intestinal microbial metabolites are linked to severity of myocardial infarction in rats. *PLoS ONE*. 2016; 11:e0160840.
 84. Andraws R. Berger J.S. Brown D.L. Effects of antibiotic therapy on outcomes of patients with coronary artery disease: a meta-analysis of randomized controlled trials. *JAMA*. 2005; 293: 2641-2647.
 85. Winkel P. Hilden J. Hansen J.F. Kastrup J. Kolmos H.J. Kjoller E. et al. Clarithromycin for stable coronary heart disease increases all-cause and cardiovascular mortality and cerebrovascular morbidity over 10 years in the claricor randomised, blinded clinical trial. *Int J Cardiol*. 2015; 182: 459-465.
 86. Knoop K.A. McDonald K.G. Kulkarni D.H. Newberry R.D. Antibiotics promote inflammation through the translocation of native commensal colonic bacteria. *Gut*. 2016; 65: 1100-1109.
 87. Colman RJ, Rubin DT. Fecal microbiota transplantation as therapy for inflammatory bowel disease: a systematic review and meta-analysis. *J Crohns Colitis*. 2014;8(12): 1569-1581.
 88. Vrieze A, Van Nood E, Holleman F, Salojärvi J, Kootte RS, et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology*. 2012;143(4): 913-916.
 89. Petrof E, Gloor G, Vanner S, Weese SJ, Carter D, et al. Stool substitute transplant therapy for the eradication of *Clostridium difficile* infection: 'repopulating' the gut. *Microbiome*. 2013;1(1): 3.