Primljen/ Received on 8.5.2013.
Revidiran/ Revised on 14.5.2013.
Prihvaćen/ Accepted on 20.5.2013.

CLINICAL
ARTICLE

doi: 10.5937/asn1367255M

ENVIRONMENTAL AND GLOVES' CONTAMINATION BY STAPHYLOCOCCI IN DENTAL HEALTHCARE SETTINGS

Giuseppe A. Messano¹, Virgilio De Bono², Renato Architrave¹, Stefano Petti¹

¹SAPIENZA UNIVERSITY, DEPARTMENT OF PUBLIC HEALTH AND INFECTIOUS DISEASES, ROME, ITALY;

²ITALIAN MEDICAL ASSOCIATION, ROMAN SECTION, ROME, ITALY

Abstract

Introduction. Each year, 37,000 people in Europe die as direct consequence of healthcare-associated infections. *Staphylococcus aureus* (especially methicillin-resistant -MRSA) and coagulase-negative staphylococci (CNS) are frequently isolated in these episodes. Environmental contamination by *S. aureus*, MRSA and CNS in dental healthcare settings is reported moderately frequently, although the associated risk for infection is not clear.

Aim. To investigate contamination of disposal gloves and of clinical contact surfaces by several types of staphylococci in dental offices soon after dental therapy.

Material and methods. 136 general dental practitioners (GDPs) voluntarily participated. At each sampling occasion, environmental samples were collected from the tray and from the gloved dominant hand, soon after dental therapy of the second or third patient of the working session. Contact plates containing Mannitol Salt Agar were used. Overall staphylococci, *S. aureus*, CNS and *Staphylococcus epidermidis* (member of CNS group) were presumptively identified and resistance to oxacillin was tested to identify methicillin-resistant (MR) strains.

Results. Staphylococci were detected in 41% and 57% samples from trays and from gloves, respectively; *S. aureus* in 5% and 5%, CNS in 36% and 52%, *S. epidermidis* in 18% and 44%, methicillin-resistant *S. aureus* (MRSA) in 1.5% and 1.5%, MR-CNS in 1.5% and 2.2%, MR-*S. epidermidis* in 1.5% and 1.5%. The samples collected from the trays were correlated with those collected from hands for all these types of staphylococci.

Conclusion. Although it was not possible to ascertain the main source of staphylococci contamination -patient or GDP, dominant hands and clinical contact surfaces were frequently contaminated.

Key words: dentistry, clinical contact surface, gloves, general dental practitioner, *Staphylococcus aureus*, coagulase-negative staphylococci, *Staphylococcus epidermidis*

Introduction

Healthcare-associated infections (HAI) occur after exposure to healthcare. Each year, 4,000,000 people in Europe acquire HAIs and 37,000 of them die as the direct consequence of infection. In addition, HAIs may also interest healthcare workers and administrative staff. The most frequent HAI types are urinary tract infections, pneumonia, surgical site infections, bloodstream infections and gastrointestinal infections, while the most frequently isolated microorganisms in HAI overall are *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterococcus* species, coagulase-negative staphylococci and *Candida* species¹. HAIs are generally treated through antibiotic and/or surgical therapy. Therefore, HAI treatment and prevention is an important cause of emergence of highly antibiotic resistant strains, another serious public health problem².

In the specific field of dental healthcare settings, environmental contamination is reported for methicillin-resistant and methicillin-sensitive *S. aureus* (MRSA and MSSA), although the risk for infection associated with contamination is not clear³⁻⁵. One problem with MRSA in dentistry is that these micro-organisms may survive up to six months on clinical contact surfaces in the dental environment⁶. *P. aeruginosa* and *Legionella pneumophila*, microorganisms responsible for HAIs, also are frequently de-

Address for correspondence:

Dr. Giuseppe Alessio Messano
Department of Public Health and Infectious Diseases
Sapienza University
P.le Aldo Moro 5, 00185 Rome, Italy
Phone/Fax: +3906 4991 4667
Email: giuseppe.messano@yahoo.it

© 2013 Faculty of Medicine in Niš. Clinic of Dentistry in Niš.
All rights reserved / © 2013 Medicinski fakultet Niš. Klinika za
stomatologiju Niš. Sva prava zadržana

tected in dental unit waterlines, along with oral streptococci, biological markers of bloodborne/airborne pathogens⁷⁻⁹. The study of environmental contamination and consequent infection transmission in dental healthcare settings is, therefore, essential to produce evidence-based guidelines¹⁰⁻¹².

Aim

The aim of the present study was to investigate the contamination level of the environment of dental offices and of the hands of the dental staff by several types of staphylococci.

Material and Methods

One hundred thirty-six General Dental Practitioners (GDPs) working in private offices in Rome, Italy, were considered. Details regarding their recruitment were previously described¹³; GDPs were invited to participate before registering to Continuing Medical Education courses. Participation was on a voluntary basis and there were no incentives. Data protection and anonymity were guaranteed. The study protocol was approved by the Review Board of the Medical and Dental Association of Rome.

Samples were collected in the midmorning/midafternoon. GDPs must have treated at least two patients before the sampling occasion. Environmental samples were collected from the tray in front of the patient, a clinical contact surface¹⁰, immediately after patient treatment and before surface cleaning, using disposable swabs previously imbibed in sterile water and passed on a 10x10 cm area of the tray. ATP bioluminescence (Lumicontrol II, PBI International, Milan, Italy) was used to assess the overall contamination level, qualitatively. ATP bioluminescence is a rapid method to detect the total viable flora, with a reasonably good accuracy of 114% at levels around 100 Relative Light Units (RLUs), roughly corresponding to 2.5 colony forming units (CFU)/cm². 100 RLUs is generally considered the benchmark for high level of environmental cleanliness in hospitals^{14,15}.

Samples which yielded RLU levels >100 were monitored using Replicate Organism Detection and Counting (Rodac) plates (Becton Dickinson Italia, Buccinasco, Italy) containing

Mannitol Salt Agar (MSA - Becton Dickinson Italia), for the enumeration of staphylococci. Plates were pressed on the surface of the tray different from the area sampled with the swab for 30 s at an approximate pressure of 20-25 g/cm². Rodac plates were preferred to swab because they provide repeatable results and provide comparable levels of recovery for general bacterial contamination of a surface¹⁶. Plates were aerobically incubated at 37°C for 48 h. Colonies grown on MSA were Gram's stained, tested for coagulase and catalase and presumptively identified using VITEK-2 "Gram-Positive Identification" and "Antibiotic Susceptibility Testing" cards (BioMérieux, Italia; Bagno a Ripoli, Italy). Further biochemical identification tests were not made, therefore, micro-organisms were presumptively classified as overall staphylococci, *S. aureus*, coagulase negative staphylococci (CNS) and *S. epidermidis*, a subgroup of CNS.

Following the criteria of the US Centers for Disease Control, isolated staphylococci strains were classified as methicillin-resistant on the basis of oxacillin MIC test, which implied the presence of the staphylococcal cassette chromosome *mec* (SCC*mec*). Namely, ≥ 4 µg/mL for *S. aureus* and ≥ 0.5 µg/mL for CNS (guidelines available at, <http://www.cdc.gov/mrsa/lab/lab-detection.html>).

Right hand samples among right-handed GDPs (from left hand among left-handed GDPs) were collected directly from the gloved hand of GDPs soon after patient treatment and before glove removal using four Rodac plates containing the aforementioned media and following the same laboratory procedures used for the environmental samples.

Staphylococci prevalence estimates on clinical contact surfaces and on GDPs' gloved dominant hands were assessed. The association between clinical contact surface and gloved hand contamination by the various Staphylococcus species also was assessed using the non-parametric Spearman correlation test, giving score 1 to positive samples and score 0 to negative samples.

GDPs were informed in the event that samples were positive for methicillin-resistant staphylococci. Environmental samples and samples from GDPs' nares were collected using sterile swabs one week after the sampling occasion to check whether additional procedures

were necessary to eradicate these microorganisms.

Results

Seventy-six (55.9%) clinical contact surfaces provided RLU levels higher than 100. Overall presumptive staphylococci prevalence was 41.2% (n=56, 73.7% of all ATP bioluminescence positive samples), with 5.1% *S. aureus* (n=7) and 36.0% (n=49) CNS, of whom 18.4% (n=25) *S. epidermidis* (Table 1). Two presumptive *S. aureus* species, corresponding to 1.5%

S. epidermidis (1.5%) were methicillin resistant (Table 2).

A good correlation was found between positive environmental samples and positive samples from gloves for overall staphylococci, *S. aureus* and CNS (Table 1; $p < 0.0001$). As for methicillin-resistant staphylococci, all the MRSA (n=2) were isolated from the environmental sample and on GDP's glove at the same time, while one of four CNS (*S. epidermidis*) was isolated from both samples (Table 2). In the three sampling occasions where methicillin-resistant staphylococci were isolated from the tray and from the glove of the dominant hand at

Table 1. Prevalence (95% confidence interval -95CI- between square brackets) of samples with ATP bioluminescence levels >100 Relative Light Units (RLU -corresponding to 2.5 CFU/cm²) and of the investigated staphylococci detected immediately after dental therapy on the 136 clinical contact surfaces and GDPs' gloved hands. Association between staphylococci detected on clinical contact surfaces and staphylococci detected on GDP's gloved hands (Spearman correlation coefficient ρ -rho- corrected for ties).

	>100 RLUs	staphylococci	<i>S. aureus</i>	CNS	<i>S. epidermidis</i>
clinical contact surfaces	55.9% (n=76) [47.6-64.2]	41.2% (n=56) [32.9-49.5]	5.1% (n=7) [1.4-8.8]	36.0% (n=49) [27.9-44.1]	18.4% (n=25) [11.9-24.9]
gloved hands		57.4% (n=78) [49.1-65.7]	5.1% (n=7) [1.4-8.8]	52.2% (n=71) [43.8-60.6]	44.1% (n=60) [35.8-52.4]
Spearman ρ		0.66 ($p < 0.0001$)	0.70 ($p < 0.0001$)	0.72 ($p < 0.0001$)	0.31 ($p = 0.0004$)

CNS: coagulase negative staphylococci (the species *S. epidermidis* is a member of CNS)

of all samples (3.6% of the environmental detected staphylococci, 28.6% of *S. aureus* isolates), resulted methicillin resistant. Two *S. epidermidis* isolates (1.5% of all samples, 3.6% of staphylococci isolates) were methicillin resistant (Table 2).

the same time, the two microorganisms exhibited the same antibiotic profile, suggesting that the strain detected on the tray was the same as the strain detected from the glove (data not in Table).

Table 2. Prevalence (95% confidence interval -95CI- between square brackets) of methicillin-resistant staphylococci isolated from the 136 clinical contact surfaces and the GDPs' gloved hands.

	staphylococci	<i>S. aureus</i>	CNS	<i>S. epidermidis</i>
clinical contact surfaces	2.9% (n=4) [<0.1-5.7]	1.5% (n=2) [<0.0-3.5]	1.5% (n=2) [<0.0-3.5]	1.5% (n=2) [<0.0-3.5]
gloved hands	3.7% (n=5) [0.5-6.9]	1.5% (n=2) [<0.0-3.5]	2.2% (n=3) [<0.0-4.7]	1.5% (n=2) [<0.0-3.5]

The two methicillin-resistant *S. aureus* isolates and two out of 3 methicillin-resistant CNS isolates (one *S. epidermidis*) were isolated on the clinical contact surface and on the GDP's gloved hand at the same sampling occasions.

Overall staphylococci prevalence in samples from gloves soon after dental therapy was 57.4% (n=78), *S. aureus* accounted for 5.1% (n=7) of the samples while the remaining 52.2% (n=71) were CNS, including 44.1% (n=60) *S. epidermidis* (Table 1). Two presumptive *S. aureus* species, corresponding to 1.5% of samples (2.6% of the detected staphylococci, 28.6% of all *S. aureus* isolates), were methicillin resistant. Three CNS (2.2%) and, among them, two

In the six dental offices where methicillin-resistant staphylococci were detected, environmental samples, collected one week following the first sampling occasion, provided negative results regarding the occurrence of methicillin-resistant staphylococci. Samples from GDPs' nares also resulted negative.

Discussion

The present study is one of the papers presented at the workshop “Advances in Infection Epidemiology and Control in Dental Healthcare Settings”, Department of Public Health and Infectious Diseases, Sapienza University, Rome, Italy on February 9th, 2013¹⁷⁻²³.

MRSA infection transmission is a serious public health problem and a cause for concern among dental healthcare workers and patients²⁴. While MRSA infection can be transmitted to patients and to healthcare workers particularly in certain settings, such as intensive care or surgical units²⁵, there are no ascertained cases of transmission to dental healthcare workers and MRSA carriage rates among GDPs are generally similar to or even lower than the general population, thus suggesting that the occupational risk of MRSA infection is probably minimal³. The question regarding the risk of acquiring MRSA infection among dental patients is more complex. Among special patients, such as special care patients, hospitalized patients, head and neck cancer patients and oral or maxillofacial surgery patients, high MRSA carriage rates are reported, as well as frequent episodes of infection and colonization, other than environmental contamination of clinics and units. These elements suggest that the risk of infection is high among these patients. The situation among the remaining dental patients is different, as there are only two reported episodes of MRSA infection, one of them due to the lack of glove use by the dentist. Carriage rate among healthy adult dental patients also is low, while there was only one case of MRSA environmental detection in non-special and non-surgical dental departments or offices^{3,4}. These elements collectively suggest that the risk for infection is generally low.

The present study showed that MRSA spread in dental healthcare settings is possible. Indeed, MRSA were detected in two out of the 136 sampling occasions collected soon after dental therapy and in these occasions both trays and gloves from the dominant hand resulted contaminated by these microorganisms at the same time. Although no further analysis was made to ascertain whether the strains detected on the gloves and those detected on the trays

were the same, it is likely that it was so, because the strain detected from the glove exhibited the same antibiotic profile as the strain detected from the tray. This assumption, along with the reported high correlation between staphylococci, CNS, *S. aureus* and *S. epidermidis* detected in the environment and those detected on the gloves may suggest two possible hypotheses to explain MRSA spread in the dental offices and the consequent risk for infection. First hypothesis: GDPs were MRSA carriers and picked these microorganisms from their skin/mucosae while they were wearing gloves, spreading them in the environment. Second hypothesis: the dental patient under treatment was MRSA carrier and was touched by GDP's gloved dominant hands, which in turn, contaminated the tray in front of the patient or, alternatively, airborne MRSA from carrier patient contaminated the tray. In favour of the second hypothesis the fact that GDPs resulted free from MRSA one week after MRSA detection in their offices, therefore, these two GDPs could only be transient carriers. Dispersion of airborne staphylococci during dental therapy also is corroborated by studies reporting CNS prevalence of 10-35% in air samples from dental clinics and offices²⁶⁻²⁸ a similar detection rate as that reported in the present study (Table 1).

Conclusion

In conclusion, environmental contamination by methicillin-resistant and methicillin-sensitive staphylococci during dental therapy is possible and may pose a risk for infection, particularly among immune-depressed patients and those with open lesions in mouth. However, it seems likely that routine glove change between patients and cleaning/disinfection of clinical contact surfaces could be sufficient to control such a risk for infection, but to be sure that these simple methods are applied, it is necessary that the overall level of knowledge and awareness among dental healthcare workers is improved, perhaps through specific Continuing Medical Education courses, as issues regarding infection control are generally neglected among dental healthcare workers²⁹.

LITERATURA / REFERENCES

1. European Centre for Disease Prevention and Control. Surveillance of healthcare-associated infections in Europe, 2007. Stockholm: ECDC; 2012.
2. Pittet D, Allegranzi B, Sax H et al. Considerations for a WHO European strategy on health-care-associated infection, surveillance, and control. *Lancet Infect Dis* 2005;5(4):242-50.
3. Petti S, Polimeni A. Risk of methicillin-resistant *Staphylococcus aureus* transmission in the dental health-care setting: a narrative review. *Infect Control Hosp Epidemiol* 2011;32(11):1109-15.
4. Roberts MC, Soge OO, Horst JA, Ly KA, Milgrom P. Methicillin-resistant *Staphylococcus aureus* from dental school clinic surfaces and students. *Am J Infect Control* 2011;39(8):628-32.
5. Petti S, Polimeni A. Methicillin-resistant *Staphylococcus aureus* infection transmission in dental health care settings: myths and facts. *Am J Infect Control* 2012;40(3):287-8.
6. Petti S, De Giusti M, Moroni C, Polimeni A. Long-term survival curve of methicillin-resistant *Staphylococcus aureus* on clinical contact surfaces in natural-like conditions. *Am J Infect Control* 2012;40(10):1010-2.
7. Walker JT, Bradshaw DJ, Finney M et al. Microbiological evaluation of dental unit water systems in general dental practice in Europe. *Eur J Oral Sci* 2004;112(5):412-8.
8. O'Donnell MJ, Boyle MA, Russell RJ, Coleman DC. Management of dental unit waterline biofilms in the 21st century. *Future Microbiol* 2011;6(10):1209-26.
9. Petti S, Moroni C, Messano GA, Polimeni A. Detection of oral streptococci in dental unit water lines after therapy with air turbine handpiece: biological fluid retraction more frequent than expected. *Future Microbiol* 2013;8(3):413-21.
10. Kohn WG, Collins AS, Cleveland JL et al. Guidelines for infection control in dental health-care settings, 2003. *MMWR Recomm Rep* 2003;52(RR-17):1-61.
11. Petti S, Polimeni A. The rationale of guidelines for infection control in dentistry: precautionary principle or acceptable risk? *Infect Control Hosp Epidemiol* 2010;31(12):1308-10.
12. Laheij AM, Kistler JO, Belibasakis GN, Välimaa H, de Soet JJ; European Oral Microbiology Workshop (EOMW) 2011. Healthcare-associated viral and bacterial infections in dentistry. *J Oral Microbiol* 2012. doi: 10.3402/jom.v4i0.17659.
13. Messano GA, Petti S. General dental practitioners and hearing impairment. *J Dent* 2012;40(10):821-8.
14. Lewis T, Griffith C, Gallo M, Weinbren M. A modified ATP benchmark for evaluating the cleaning of some hospital environmental surfaces. *J Hosp Infect* 2008;69(2):156-63.
15. Mulvey D, Redding P, Robertson C, Woodall C, Kingsmore P, Bedwell D, Dancer SJ. Finding a benchmark for monitoring hospital cleanliness. *J Hosp Infect* 2011;77(1):25-30.
16. Galvin S, Dolan A, Cahill O, Daniels S, Humphreys H. Microbial monitoring of the hospital environment: why and how? *J Hosp Infect* 2012;82(1):143-51.
17. Messano GA, Sofan AAA, Petti S. Quality of air and water in dental healthcare settings during professional toothcleaning. *Acta Stomatol Naissi* 2013; 29(67): 1230-35; doi: 10.5937/asn1367230M.
18. Messano GA, Masood M, Palermo P, Petti S. Predictors of *Legionella* occurrence in dental unit waterlines of a highly colonized dental hospital. *Acta Stomatol Naissi* 2013; 29(67): 1236-41; doi: 10.5937/asn1367236M.
19. Messano GA, Masood M, Palermo P, Petti S. Prevalence of reactive tuberculin skin test in dental healthcare workers and students. *Acta Stomatol Naissi* 2013; 29(67): 1242-48; doi: 10.5937/asn1367242M.
20. Petti S, Sofan AAA, Messano GA. *Streptococcus pneumoniae* carriage rate in healthy preadolescent dental patients. *Acta Stomatol Naissi* 2013; 29(67): 1249-54; doi: 10.5937/asn1367249P.
21. Messano GA. Bacterial and fungal contamination of dental hygienists' hands with and without finger rings. *Acta Stomatol Naissi* 2013; 29(67): 1260-64; doi: 10.5937/asn1367260M.
22. Petti S, Messano GA, Polimeni A, Dancer SJ. Effect of cleaning and disinfection on naturally contaminated clinical contact surfaces. *Acta Stomatol Naissi* 2013; 29(67): 1265-72; doi: 10.5937/asn1367265P.
23. Petti S, Messano GA, Scully C. Antibody level and immunity against Hepatitis B virus infection among general dental practitioners. *Acta Stomatol Naissi* 2013; 29(67): 1273-78; doi: 10.5937/asn1367273P.
24. Klevens RM, Gorwitz RJ, Collins AS. Methicillin-resistant *Staphylococcus aureus*: a primer for dentists. *J Am Dent Assoc* 2008;139(10):1328-37.
25. Albrich WC, Harbarth S. Health-care workers: source, vector, or victim of MRSA? *Lancet Infect Dis* 2008;8(5):289-301.
26. Rautemaa R, Nordberg A, Wuolijoki-Saaristo K, Meurman JH. Bacterial aerosols in dental practice - a potential hospital infection problem? *J Hosp Infect* 2006;64(1):76-81.
27. Szymańska J, Dutkiewicz J. Concentration and species composition of aerobic and facultatively anaerobic bacteria released to the air of a dental operation area before and after disinfection of dental unit waterlines. *Ann Agric Environ Med* 2008;15(2):301-7.
28. Kimmerle H, Wiedmann-Al-Ahmad M, Pelz K, Wittmer A, Hellwig E, Al-Ahmad A. Airborne microbes in different dental environments in comparison to a public area. *Arch Oral Biol* 2012;57(6):689-96.
29. Petti S, Messano GA, Polimeni A. Dentists' awareness toward vaccine preventable diseases. *Vaccine* 2011;29(45):8108-12.