Maja Račić¹, Vedrana R. Joksimović¹,
Bojan N. Joksimović¹, Srebrenka Kusmuk¹,
Ljilja Kozomara²

¹School of Medicine,
University East Sarajevo, Bosnia and Herzegovina

²Health Center Banja Luka,
Bosnia and Herzegovina

Alcohol consumption among the elderly primary health care patients: The impact of alcohol on overall health

Key words:

alcoholism,
screening,
elderly population,
comprehensive geriatric assessment

Abstract

Introduction: Alcoholism is the third most common psychiatric disorder among the elderly, yet it often goes undiagnosed in primary care setting.

Objective: To estimate the prevalence of alcohol use disorders among elderly primary care patients in Bosnia and Herzegovina and to determine the presence of different health problems related to alcohol consumption.

Method: The study was conducted in 10 family medicine practices. Family physicians randomly selected a group of 40 patients over 60 years of age registered with their practice. As a part of Comprehensive Geriatric Assessment, patients were asked to complete Alcohol Consumption Screening Questionnaire, Functional Status Questionnaire, Geriatric Depression Scale (GDS), Mini mental screening examination (MMSE), and Hamilton Anxiety Scale. Screening results were followed by additional clinical evaluation. To estimate the feasibility of Alcohol Consumption Screening Questionnaire, one family member or a caregiver of each patient was questioned about patient's habits and health problems.

Results: Eighty-nine (22%) patients were current drinkers. Of those, 59% were harmful drinkers, 26% hazardous and 15% nonhazardous drinkers. Women accounted for 27% of current drinkers. MMSE revealed dementia symptoms in 37%, and symptoms of mild cognitive impairment (MCI) in 25% of current drinkers. Depression symptoms were found in 38% and anxiety symptoms in 6% of current drinkers. Functional status was decreased in hazardous and harmful drinkers.

Conclusion: High percentage of older adults in Bosnia and Herzegovina is regular users of alcohol. Hazardous and harmful drinking is associated with significant morbidity. These findings demonstrate the usefulness and importance of the excess alcohol use screening in all primary care settings serving adults over age 60.

Introduction

Alcoholism is the third most common psychiatric disorder among older adults. Up to 16% of men and 8% of women have alcohol use disorders¹.

Also, alcoholism is in the differential diagnosis, either as a cause or exacerbating factor, in majority of geriatric syndromes. It is estimated that 10% of demented individuals suffer from an alcohol-induced dementia which often goes unrecognized, despite being one of the most responsive to treatment².

Alcohol abuse often coexists with depression or anxiety and increases risk of osteoporosis, falls and fractures³.

Despite the fact that many textbooks and research results talk about alcohol abuse among the elderly, it is still happening that elderly patients are treated symptomatically for alcohol-related conditions without recognition of the underlying problem. Why?

Partially, problem is in the fact that people have different attitudes toward alcohol consumption. There is continued debate, even among the physicians, regarding the beneficial effects of low-dose drinking. Many elderly people think that alcohol consumption improves overall health status. However, evidence of the benefits from drinking is still unclear, but according to some researchers, even low levels of consumption can cause adverse health effects because of age-related physiologic changes and the interaction between alcohol, ageing, impaired functional status and medication use⁴. Sometimes, patients withhold information because of shame or fear of stigmatization (particularly women). This could lead to missed information about medical and psychiatric conditions, unexpected alcohol withdrawal symptoms, drug interactions, and lost opportunities for prevention what can cause serious complications^{5,6}.

The other reason might be a worldwide tendency for primary care physicians to neglect or to be unaware of symptoms and signs of alcohol abuse. In Bosnia and Herzegovina, development of Family Medicine Departments brought the series of teaching sections regarding alcohol abuse during undergraduate and postgraduate studies, but still, a large number of physicians have inappropriate attitudes toward the screening. Very often, physicians find that available screening procedures take too much of their time and that screening measures are not sensitive enough to detect the alcoholism among the elderly, particularly of female gender. Sometimes, physicians do not have confidence to search for alcohol consumption in their elderly patients, especially when this is considered to be culturally inappropriate⁷.

Also, behaviours that lead to diagnosis in young people are considered to be less likely occurring in older persons and many of the classical clues are mistakenly attributed to age related changes or disease common in old age¹.

Current standardized screening questionnaires are gold

standards for diagnosis, but most of them have been designed to detect abusive and dependant drinking. The CAGE questionnaire⁸ and the Geriatric version of the Michigan Alcoholism Screening Test (MAST)⁵ are widely used screening measures, which can identify abusive (e.g. failure to fulfill social obligations) and dependent (e.g. having withdrawal symptoms) drinking. However, these instruments do not distinguish recent from remote drinking behaviour, and among patients aged 60 years and older, the CAGE screen is insensitive with usual scoring for detecting binge drinking. Also, those two questionnaires do not provide information about the relation between alcohol use and functional status or overall health. Therefore, supplemental information about the current quantity, frequency and pattern of alcohol use should be obtained^{9,10}.

The Alcohol Use Disorders Identification Test (The AUDIT) is another frequently used screening questionnaire, which has been shown to detect hazardous and harmful drinking in younger people¹¹. Yet, it does not provide information on alcohol use and health or on nonhazardous drinking, what can be important for clinical understanding of potential beneficial effects of low alcohol consumption on certain diseases prevention, such as stroke, dementia or cardiovascular diseases^{12,13,14}.

The Alcohol-Related Problems Survey (The ARPS) is another recently developed tool. It is an 18-item self administered screening measure that focuses on the relation between alcohol use and medical problems, medication use and functional status. ARPS classifies drinking as non-hazardous, hazardous or harmful, according to algorithms developed from combining patients' responses^{15,16}. According to the study results, the ARPS and its short version (sharps) are quite sensitive in identifying older drinkers with a spectrum of alcohol use disorders. Sensitivity and specificity of the ARPS are 93% and 63%, what makes it more sensitive than the AUDIT and the SMAST-G in identifying older persons who may be at risk or experiencing harm as a result of their alcohol use¹⁷. They also provide information on specific risks associated with alcohol use, what cannot be obtained by other screening questionnaires.

However, our experience showed that asking questions regarding alcohol consumption was culturally unacceptable in our country, what instigated us to improve screening methods and make it easier in primary care settings. According to the previous studies, we made Alcohol Consumption Screening Questionnaire which includes the questions about quantity and frequency of alcohol drinking, symptoms of alcohol abuse and alcoholism, remote drinking, attitudes toward alcohol consumption and presence of alcoholism in the family. Further one, we wanted to assess the connection between alcohol consumption and different mental health problems, elderly abuse and functional status of geriatric patients.

Method

Before the study was conducted, Alcohol Consumption Screening Questionnaire was tested with a sample of 112 older adults. Terminology understanding, ways of administration and validity were analyzed. The results showed that Alcohol Consumption Screening Questionnaire is valid alcohol consumption screening tool, understandable to all elderly people regardless of education level, and that the best way of using it comes when questionnaire is administered by physician and incorporated in Comprehensive Geriatric Assessment.

Participants

Participants were recruited from ten family medicine practices, placed in eastern parts of Bosnia and Herzegovina. Selection criteria included: unknown alcohol consumption status and unknown/undocumented mental disorders. Family physicians randomly chose a group of 40 patients, who were 60 years of age or older and who were visiting them for different reasons.

Procedure

During first study consultation, physicians explained the importance of Comprehensive Geriatric assessment and made the plan, together with the patient, how to provide it gradually during next few consultations.

On the same occasion, physicians have done complete medical assessment, medication revision and functional status assessment, using Functional Status Questionnaire.

During the second consultation, depression and dementia screening were done, using Short Geriatric Depression Scale (GDS) and Mini Mental Screening Examination (MMSE).

During the third consultation, Alcohol Consumption Screening Questionnaire was administered to the patient by physician. According to the answers, participants were classified as abstinent, nonhazardous, hazardous and harmful drinkers.

Participants who have not been taking any alcohol in last five years were classified as abstinent. Definition of nonhazardous, hazardous and harmful drinking was derived by an expert panel that used standardized panel agreement methods to reach consensus on the classification scheme⁴. Nonhazardous drinking is a type of drinking that does not result in risks of medical or psychosocial damage. Hazardous drinking is a type of drinking that indicates risks for future problems. Harmful drinking is defined by the presence of health problems that can be worsened by alcohol use and includes alcohol abuse or dependence.

Home visit was a fourth consultation. Physician or nurse was estimating social functioning, environment factors, compliance.

On the same occasion, caregiver or family member was interviewed separately. She/he was asked about the patient's alcohol consumption habits, potential mental disorders, family history and other health problems that a patient might experience. During fifth consultation, anxiety screening was done, using Hamilton Anxiety Scale (HAS). Data collection took place from December 2010-April 2011.

Statistical Analysis

Descriptive analysis in the form of frequencies and percentages was used to view the sample and the formation of certain categories of patients. Chi-square was used to test associations between age, gender and category of drinking.

The data were analyzed using SPSS program, version 20 (Statistical Program for Social Sciences).

Results

Of the 400 patients who had a screening for alcohol consumption, 182 (45%) were male and 218 (55%) were female. 208 (52%) had primary school education, while 104 (26%) had the basic level of education. 180 (45%) patients belonged to the age group between 60-69, 169 (42%) to the age group between 70-79 and 51 (13%) to the age group above 80. Most of the patients were coming from semi rural environment (78%), (Table1).

Table 1. Distribution of geriatric patients according to gender, age, education and living environment (N=400)

Characteristics	Number	Percent
Gender		
Male	182	45
Female	218	55
Age		
60-69	180	45
70-79	169	42
≥80	51	13
Education		
4 th grade of primary school	68	17
Complete primary school	208	52
High school	104	26
University degree	20	5
Environment		
Rural	65	16
Semirural	312	78
Urban	23	6

Alcohol consumption screening revealed that 89 (22%) geriatric patients were current drinkers and 311 (78%) were abstinent. We found out that 29 (7.25%) current drinkers drink daily, and that 37 (9.25%) drink 2-3 times per week. 37 (9%) patients were drinking 3-4 drinks per occasion, and 16 (4%) were drinking >5 drinks per occasion, (Table 2).

Table 2. Alcohol consumption screening results in patients (N=400)

Alcohol consumption	Number of patients	Percent
Frequency		
abstinence	311	78
1 time a month or less	13	3
2-4 time a month	10	2.5
2-3 times a week	37	9.25
Every day	29	7.25
Number of drinks		
1 or less	336	84
2	11	3
3-4	37	9
>5	16	4

According to the quantity and frequency of alcohol consumption, current drinkers were divided into three categories: nonhazardous, hazardous and harmful drinkers.

The data derived from patients responses and data given by caregiver were completely the same in the groups of hazardous and nonhazardous drinkers, while the harmful drinkers gave false negative results in 2% of cases. Fifty-three (59%) patients were classified as harmful drinkers, 23 (26%) as hazardous and 13 (15%) as nonhazardous. Statistically significant difference in the drinking category between age groups was not found ($\chi^2=4,909$; $DF=2$; $p=0,086$). Also, there was no statistically significant difference in terms of age of patients ($\chi^2=4,957$; $DF=4$; $p=0,292$), (Table 3).

Table 3. Distribution of nonhazardous, hazardous and harmful drinking according to gender and age (n=89)

Drinking classification	Men (N=65) No (%)	Women (N=24) No (%)
Nonhazardous (n=13)		
60-69	4 (6)	2 (8)
70-79	2 (3)	3 (12.5)
≥ 80	1 (2)	1 (4)
Hazardous (n=23)		
60-69	3 (5)	5 (21)
70-79	10 (15)	3 (12.5)
≥ 80	2 (3)	0
Harmful (n=53)		
60-69	15 (23)	6 (25)
70-79	27 (41)	4 (17)
≥ 80	1 (2)	0

Attitudes toward alcohol consumption were different among geriatric patients. Two hundred twenty-eight (57%) patients stated that alcohol consumption is bad for the health, 100 (25%) reported that alcohol's effects vary with the quantity or frequency of drinking. Seventy-two (18%) geriatric patients stated that alcohol consumption has very good influence on overall health, whatever or quantity or frequency (all from harmful drinking category).

Of 311 abstinent, 28 (9%) were taking alcohol during some part of their life. 21 (75%) were nonhazardous drinkers, 6 (21%) hazardous drinkers, and 1 (4%) was harmful drinker who quitted alcohol use 15 years ago.

Family history of alcoholism was positive in 76 (19%) geriatric patients. Of those, family history was positive in 24 (31,6%) abstinent and 52 (68,4%) current drinkers, of those 39 (75%) harmful drinkers and 13 (25%) hazardous drinkers. Many current drinkers had medical conditions that could be affected by alcohol consumption, such as hypertension (31%), congestive heart failure (15%) and ulcer disease (6%), (Table 4).

Table 4. Chronic medical diseases in patients who consume alcohol (N=89)

Disease	Number of patients No (%)
Hypertension	28 (31)
Osteoarthritis	11 (12)
Congestive heart failure	13 (15)
Diabetes	12 (13)
Asthma	7 (8)
Ulcer disease	5 (6)
Without chronic disease	13 (15)

Almost 24 (27%) of current drinkers were taking antihypertensive medication, 10 (11%) sedatives and 7 (8%) pain killers, such as nonsteroidal anti-inflammatory agents, (Table 5).

Table 5. Medication use in patients who consume alcohol (N=89)

Medication	Number of patients No (%)
Antihypertensives	24 (27)
Sedatives	10 (11)
NSAID	7 (8)
Antacids	10 (11)
H ₂ blockers	12 (14)
None	26 (29)

Mini Mental Screening Examination revealed that 33 (37%) current drinkers had the symptoms of dementia, while 22 (25%) had the symptoms of mild cognitive impairment.

If we analyze total number of harmful drinkers, we can see that 27 (62%) male harmful drinkers had symptoms of either dementia or mild cognitive impairment. Also, 8 (80%) female harmful drinkers had the positive score on MMSE for dementia, and 2 (20%) for mild cognitive impairment, (Table 6).

Table 6. Dementia screening results in current drinkers (N=89)

Depression screening positively revealed depression symptoms in 34 (38%) current drinkers. Of those, mild depression symptoms were present in 7 (21%) cases, moderate depression symptoms in 18 (53%), and severe depression symptoms in 9 (26%). Depression symptoms were found in 15 (35%) male harmful drinkers and 7 (70%) female harmful drinkers, (Table 7).

Table 7. Depression screening results in current drinkers (N=89)

Anxiety was experienced by 5 (6%) current drinkers.

Functional status screening showed that 213 (68%) abstinent, 10 (80%) nonhazardous drinkers, 11 (48%) hazardous drinkers and 12 (23%) of harmful drinkers had good score in the area of physical functioning. In the area of psychological functioning, 203 (65%) abstinent, 10 (80%) nonhazardous drinkers, 15 (65%) hazardous drinkers and 8 (15%) harmful drinkers had good score. In the area of role functioning, harmful drinking group had good score in 274 (42%) of cases, while in other groups, good score was present

in >88% of cases. Social functioning score was in the warning zone in almost 32 (60%) harmful drinkers. Also, harmful drinkers had warning score in the area of interaction in almost 39(74%) cases. Statistically significant differences were found in every area of functioning between the drinking categories, (Table 8).

Table 8. Functional status screening in geriatric patients (N=400)

Functional status measures	Drinking classification			
	Abstinence (N=311) No (%)	Non-hazardous (N=13) No (%)	Hazardous (N=23) No (%)	Harmful (N=53) (P value) No (%)
Physical function				
Warning zone	98 (32)	3 (20)	12 (52)	41 (77) 43,337
Good	213 (68)	10 (80)	11 (48)	12 (23) (0,000)
Psychological function				
Warning zone	109 (35)	3 (20)	8 (35)	45 (85) 48,784
Good	202 (65)	10 (80)	15 (65)	8 (15) (0,000)
Role function				
Warning zone	37 (12)	1 (8)	3 (10)	31 (58) 68,037
Good	274 (88)	12 (92)	20 (90)	22 (42) (0,000)
Social function				
Warning zone	78 (25)	3 (20)	5 (22)	32 (60) 54,485
Good	233 (75)	10 (80)	18 (78)	21 (40) (0,000)
Quality of interaction				
Warning zone	62 (20)	1 (8)	5 (21)	39 (74) 69,406
Good	249 (80)	12 (92)	18 (79)	14 (26) (0,000)

Discussion

This study showed that a significant number of older adults in BiH consume alcohol at harmful level, what has an immense impact on their overall health and should raise the awareness of general practitioner on this issue. In international studies, the prevalence of harmful drinking among the elderly was lower^{18,19}.

Alcohol consumption Screening Questionnaire appeared to be valid for use in primary health care. Yet, to avoid false negative results and to provide complete assessment of patients with possible alcohol-related problems, informa-

tion had to be gathered from several resources, including the patient's medical records and collateral informants, such as family member or caregiver.

Majority of patients who were drinking with certain risk presented with medical problems not obviously due to drinking. It included hypertension, presented in 31% of harmful and hazardous drinkers, which could have been secondary and potentially reversible. Also, hypertension in those two groups was more resistant to a treatment, compared to the hypertension in a group of nonhazardous drinkers or in abstinent group. Standard treatment of ulcer disease did not lead to the complete healing of ulcer in majority of participants who were harmful drinkers. A large percentage of patients were on medication that could be adversely affected by alcohol abuse. Alcohol adversely affected adherence to treatment.

Family history was positive in 58% of current drinkers, what shows that having an alcoholic parent present an important risk factor for developing alcoholism or alcohol abuse. Although environmental and interpersonal factors are important, a genetic predisposition underlies alcohol abuse, particularly in harmful drinking pattern²⁰.

This research showed that in abstinent group depression symptoms were also prevalent (32%), but in less percentage than in the groups of harmful and hazardous drinkers (41%). Also, in 87% of abstinent, depression symptoms were mild compared to the groups of harmful and hazardous drinkers, where 79% of depression symptoms were classified as moderate or severe. The extent of comorbidity between depression and alcoholism was demonstrated by the results of several large epidemiological studies. Although the association between major depression and alcoholism is well established, the reason for this association is not completely understood. Possible causes of the association may be shared etiology, the direct result of the pharmacology of heavy alcohol consumption, or an indirect link between the outcomes of alcohol abuse that lead to increased risk factors for depression. It is now known that some of the systems that are involved in producing the symptoms of low mood, anxiety, poor sleep and reduced appetite in depression are also affected by alcohol. Also, alcohol can have a toxic effect on their serotonin neuro-transmitters and contributes to the development of depression, increasing the frequency and severity of depression episodes, but not all heavy drinkers will become depressed, as we were able to see in this study results^{21, 22, 23, 24, 25}.

The results of the research conducted in Centre for Addiction and Mental Health, London, Ontario Department of Psychology, University of Western Ontario, London, Ontario, Canada, and presented by Agnes Massak and Kathryn Graham, using the Composite International Diagnostic Interview (CIDI) and Alcohol Use Disorders Identification Test (AUDIT), found that depression was more strongly and more generally related to hazardous drinking pattern for women than for men, but alcohol dependence symptoms and harmful con-

sequences from drinking were significantly associated with depression for both men and women.²⁶ However, establishing the association between major depression and alcoholism in geriatric patient, especially women, should provide physicians who care for these patients with a better understanding of the dual diagnoses, precaution and better plan for treatment.

People suffering from depression will often turn to alcohol in an attempt to make themselves feel better. The same is happening to the elderly people who have history of abuse. All types of abuse lead to emotional distress, and sometimes abused people cope with it. In some other cases, they can't see a solution for their problem, and start to drink in order to feel better. Further one, long-term alcohol abuse could upset chemical balances in the brain and thus could promote the onset of depressive episodes. If a patient is already depressed, the affect that heavy drinking has on the central nervous system will be more detrimental to the wellbeing of depressed patients than non depressed individuals, so all primary care patients, who are predisposed to depression, should be advised to abstain from drinking²⁷.

A large percentage of geriatric patients in abstinent group had a positive score on MMSE for dementia (usually vascular) or mild cognitive impairment (52,4%). This finding could come from the fact that 62% of study participants appeared to have insufficient or basic education (one of the risk factors for dementia). However, excessive drinking over a period of years might lead to an alcohol dementia in a group of harmful drinkers, causing problems with memory, learning and other cognitive skills.

In the study conducted by Mukamal KJ, Kuller LH and coworkers, it was found that compared with abstinence, consumption of 1 to 6 drinks weekly is associated with a lower risk of incident dementia among older adults²⁸. According to other studies, alcohol has a direct affect on brain cells, resulting in poor judgment, difficulty making decisions and lack of insight, and lowers initial repetition, recall, and recognition scores, especially when associated with depression. Current drinkers suffering from dementia, had little ability to learn new things, but few of them kept other mental abilities still highly functioning, what perhaps misled the physicians in establishing proper diagnosis^{29, 30}. Eventually, alcohol-induced dementia is reversible condition, where cessation of drinking leads to the improvement in cognition and function, and for that reason, physicians should think about the alcohol abuse as a potential cause of cognitive impairments, and do the screening of alcohol consumption among elderly patients in every day practice.

Results of the study of alcoholism, based on animal models, indicated that alcohol acts as anxiolytic agent during early stages of consumption and as an anxiogenic agent when absent after chronic or acute doses, and that the administration of ethanol provokes anxiolytic effects³¹. That could be an explanation why some current drinkers experienced anxiety

symptoms (6%).

Functional status was in decline in a group of hazardous and harmful drinkers. Perhaps, explanation could be analyzed through the understanding of alcohol effect on overall health. Alcohol leads to the anxiety, mood disorder, polyneuropathy, confusion, dizziness and drowsiness, which can influence the functional status. On the contrary, nonhazardous drinking did not cause any particular decline comparing to the abstinent group, what can follow other study results, which indicate that patients who drink infrequently and low-quantity of alcohol (nonhazardous drinkers) have better overall functional status and health than patients from other consumption groups³².

The fact the majority of harmful and hazardous drinkers come from rural areas is very important. It could be explained by the tradition and customs in rural areas of Bosnia, where domestic alcohol production is freely engaged into, without any legal repercussions, as the majority of the social events in such areas, like celebrations of Saint Patron's days and feasts, with large alcohol consumption are socially acceptable. Current drinkers from rural areas had also specific attitudes toward alcohol consumption, where a distinguishing between alcohol use and alcohol dependence or abuse in reality does not exist. Such group in particular should be a target for health promotion in family practice settings.

Conclusion

Alcohol abuse and alcoholism are common but under-recognized problems among elderly population in Bosnia and Herzegovina. Hazardous and harmful drinking was associated with significant morbidity. These findings demonstrate the usefulness and importance of the excess alcohol use screening in all primary care settings serving adults over age 60. Several practical screening tools for alcoholism are available but, supplemental information about the current quantity, frequency and pattern of alcohol use should be obtained

When caring for elderly patients who have problems related to alcohol use, family physician should encounter interrelated medical, behavioural, social and environmental factors and must maintain high index of suspicious and non-judgmental attitude.

Маја Рачић¹, Ведрана Р. Јоксимовић¹,
Бојан Н. Јоксимовић¹, Сребренка Кусмук¹,
Љиља Козомара²

¹Медицински факултет Фоча,

Универзитет Источно Сарајево, Босна и Херцеговина

²Дом здравља Бања Лука, Босна и Херцеговина

Конзумирање алкохола код старијих пацијената примарне здравствене заштите: Утицај алкохола на целокупно здравље

Кључне речи:

алкохолизам,

скрининг,

популација старијег животног доба,

свеобухватна геријатријска процена

Сажетак

Увод. Алкохолизам се налази на трећем месту најчешћих психијатријских поремећаја код особа старијег животног доба, али га лекари примарне здравствене заштите често не дијагностикују.

Циљ рада. Проценити преваленцију поремећаја изазваних употребом алкохола код пацијената старијег животног доба, који се лече на нивоу примарне здравствене заштите у Босни и Херцеговини и утврдити присуство различитих здравствених проблема у вези са конзумирањем алкохола.

Метод. Студија је спроведена у 10 тимова породичне медицине. Љекари породичне медицине су насумично одабрали групу од 40 пацијената старијих од 60 година, регистрованих у њиховом тиму породичне медицине. Као део свеобухватне геријатријске процене, пацијенти су замољени да испуне скрининг упитник о конзумирању алкохола (*Alcohol Consumption Screening Questionnaire*), упитник о процени функционалног статуса (*Functional Status Questionnaire*), геријатријску скалу депресије (*Geriatric Depression Scale - GDS*), мини ментал скрининг тест (*Mini mental screening examination - MMSE*) и Хамилтонову скалу анксиозности. Како би се проценила тачност скрининг упитника о конзумирању алкохола, члану породице или неговатељу сваког пацијента су постављена питања о пацијентовим навикама и здравственим проблемима.

Резултати. Осамдесет девет (22%) пацијената редовно конзумира алкохол. Од тога, код 59% се радило о штетном, код 26% хазардном и код 15% нехазардном конзумирању алкохола; 27% особа које редовно конзумирају алкохол су женског пола. Мини ментал скрининг тестом пронађени су симптоми деменције код 37% особа, симптоми благог когнитивног поремећаја код 25%, симптоми депресије код 38% и симптоми анксиозности код 6% особа које редовно конзумирају алкохол. Код особа које хазардно и штетно конзумирају алкохол, функционални статус је био снижен.

Закључак. Висок проценат особа старијег животног доба у Босни и Херцеговини редовно конзумира алкохол. Хазардно и штетно пијење је удружено са значајним морбидитетом. Ови резултати указују на корисност и важност скрининга прекомјерне употребе алкохола у свим здравственим установама примарног нивоа које збрињавају здравствене проблеме особа старијих од 60 година.

Literatura References

- Campbell JW. *Use of Alcohol, Tobacco & Nonprescribed Drugs*. In: CS. Landefeld et al, editors. *Current Geriatric Diagnosis and Treatment*. New York: McGraw Hill; 2004. p. 407-413.
- Enoch M, Goldman D. *Problem drinking and alcoholism: diagnosis and treatment*. *Am Fam Physician* 2002; 65(3):441.
- Yuan Z, et al. *Effects of alcohol related disease on hip fracture and mortality: a retrospective cohort study of hospitalized Medicare beneficiaries*. *Am J Public Health* 2001;91:1089.
- Moore AA, Morton SC, Beck JC, et al. *A new paradigm for alcohol use in older persons*. *Med Care* 1999;37:165-79.
- Blow FC. *Treatment of older women with alcohol problems: meeting the challenge for a special population*. *Alcohol Clin Exp Res* 2000;24(8):1257-66.
- Fielling DA, Reid MS, O'Connor PG. *Screening for alcohol problems in primary care: a systematic review*. *Arch Intern Me*. 2000;160:1977-89.
- Dufour M, Fuller RK. *Alcohol in the elderly*. *Annu Rev Med* 1995;46:123-32.
- Hinshin CH, Castellon SA, Dickson-Fuhrman E. *Screening for drug and alcohol abuse among older adults using a modified version of the CAGE*. *Am J Addict* 2001;10:319.
- Adams WL, Berry KL, Fleming MF. *Screening for problem drinking in older primary care patients*. *JAMA* 1996; 276:1964-67.
- Hays RD, Merz JF, Nicholas R. *Response burden, reliability, and validity of the CAGE, Short MUST and Audit alcohol screening measures*. *Behav Res Meth Instrum Comput* 1995;27:277-80.
- Graham K. *Identifying and measuring alcohol abuse among the elderly: serious problem with existing instrumentation*. *J Stud Alcohol* 1986; 47:322-6.
- Klatsky AL, Armstrong MA, Friedman GD. *Alcohol and mortality*. *Ann Intern Med* 1992;117:464-54.
- Thun MJ, Peto R, Lopez AD, et al. *Alcohol consumption and mortality among middle aged and elderly US adults*. *N Engl J Med* 1997;337:1705-14.
- Sacco RL, Elkind M, Boden-Albala B, et al. *The Protective Effect of Moderate Alcohol Consumption on Ischemic Stroke*. *JAMA* 1999; 281:53-60.
- Nguyen K, Fink A, Beck J, Higa J. *Feasibility of Using an Alcohol-Screening and Health Education System with older primary care patients*. *J Am Board Fam Pract* 2001;14:7-15.
- Fink A, Morton S, Beck JC, et al. *The Alcohol-Related Problems Survey: Identifying Hazardous and Harmful Drinking in Older Primary Care Patients*. *J Am Geriatr Soc* 2002; 50(10):1717-22.
- Moore AA, Beck JC, Babor TF. *Beyond alcoholism: identifying older, at-risk drinkers in primary care*. *J Stud Alcohol*. 2002; 316-24.
- Mirand AL, Welte JW. *Alcohol consumption among the elderly in a general population, Erie County, New York*. *Am J Public Health* 1996 ; 86(7): 978-984.
- Reid MC, Anderson PA. *Geriatric substance use disorders*. *Med Clin North Am* 1997;81: 999-1016
- Enoch MA, Goldman D. *Genetics of alcoholism and substance use*. *Psychiatr Clin North Am* 1999; 21:370-4.
- Spaner D, Bland RC, Newman SC. *Major depressive disorder*. *Acta Psychiatr Scand* 1994;77:7-15.
- Grant BF, Harford TC. *Comorbidity between DSM-IV alcohol use disorders and major depression: results of a national survey*. *Drug Alcohol Depend* 1995;39:197-206.
- Ross HE. *DSM-III-R alcohol abuse and dependence and psychiatric comorbidity in Ontario: results from the mental health supplement to the Ontario Health Survey*. *Drug Alcohol Depend* 1995;39:111-28.
- Kessler RC, Nelson CB, McGonagle KA, Liu J, Swartz M, Blazer DG. *Comorbidity of DSM-III-R major depressive disorder in the general population: results from the US National Comorbidity Survey*. *Br J Psychiatry* 1996;168(30):7-30.
- Kessler RC, Crum RM, Warner LA, Nelson CB, Schulenberg J, Anthony JC. *Lifetime co-occurrence of DSM-III-R alcohol abuse and dependence with other psychiatric disorders in the national Comorbidity Survey*. *Arch Gen Psychiatry* 1997; 54:313-24.
- Massak A, Graham K. *The relationship between depression, alcohol use disorder and coping motives for drinking among canadian men and women*. In: Leo Sher, editors. *Comorbidity of Depression and Alcohol Use Disorders*. New York: Nova Science Publishers; 2009. p. 11-30.
- Racic M, Kusmuk S, Kozomara L, Debelnogi B. *The prevalence of mistreatment among the elderly with mental disorders in primary health care settings*. *J Adult Prot* 2006; 8 (4):20-24.
- Mukamal KJ, Kuller LH, Fitzpatrick AL, Longstreth WT Jr, Mittelman MA, Siscovick DS. *Prospective study of alcohol consumption and risk of dementia in older adults*. *JAMA* 2003;289(11):1405-13./29
- Query WT, Megran J. *Influence of Depression and Alcoholism on Learning, Recall, and Recognition*. *J Clin Psychol* 1984;40:1097.
- Liappas I. *Alcoholism and dementia*. *Ann Gen Psychiatry* 2003; 2 Suppl 1:39.
- Mustaca AE, Kamenetzky GV. *Alcoholismo y Ansiedad: Modelos Animales [Alcoholism and Anxiety: Animal Models]*. *Rev Int Psicol Ter Psicol* 2006; 6(3):343-364.
- Volk RJ, et al. *Alcohol use disorders, consumption patterns, and health-related quality of life of primary care patients*. *Alcohol Clin Exp Res* 1997; 21(5):899-905.

Примљен • Received: 31/08/2013
Прихваћен • Accepted: 14/01/2014