UDC 631.415.2; 634.11
COBISS.SR-ID: 220312076
Original research paper

Acta Agriculturae Serbica, Vol. XX, 40 (2015); 165-171

Effect of high Al content in the soil on vegetative rootstocks for apple

Gorica Paunović, Ljiljana Bosković-Rakočević, Radmila Ilić

Faculty of Agronomy, University of Kragujevac, Serbia

Abstract: Proper nutrition of fruit trees depends not only on the availability of major chemical elements in the soil, but also on their inter-relationships, their existing forms in the soil and the capability of the root system to absorb nutrients. The intensity of nutrient uptake is also highly dependent on soil solution pH. In recent years, large areas of land have been increasingly underused for fruit production. This can be partly attributed to soil acidification which also leads to increased concentrations of H, Al, Fe and Mn ions in its adsorptive complex. In order to neutralize soil acidity and Al toxicity, neutralizing agents should be used to help the soil become capable of fruit production. For this purpose, the following lime materials were used: CaO, bentonite and lifos. The commonly used vegetative rootstocks of different vigor for apple M9, M26 and MM106 were planted as test plants. The impact of lime materials was evaluated with respect to the leaf assimilating surface area, vegetative growth and tree diameter of the rootstocks used. The best soil amending effect was achieved by CaO, followed in declining order by lifos and bentonite.

Key words: acid soil, mobile Al, vegetative rootstocks for apple, lime materials

Introduction

Proper nutrition of fruit trees depends not only on the supply of major chemical elements in the soil, but also on their relationship, their existing forms in the soil and the capability of the root system to absorb nutrients. The uptake of nutrients from the soil is largely affected by soil pH value, which is of special importance given the low pH of a large part of our land, especially in highland areas, where fruit production predominates.

If these soils contain a high concentration of aluminum, usually at a pH below 5, fruit yields are low or fruit production is even impossible. In addition to aluminum toxicity, such soils have a reduced content of base cations in the soil adsorption complex, mineral (P, Ca, Mg) deficiencies and excess soluble Mn and Fe (Dugalić, 1997; Radanović and Predić, 1997; Narro et al., 2001; Bošković Rakočević and Bokan, 2002, Sumner, 2004).

Most fruit trees require a pH of 5.5 to 6.5 for their flourishing because in such soils their root system can develop normally, which is particularly important because they are perennial plantations. Adverse conditions for root development, due to the law of correlation, have a negative impact on the development of the above-ground system, which causes abnormal growth and development of fruit trees (Lučić et al., 1996). At the same time, nutrient uptake in such soils is limited, and very often there are symptoms of nutrient deficiencies, particularly of phosphorus, calcium and magnesium, on leaves and fruits (Ubavić et al., 2001).

In order to neutralize soil acidity and aluminum toxicity, proper liming agents should be used to neutralize the soil and, hence, recover the land for fruit production, and achieve regular and stable yields.

The aim of this study was to investigate possibilities for neutralization of soil acidity and aluminum toxicity using CaO, bentonite and lifos, and test their effects on breeding vegetative rootstocks of differing vigor for apple which are commonly used in the production of apple seedlings.

Materials and methods

The experiment was set up in vegetative containers filled with the soil obtained from the site Teocin, Mt. Suvobor.

The soil was acidic (pH / KCl 3.12), as determined by the content of 25 mg / 100 g of mobile aluminum.

To neutralize soil acidity, the following treatments were used:

- Calcium oxide 3.6 t / ha (T1)
- Bentonite 48 t / ha (T2)
- Lifos 25.5 t / ha (T3)
- Control (T4)

Vegetative apple-tree rootstocks of differing vigor M9, M26 and MM106 were planted as test plants.

Trials were set up in three variants, each with three replicates, with two bases at each iteration, plus controls. Care, protection, irrigation and fertilization experiments were uniform in all iterations and variants.

After the experiment, substitutional acidity in soil pH M KCl was performed (in suspension 1: 2.5, electrometric) and measurements were made of the assimilation leaf surface, vegetative growth and diameter of the aboveground part of the rootstock.

Statistical analysis of the results was performed by analysis of variance, and significant differences were found by LSD test.

Results and Discussion

Given the extremely high soil acidity of the soils (pH / KCl 3.12), after treatment with lime materials, the soil pH was checked, and the data showed that acidity decreased in all variants. The highest pH value was determined in treatment with CaO (pH / KCl 4.08), followed in declining order by lifos (pH / KCl 3.67) and bentonite (pH / KCl 3.59), which are still very low values, but changes in acidity were found by Bošković Rakočević et al. (2003).

The effect of treatment on the leaf assimilating surface area

Research results on the effect of different agents used to neutralize soil acidity on the size of the leaf assimilating surface area in apple rootstocks are given in Table 1.

Table 1. The effect of treatment on the leaf assimilating surface area

Treatments	Rootstock			Average
	M9	M26	MM106	
T ₁	25.91	21.39	25.68	24.33
T ₂	20.54	18.92	24.65	21.37
T ₃	25.25	19.55	26.36	23.72
T ₄	19.73	16.00	17.19	17.64
Average	22.85	18.96	23.47	

LSD_{0.05} = 16.25

LSD_{0.01} = 24.61

The analysis of the results shows that the highest leaf assimilating surface area was formed in treatment with CaO (T₁), followed by treatment with lifos

(T3), and the control (T4). As shown by the statistical analysis, no statistically significant differences were observed between the surface M9 and MM106, while in relation to the base M26 the differences were statistically significant.

The effect of treatment on vegetative growth

The results (Tab. 2) show that the greatest vegetative growth was obtained under CaO treatment i.e. an average 152.3 cm, which was a 64.1 cm increase relative to the control, and the increase was statistically significant. The weakest effect was achieved by the use of bentonite (average 93.8 cm), which was an increase of only 5.6 cm compared to the control, but without statistical significance.

Table 2. The effect of treatment on vegetative growth

Treatments	Rootstock			
	M9	M26	MM106	Average
T ₁	129.5	168.8	158.5	152.3
T ₂	103.8	99.2	78.4	93.8
T ₃	104.5	112.0	137.9	118.1
T ₄	94.3	77.0	93.4	88.2
Average	108.0	114.2	117.1	

LSD_{0.05} = 16.25

LSD_{0.01} = 24.61

The comparison of the tested rootstocks shows that although they differ in vigor, there are no statistically significant differences in vegetative growth. However, the interaction between the vegetative surface and lime material showed statistically significant differences in the application of CaO and lifos compared to all other treatments.

The effect of treatment on the diameter of the aboveground part of the rootstock

The diameter of the aboveground part of the rootstock was measured at a height of 15 cm above the surface. As with the previously analyzed parameters, better effects on the diameter of the aboveground part of the rootstock were obtained by the use of CaO (12.3 mm), and the same average value was determined after lifos application (Tab. 3). Statistical analysis revealed a highly significant difference between the acidity neutralizing agents, while the rootstock had no significant influence.

Table 3. The effect of treatment on the diameter of the aboveground part of the rootstock

Treatments	Rootstock			Average
	M9	M26	MM106	
T ₁	11.6	12.7	12.5	12.3
T ₂	10.6	12.3	12.2	11.7
T ₃	12.6	12.6	11.7	12.3
T ₄	11.4	9.4	10.1	10.3
Average	11.5	11.7	11.6	

LSD_{0.05} = 0.53LSD_{0.01} = 0.81

Conclusion

The research results suggest the following:

Upon completion of the experiment, the maximum pH value of the soil was in the treatment with CaO (4.08), followed by lifos (3.67) and bentonite (3:59) applications.

The highest assimilating surface area of the leaf was obtained under CaO treatment, lifos and bentonite had weaker effects, and the lowest surface area was found in the control.

Maximum vegetative growth was achieved after CaO treatment.

The diameter of the aboveground part of the rootstock was the same in treatments with CaO and lifos, and somewhat lower after bentonite application.

Among the lime materials used, the best effect on all tested parameters was achieved by CaO treatment.

All agents used to neutralize soil acidity showed a positive effect on the test parameters in vegetative rootstocks for apple. This indicates the need to increase the pH value of the soil to ensure the production of quality planting material.

References

- Bošković Rakočević Lj., Bokan N. (2002): Mogućnosti rešavanja problema kiselih zemljišta i toksičnosti aluminijuma. Eko-konferencija, Tematski zbornik, Novi Sad, 113-118.
- Bošković Rakočević Lj., Jakovljević M., Ubavić, M., Milivojević J. (2003): The Changes in Soil Acidity Depending on Amelioration Measures. *Journal of Agricultural Sciences*, 48(2): 149-158.
- Dugalić, G. (1997): Karakteristike Kraljevačkog pseudogleja i iznalaženje mogućnosti za povećanje njegove produktivne sposobnosti. Doktorska disertacija, Poljoprivredni fakultet, Zemun.
- Lučić, P., Đurić, G., Mičić, N. (1996): Voćarstvo. Nolit-Partenon, Beograd.
- Narro, L. A., Perez, C. J., Pandey, S., Crossa, J., Salazar, F., Arias, P. M. (2001): Implications of soil-acidity tolerant maize cultivars to increase production in developing countries. In: N. Ae et al. (ed.). *Plant nutrient acquisition: New perspectives*. NIAES series 4. Springer Verlag, Japan, 447-463.
- Radanović, D., Predić, T. (1997): Primeri toksičnosti Al i Mn na usevima pšenice i ječma na distričnom pseudogleju i degradiranom smeđem zemljištu. IX Kongres JDPZ "Uređenje, korišćenje i očuvanje zemljišta", Zbornik radova, Novi Sad, 235-243.
- Sumner, M. E. (2004): Food production on acid soils in the developing world: problems and solutions. *Proceedings of the 6th International Symposium on Plant-Soil Interactions at Low pH* (Editors: Matsumoto H. et al.), August 1-5, Sendai, Japan, 2-3.
- Ubavić, M., Kastori, R., Oljača, R., Marković, M. (2001): Ishrana voćaka. Naučno voćarsko društvo Republike Srpske i Poljoprivredni fakultet u Banjaluci, Banjaluka.

TOLERANTNOST VEGETATIVNIH PODLOGA ZA JABUKU NA POVEĆANU KISELOST ZEMLJIŠTA

Gorica Paunović, Ljiljana Bošković-Rakočević, Radmila Ilić

Agronomski fakultet, Univerzitet u Kragujevcu, Srbija

Rezime

Usvajanje hranljivih materija iz zemljišta od strane korenovog sistema voćaka ne zavisi samo od snabdevenosti zemljišta tim elementima, oblika u kome se oni nalaze, već i od reakcije sredine, tj. pH zemljišta. Poslednjih godina sve prisutniji problem u voćarskoj proizvodnji jeste pojava acidifikacije zemljišta, koja ima za posledicu povećanje sadržaja H, Al, Fe i Mn jona u adsorptivnom kompleksu zemljišta. Kisela zemljišta ograničavaju izbor voćnih vrsta i sorti koje se na njima mogu uspešno gajiti, pa je potrebno ispitivati tolerantnost i podloga i sorti na povećanu kiselost zemljišta. U cilju neutralizacije kiselosti i toksičnosti aluminijuma, potrebno je pronaći sredstva za njihovo neutralisanje i na taj način ponovo osposobiti zemljište za uspešnu voćarsku proizvodnju. U tu svrhu primenjeni su sledeći krečni materijali: CaO, bentonit i lifos. Kao test biljke zasadene su vegetativne podloge za jabuku različite bujnosti: M9, M26 i MM106, koje su u proizvodnoj praksi najčešće korišćene. Dejstvo krečnih materijala je ispitivano u odnosu na asimilacionu površinu lista, vegetativni porast i prečnik nadzemnog dela korišćenih podloga. Najbolji efekat u popravci zemljišta postignut je uz pomoć CaO, nešto slabiji korišćenjem lifosa, a najslabiji primenom bentonita.

Ključne reči: pH zemljišta, mobilni Al, vegetativne podloge za jabuku, krečni materijali.