Oscillation of second-order nonlinear difference equations with sublinear neutral term

MARTIN BOHNER, HASSAN A. EL-MORSHEDY,
SAID R. GRACE, ILGIN SAĞER

ABSTRACT. We establish some new criteria for the oscillation of second-order nonlinear difference equations with a sublinear neutral term. This is accomplished by reducing the involved nonlinear equation to a linear inequality.

1. INTRODUCTION

This paper deals with oscillatory behavior of all solutions of nonlinear second-order difference equations with a sublinear neutral term of the form

$$(1) \quad \Delta(a_n \Delta(x_n + p_n x_{n-k}^\alpha)) + q_n x_{n+1-m}^\beta = 0.$$

We assume that

- (H₁) $0 < \alpha < 1$ and $\beta > 0$ are ratios of positive odd integers,
- (H₂) $\{a_n\}$, $\{p_n\}$, $\{q_n\}$, $n \geq n_0$, are positive real sequences,

$$\lim_{n \rightarrow \infty} p_n = 0 \quad \text{and} \quad \sum_{s=n_0}^{\infty} \frac{1}{a_s} < \infty,$$

- (H₃) $k \in \mathbb{N}$ and $m \in \mathbb{N}_0$.

Let $\xi = \max\{k, m - 1\}$. By a solution of (1), we mean a real sequence $\{x_n\}$ defined for all $n \geq n_0 - \xi$ that satisfies (1) for $n \geq n_0$. A solution of (1) is said to be oscillatory if its terms are neither eventually positive nor eventually negative, and otherwise it is called nonoscillatory. Equation (1) is said to be oscillatory if all its solutions are oscillatory.

In recent years, there has been a great interest in establishing criteria for the oscillation and asymptotic behavior of solutions of various classes of second-order difference equations, see [1, 2, 4, 9–12, 15, 18, 20, 21, 24] and the references cited therein. However, it seems that there are no known results regarding the oscillation of second-order difference equations with positive

2010 *Mathematics Subject Classification.* Primary: 39A10, 39A21.

Key words and phrases. Neutral term, nonlinear difference equation, oscillation.

Full paper. Received 28 November 2018, accepted 28 December 2019, available online 17 January 2019.

sublinear neutral term. More exactly, the existing literature does not provide any criteria which ensure oscillation of all solutions of (1). In view of this motivation, our aim in this paper is to present sufficient conditions which ensure that all solutions of (1) are oscillatory. For related results concerning second-order differential equations with sublinear neutral term, we refer the reader to [3, 16, 17, 23]. Some related results concerning second-order dynamic equations on time scales can be found in [6–8, 13, 14, 19, 22].

2. MAIN RESULTS

For $n \geq n_0^*$ for some $n_0^* \geq n_0$, we let

$$A_n = \sum_{s=n}^{\infty} \frac{1}{a_s}.$$

For convenience, for some $0 < \nu \leq 1$ and $n \geq n_0^*$, we set

$$y_n = x_n + p_n x_{n-k}^{\alpha},$$

$$P_n = 1 - p_n \frac{A_{n-k}^{\alpha}}{A_n^{1+(1-\alpha)\nu}} \geq 0$$

and

$$Q_n = q_n A_{n+1}^{(1+\nu)(\beta-1)} P_{n+1-m}^{\beta}.$$

In the following, we establish a new oscillation result for (1) when $\beta \geq 1$.

Theorem 2.1. *Let $\beta \geq 1$. Assume (H_1) – (H_3) . If*

$$(2) \quad \limsup_{n \rightarrow \infty} \sum_{s=n_0^*}^n \left[Q_s A_{s+1} - \frac{1}{4a_s A_{s+1}} \right] = \infty,$$

then (1) is oscillatory.

Proof. Let x_n be a nonoscillatory solution of (1), say $x_n > 0$, $x_{n+1-m} > 0$, $x_{n-k} > 0$, and $y_n > 0$ for $n \geq n_1$ for some $n_1 \geq n_0^*$. It is easy to see that $y_n > 0$, $n \geq n_1$, and (1) becomes

$$(3) \quad \Delta(a_n \Delta y_n) + q_n x_{n+1-m}^{\beta} = 0.$$

Thus $\Delta(a_n \Delta y_n) \leq 0$ for $n \geq n_1$, which implies that y_n is bounded. Also, the decreasing nature of $a_n \Delta y_n$ implies that (I) $\Delta y_n > 0$ or (II) $\Delta y_n < 0$ for $n \geq n_1^* \geq n_1$. Therefore, y_n converges, and hence

$$\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} (x_n + p_n x_{n-k}^{\alpha}) = \lim_{n \rightarrow \infty} x_n,$$

since $\lim_{n \rightarrow \infty} p_n = 0$. Now, we consider Case (I). Since y_n is a positive increasing sequence, there exist $n_2 \geq n_1^*$ and $d > 0$ such that

$$(4) \quad x_n > d \quad \text{for } n \geq n_2.$$

Substituting (4) into (3), we get

$$(5) \quad \Delta(a_n \Delta y_n) + q_n d^\beta < 0 \quad \text{for } n \geq n_2.$$

Summing (5) from n_2 to $n-1$, we obtain

$$(6) \quad a_n \Delta y_n - a_{n_2} \Delta y_{n_2} + d^\beta \sum_{s=n_2}^{n-1} q_s < 0 \quad \text{for } n \geq n_2.$$

But (2) implies that

$$\sum_{n=n_2}^{\infty} q_n = \infty,$$

which together with (6) yields

$$\lim_{n \rightarrow \infty} a_n \Delta y_n = -\infty,$$

a contradiction due to the eventual positivity of $a_n \Delta y_n$.

Next, we consider Case (II). Define the sequence $\{v_n\}$ by

$$(7) \quad v_n = \frac{a_n \Delta y_n}{y_n} \quad \text{for } n \geq n_1.$$

Then $v_n < 0$ for $n \geq n_1$. Also, the decreasing nature of $a_n \Delta y_n$ implies that

$$(8) \quad \Delta y_s \leq \frac{a_n}{a_s} \Delta y_n \quad \text{for } s \geq n \geq n_1.$$

Summing (8) from n to $r-1 \geq n$, we obtain

$$y_r - y_n \leq a_n \Delta y_n \left(\sum_{s=n}^{r-1} \frac{1}{a_s} \right),$$

which, by letting $r \rightarrow \infty$, leads to

$$(9) \quad \frac{a_n \Delta y_n}{y_n} A_n \geq -1 \quad \text{for } n \geq n_1,$$

i.e.,

$$(10) \quad v_n A_n \geq -1 \quad \text{for } n \geq n_1.$$

On the other hand, we find from (9) that

$$\Delta \left(\frac{y_n}{A_n} \right) = \frac{A_n \Delta y_n - y_n \Delta A_n}{A_n A_{n+1}} = \frac{A_n \Delta y_n + \frac{y_n}{a_n}}{A_n A_{n+1}} \geq 0,$$

for $n \geq n_1$, and thus

$$(11) \quad \frac{y_n}{A_n} \geq \frac{y_{n-k}}{A_{n-k}} \quad \text{for } n \geq n_1 + k.$$

Now,

$$x_n = y_n - p_n x_{n-k}^\alpha \geq y_n - p_n y_{n-k}^\alpha \quad \text{for } n \geq n_1 + k,$$

and using (11), we obtain

$$(12) \quad x_n \geq y_n - p_n \frac{A_{n-k}^\alpha}{A_n^\alpha} y_n^\alpha = \left(1 - p_n \frac{A_{n-k}^\alpha}{A_n^\alpha} y_n^{\alpha-1}\right) y_n.$$

Since y_n/A_n is positive and increasing, we get

$$(13) \quad \frac{y_n}{A_n} \geq \frac{y_{n_1}}{A_{n_1}} =: \gamma > 0 \quad \text{for } n \geq n_1.$$

Since $\{A_n\}$ is positive and converging to zero, there exists $n_3 \geq n_1 + k$ such that

$$(14) \quad 0 < A_n^\nu \leq \gamma \quad \text{for all } n \geq n_3.$$

Hence, by (13) and (14),

$$(15) \quad y_n \geq A_n^{1+\nu} \quad \text{for } n \geq n_3.$$

Using (15) in (12), we get

$$(16) \quad x_n \geq \left(1 - p_n \frac{A_{n-k}^\alpha}{A_n^\alpha} A_n^{(1+\nu)(\alpha-1)}\right) y_n = P_n y_n \quad \text{for } n \geq n_3.$$

By (16), from (3), we have

$$(17) \quad \begin{aligned} \Delta(a_n \Delta y_n) &= -q_n x_{n+1-m}^\beta \\ &\leq -q_n P_{n+1-m}^\beta y_{n+1-m}^\beta \\ &\leq -q_n P_{n+1-m}^\beta y_{n+1}^\beta \quad \text{for } n \geq n_3, \end{aligned}$$

where we also used the decreasing nature of y_n in the last estimate. Now (17), in view of (15), leads to

$$(18) \quad \begin{aligned} \Delta(a_n \Delta y_n) &\leq -q_n A_{n+1}^{(1+\nu)(\beta-1)} P_{n+1-m}^\beta y_{n+1} \\ &= -Q_n y_{n+1} \quad \text{for } n \geq n_3. \end{aligned}$$

Taking the difference of both sides of (7) and using the decreasing nature of $a_n \Delta y_n$, we get

$$(19) \quad \begin{aligned} \Delta v_n &= \frac{y_n \Delta(a_n \Delta y_n) - a_n (\Delta y_n)^2}{y_n y_{n+1}} \\ &= \frac{\Delta(a_n \Delta y_n)}{y_{n+1}} - \frac{y_n}{a_n y_{n+1}} v_n^2 \\ &\leq \frac{\Delta(a_n \Delta y_n)}{y_{n+1}} - \frac{v_n^2}{a_n} \quad \text{for } n \geq n_3, \end{aligned}$$

where we have used again the decreasing nature of y_n . Combining (19) and (18), we have

$$(20) \quad \Delta v_n \leq -Q_n - \frac{v_n^2}{a_n} \quad \text{for } n \geq n_3.$$

Using (20), we get

$$\begin{aligned}
 \Delta(A_n v_n) &= v_n \Delta A_n + A_{n+1} \Delta v_n \\
 &= -\frac{v_n}{a_n} + A_{n+1} \Delta v_n \\
 &\leq -\frac{v_n}{a_n} - A_{n+1} Q_n - \frac{A_{n+1} v_n^2}{a_n} \\
 &\leq -A_{n+1} Q_n + \frac{1}{4a_n A_{n+1}},
 \end{aligned}$$

and summing this resulting inequality from n_3 to n and using (10) yields

$$\begin{aligned}
 \sum_{s=n_3}^n \left[Q_s A_{s+1} - \frac{1}{4a_s A_{s+1}} \right] &\leq A_{n_3} v_{n_3} - A_{n+1} v_{n+1} \\
 &\leq 1 + A_{n_3} v_{n_3} < \infty \quad \text{for } n \geq n_3,
 \end{aligned}$$

contradicting (2). This completes the proof. $\square$

When $\beta = 1$, we have the following immediate corollary from Theorem 2.1.

Corollary 2.1. *Let $\beta = 1$. Assume $(H_1)-(H_3)$. If*

$$(21) \quad \limsup_{n \rightarrow \infty} \sum_{s=n_0^*}^n \left[q_s P_{s+1-m} A_{s+1} - \frac{1}{4a_s A_{s+1}} \right] = \infty,$$

then (1) is oscillatory.

Next, we establish an oscillation result when $0 < \beta < 1$.

Theorem 2.2. *Let $0 < \beta < 1$. Assume $(H_1)-(H_3)$. If*

$$(22) \quad \limsup_{n \rightarrow \infty} \sum_{s=n_0^*}^n \left[L q_s P_{s+1-m}^\beta A_{s+1} - \frac{1}{4a_s A_{s+1}} \right] = \infty \quad \text{for some } L > 0,$$

then (1) is oscillatory.

Proof. Let x_n be a nonoscillatory solution of (1), say $x_n > 0$, $x_{n+1-m} > 0$, $x_{n-k} > 0$, and $y_n > 0$ for $n \geq n_1$ for some $n_1 \geq n_0^*$. Proceeding as in the proof of Theorem 2.1, we obtain the two cases (I) $\Delta y_n > 0$ or (II) $\Delta y_n < 0$ for $n \geq n_1$. Next, we consider only Case (II) as Case (I) can be treated similarly as in the proof of Theorem 2.1. Recall that y_n is positive and decreasing with $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} x_n$. Then we have either $\lim_{n \rightarrow \infty} y_n = d_1 > 0$ or $\lim_{n \rightarrow \infty} y_n = 0$. The first case implies that $\lim_{n \rightarrow \infty} x_n = d_1$. Thus, there exist $d_2 > 0$ and $n_1^* \in \mathbb{N}$ such that $x_n \geq d_2$ for all $n \geq n_1^*$. Hence we can obtain a contradiction similarly as in Case (I). The other case implies that for $K := L^{1/(\beta-1)} > 0$, there exists $n_2^* \in \mathbb{N}$ such that

$$(23) \quad 0 < y_n < K \quad \text{for all } n \geq n_2^*.$$

Now proceeding as in the proof of Theorem 2.1, we obtain (17), which with (23) yields

$$\begin{aligned}
 0 &\geq \Delta(a_n \Delta y_n) + q_n P_{n+1-m}^\beta y_{n+1}^\beta \\
 &= \Delta(a_n \Delta y_n) + \frac{q_n P_{n+1-m}^\beta y_{n+1}}{y_{n+1}^{1-\beta}} \\
 &\geq \Delta(a_n \Delta y_n) + \frac{q_n P_{n+1-m}^\beta y_{n+1}}{K^{1-\beta}} \\
 &= \Delta(a_n \Delta y_n) + L q_n P_{n+1-m}^\beta y_{n+1} \quad \text{for } n \geq n_3,
 \end{aligned}$$

with some $n_3 \geq n_2^*$. The remainder of the proof is similar to that of Theorem 2.1 and hence is omitted. $\square$

3. EXAMPLES AND REMARKS

First, we give two examples for the case $\beta > 1$.

Example 3.1. Consider the second-order equation

$$(24) \quad \Delta \left(n(n+1) \Delta \left(x_n + \frac{x_{n-k}^\alpha}{n^2} \right) \right) + (n+1)^6 x_{n+1-m}^{\frac{5}{3}} = 0, \quad n \in \mathbb{N}.$$

Here, $0 < \alpha < 1$ is a ratio of positive odd integers, $\beta = 5/3$, the delays are $k \in \mathbb{N}$ and $m \in \mathbb{N}_0$, and

$$a_n = n(n+1), \quad p_n = \frac{1}{n^2}, \quad \text{and} \quad q_n = (n+1)^6.$$

We let $\nu = 1$. It is easy to see that (H_2) holds. Also,

$$A_n = \frac{1}{n} \quad \text{and} \quad P_n = 1 - \frac{1}{n^\alpha (n-k)^\alpha}.$$

Moreover,

$$Q_n A_{n+1} - \frac{1}{4a_n A_{n+1}} = (n+1)^{\frac{11}{3}} \left[1 - \frac{1}{(n+1-m)^\alpha (n+1-m-k)^\alpha} \right]^{\frac{5}{3}} - \frac{1}{4n}.$$

Thus,

$$\lim_{n \rightarrow \infty} \left(Q_n A_{n+1} - \frac{1}{4a_n A_{n+1}} \right) = \infty.$$

Therefore, (2) of Theorem 2.1 is satisfied, and hence (24) is oscillatory.

Example 3.2. Consider the second-order equation

$$(25) \quad \Delta \left(n(n+1) \Delta \left(x_n + \frac{x_{n-k}^\alpha}{n^2} \right) \right) + (n+1)^2 x_{n+1-m}^{\frac{5}{3}} = 0, \quad n \in \mathbb{N}.$$

Here, all data are the same as in Example 3.1 except

$$q_n = (n+1)^2,$$

and therefore

$$Q_n A_{n+1} = \frac{1}{(n+1)^{\frac{1}{3}}} \left[1 - \frac{1}{(n+1-m)^\alpha (n+1-m-k)^\alpha} \right]^{\frac{5}{3}} \geq \frac{1}{n} \cdot \frac{1}{2},$$

for $n \geq N$ with some $N \in \mathbb{N}$, and thus

$$\begin{aligned} \sum_{n=N}^M \left(Q_n A_{n+1} - \frac{1}{4a_n A_{n+1}} \right) &= \sum_{n=N}^M \left(Q_n A_{n+1} - \frac{1}{4n} \right) \\ &\geq \sum_{n=N}^M \frac{1}{4n} \rightarrow \infty \quad \text{as } M \rightarrow \infty. \end{aligned}$$

Hence, (2) of Theorem 2.1 is satisfied, and thus (25) is oscillatory.

Next, we give an example in the case $\beta = 1$.

Example 3.3. Consider the second-order equation

$$(26) \quad \Delta \left(n(n+1) \Delta \left(x_n + \sqrt[3]{\frac{(n-k)x_{n-k}}{8n^4}} \right) \right) + \frac{n+1}{n} x_{n+1-m} = 0, \quad n \in \mathbb{N}.$$

Here, $\alpha = 1/3$, $\beta = 1$, the delays are $k \in \mathbb{N}$ and $m \in \mathbb{N}_0$, and

$$a_n = n(n+1), \quad p_n = \sqrt[3]{\frac{n-k}{8n^4}} \quad \text{and} \quad q_n = \frac{n+1}{n}.$$

We let $\nu = 1/2$. It is easy to see that (H_2) holds. Also,

$$A_n = \frac{1}{n} \quad \text{and} \quad P_n = \frac{1}{2}.$$

Moreover,

$$q_n P_{n+1-m} A_{n+1} - \frac{1}{4a_n A_{n+1}} = \frac{q_n}{2(n+1)} - \frac{1}{4n} = \frac{1}{4n}.$$

Therefore, (21) of Corollary 2.1 is satisfied, and hence (26) is oscillatory.

Finally, we present an example in the case $0 < \beta < 1$.

Example 3.4. Consider the second-order equation

$$(27) \quad \Delta \frac{\Delta \left(x_n + 4^{(\alpha-1)(n-1)-(k\alpha+1)/2} x_{n-k}^\alpha \right)}{2^n} + n 8^n x_{n+1-m}^\beta = 0, \quad n \in \mathbb{N}.$$

Here, $0 < \alpha < 1$ and $0 < \beta < 1$ are ratios of positive odd integers, the delays are $k \in \mathbb{N}$ and $m \in \mathbb{N}_0$, and

$$a_n = \frac{1}{2^n}, \quad p_n = 4^{(\alpha-1)(n-1)-(k\alpha+1)/2} \quad \text{and} \quad q_n = n 8^n.$$

We let $\nu = 1$. It is easy to see that (H_2) holds. Also,

$$A_n = \frac{1}{2^{n-1}} \quad \text{and} \quad P_n = \frac{1}{2}.$$

Moreover,

$$Lq_n P_{n+1-m}^\beta A_{n+1} - \frac{1}{4a_n A_{n+1}} = Ln2^{2n-\beta} - 2^{2n-2},$$

which tends to infinity for any constant $L > 0$. Therefore, (22) of Theorem 2.2 is satisfied, and hence (27) is oscillatory.

Remark 3.1. The results of this paper are presented in a form that makes it easy to study extensions to higher-order equations. It would also be of interest to use the approach here to study (1) with $\alpha > 1$, i.e., (1) with superlinear neutral term.

Remark 3.2. Another possibility for extension of the presented results would be to consider the time-scales [5, 8] analogue of (1).

REFERENCES

- [1] R. P. Agarwal, *Difference equations and inequalities: Theory, methods, and applications*, Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, Inc., New York, 2nd Ed., 2000.
- [2] R. P. Agarwal, M. Bohner, S. R. Grace, D. O'Regan, *Discrete oscillation theory*, Hindawi Publishing Corporation, New York, 2005.
- [3] R. P. Agarwal, M. Bohner, T. Li, C. Zhang, *Oscillation of second-order differential equations with a sublinear neutral term*, Carpathian J. Math., 30 (1) (2014), 1–6.
- [4] R. P. Agarwal, S. R. Grace, D. O'Regan, *Oscillation theory for difference and functional differential equations*, Kluwer Academic Publishers, Dordrecht, 2000.
- [5] M. Bohner, S. G. Georgiev, *Multivariable dynamic calculus on time scales*, Springer, Cham, 2016.
- [6] M. Bohner, S. R. Grace, I. Jadlovská, *Oscillation criteria for second-order neutral delay differential equations*, Electron. J. Qual. Theory Differ. Equ., 2017, No. 60, 1–12.
- [7] M. Bohner, T. Li, *Oscillation of second-order p -Laplace dynamic equations with a nonpositive neutral coefficient*, Appl. Math. Lett., 37 (2014), 72–76.
- [8] M. Bohner, A. Peterson, *Dynamic equations on time scales: An introduction with applications*, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [9] C. Dharuman, J. R. Graef, E. Thandapani, K. S. Vidhyaa, *Oscillation of second order difference equation with a sub-linear neutral term*, J. Math. Appl., 40 (2017), 59–67.
- [10] H. A. El-Morshedy, *Oscillation and nonoscillation criteria for half-linear second order difference equations*, Dynam. Systems Appl., 15 (3-4) (2006), 429–450.

-
- [11] H. A. El-Morshedy, *New oscillation criteria for second order linear difference equations with positive and negative coefficients*, Comput. Math. Appl., 58 (10) (2009), 1988–1997.
 - [12] H. A. El-Morshedy, S. R. Grace, *Comparison theorems for second order nonlinear difference equations*, J. Math. Anal. Appl., 306 (1) (2005), 106–121.
 - [13] S. R. Grace, R. P. Agarwal, M. Bohner, D. O'Regan, *Oscillation of second-order strongly superlinear and strongly sublinear dynamic equations*, Commun. Nonlinear Sci. Numer. Simul., 14 (8) (2009), 3463–3471.
 - [14] S. R. Grace, M. Bohner, R. P. Agarwal, *On the oscillation of second-order half-linear dynamic equations*, J. Difference Equ. Appl., 15 (5) (2009), 451–460.
 - [15] S. R. Grace, H. A. El-Morshedy, *Oscillation criteria of comparison type for second order difference equations*, J. Appl. Anal., 6 (1) (2000), 87–103.
 - [16] S. R. Grace, J. R. Graef, *Oscillatory behavior of second order nonlinear differential equations with a sublinear neutral term*, Math. Model. Anal., 23 (2) (2018), 217–226.
 - [17] J. R. Graef, S. R. Grace, E. Tunç, *Oscillatory behavior of even-order nonlinear differential equations with a sublinear neutral term*, Opuscula Math., 39 (1) (2019), 39–47.
 - [18] W.-T. Li, S. H. Saker, *Oscillation of second-order sublinear neutral delay difference equations*, Appl. Math. Comput., 146 (2-3) (2003), 543–551.
 - [19] S. H. Saker, *Oscillation of superlinear and sublinear neutral delay dynamic equations*, Commun. Appl. Anal., 12 (2) (2008), 173–187.
 - [20] S. Selvarangam, M. Madhan, E. Thandapani, *Oscillation theorems for second order nonlinear neutral type difference equations with positive and negative coefficients*, Rom. J. Math. Comput. Sci., 7 (1) (2017), 1–10.
 - [21] S. Selvarangam, E. Thandapani, S. Pinelas, *Oscillation theorems for second order nonlinear neutral difference equations*, J. Inequal. Appl., 2014, 2014:417.
 - [22] A. K. Sethi, *Oscillation of second order sublinear neutral delay dynamic equations via Riccati transformation*, J. Appl. Math. Inform., 36 (3-4) (2018), 213–229.
 - [23] A. K. Tripathy, A. K. Sethi, *Oscillation of sublinear second order neutral differential equations via Riccati transformation*, In: Differential and difference equations with applications, Springer Proc. Math. Stat., 230 (2018), 543–557.
 - [24] M. K. Yildiz, H. Ögünmez, *Oscillation results of higher order nonlinear neutral delay difference equations with a nonlinear neutral term*, Hacet. J. Math. Stat., 43 (5) (2014), 809–814.

MARTIN BOHNER

DEPARTMENT OF MATHEMATICS AND STATISTICS
MISSOURI S&T
ROLLA, MO 65409-0020
USA

E-mail address: bohner@mst.edu

HASSAN A. EL-MORSHEDY

DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
DAMIETTA UNIVERSITY
NEW DAMIETTA 34517
EGYPT

E-mail address: elmorshedy@yahoo.com

SAID R. GRACE

DEPARTMENT OF ENGINEERING MATHEMATICS
FACULTY OF ENGINEERING
CAIRO UNIVERSITY
ORMAN, GIZA 12000
EGYPT

E-mail address: saidgrace@yahoo.com

ILGIN SAĞER

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
UNIVERSITY OF MISSOURI-ST. LOUIS
ST. LOUIS, MO 63121-4400
USA

E-mail address: sageri@umsl.edu