

Common fixed points under strict conditions

HAKIMA BOUHADJERA, SAID BELOUL, ACHREF EDDINE TABET

ABSTRACT. In this contribution, three new concepts called reciprocally continuous, strictly subweakly compatible and strictly subreciprocally continuous single and multivalued mappings are given for obtention some common fixed point theorems in a metric space. Our results improve and complement the results of Aliouche and Popa, Azam and Beg, Deshpande and Pathak, Kaneko and Sessa, Popa and others.

1. INTRODUCTION AND PRELIMINARIES

Let (\mathcal{X}, d) be a metric space. We denote by $CL(\mathcal{X})$ (resp., $CB(\mathcal{X})$) the nonempty closed (resp., closed and bounded) subsets of \mathcal{X} and H the Hausdorff metric on $CL(\mathcal{X})$ (resp., $CB(\mathcal{X})$)

$$H(A, B) = \max\{\sup_{x \in A} d(x, B), \sup_{x \in B} d(x, A)\},$$

where $A, B \in CL(\mathcal{X})$ or $CB(\mathcal{X})$, and

$$d(x, A) = \inf_{y \in A} \{d(x, y)\}.$$

Now, let f and g be two self-mappings of a metric space (\mathcal{X}, d) . In 1982, Sessa [11] gave the weaker concept of the commutativity, namely the weakly commuting notion. f and g are weakly commuting if

$$d(fgx, gfx) \leq d(gx, fx),$$

for all $x \in \mathcal{X}$.

In 1986, Jungck [5] gave a generalization of the weak commutativity by giving the notion of compatible mappings. He defined f and g to be compatible if

$$\lim_{n \rightarrow \infty} d(fgx_n, gfx_n) = 0,$$

whenever $\{x_n\}$ is a sequence in \mathcal{X} such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = t$ for some $t \in \mathcal{X}$.

2010 *Mathematics Subject Classification*. Primary: 47H10; Secondary: 54H25.

Key words and phrases. Strictly subweakly compatible mappings, reciprocally continuous mappings, strictly subreciprocally continuous mappings.

Full paper. Received 14 May 2020, accepted 1 July 2020, available online 10 August 2020.

Weakly commuting mappings are compatible. However, compatible mappings need not be weakly commuting (see example 2.2 of [13]).

In 1996, Jungck [6] generalized the above notion by introducing the concept of weakly compatible mappings. He defined f and g to be weakly compatible if they commute at their coincidence points; i.e., if $fu = gu$ for some $u \in \mathcal{X}$, then $fgu = gfu$.

If f and g are compatible then they are obviously weakly compatible but as shown in example 2.52 of [1] the converse is not true.

In their paper [8], Kaneko and Sessa extended the definition of compatibility to include multivalued mappings in the following way: mappings $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ are compatible if $fFx \in CB(\mathcal{X})$ for all $x \in \mathcal{X}$ and

$$\lim_{n \rightarrow \infty} H(fFx_n, Ffx_n) = 0,$$

whenever $\{x_n\}$ is a sequence in \mathcal{X} such that $\lim_{n \rightarrow \infty} Fx_n = M \in CB(\mathcal{X})$ and $\lim_{n \rightarrow \infty} fx_n = t \in M$.

To generalize the above notion, Jungck and Rhoades [7] gave the concept of weakly compatible mappings. $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ are said to be weakly compatible if they commute at their coincidence points; i.e., if $fFx = Ffx$ whenever $fx \in Fx$.

Recall that a point $t \in \mathcal{X}$ is called a **strict coincidence point** (resp. **strict common fixed point**) of mappings $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ if $Ft = \{ft\}$ (resp. $Ft = \{ft\} = \{t\}$).

2. MAIN RESULTS

Our first objective in this contribution is to generalize the above definition by introducing the concept of strictly subweakly compatible single and multivalued mappings.

Definition 1. Mappings $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ are **strictly subweakly compatible (shortly sswc)** if and only if $fFx \in CB(\mathcal{X})$ and there exists a sequence $\{x_n\}$ in \mathcal{X} such that $\lim_{n \rightarrow \infty} fx_n = t$, $\lim_{n \rightarrow \infty} Fx_n = \{t\}$ for some $t \in \mathcal{X}$ and $\lim_{n \rightarrow \infty} (fFx_n, Ffx_n) = 0$.

The example below shows that there exist sswc mappings which are not weakly compatible.

Example 1. Let $\mathcal{X} = [0, \infty)$ and $d(x, y) = |x - y|$. Define $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ by

$$fx = x^2 \text{ and } Fx = \begin{cases} [4, x + 2], & \text{if } x \in [2, 4] \cup (9, \infty), \\ \{x + 12\}, & \text{if } x \in [0, 2) \cup (4, 9]. \end{cases}$$

We have $fFx \in CB(\mathcal{X})$. Consider the sequence $\{x_n\}$ in \mathcal{X} defined by $x_n = 2 + \frac{1}{n}$ for $n = 1, 2, \dots$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} fx_n &= \lim_{n \rightarrow \infty} x_n^2 = 4 = t, \\ \lim_{n \rightarrow \infty} Fx_n &= \lim_{n \rightarrow \infty} [4, x_n + 2] = \{4\} = \{t\}, \\ \lim_{n \rightarrow \infty} H(fFx_n, Ffx_n) &= \lim_{n \rightarrow \infty} H([16, (x_n + 2)^2], \{x_n^2 + 12\}) = 0, \end{aligned}$$

therefore f and F are sswc.

On the other hand, we have $fx \in Fx$ if and only if $x \in [4, x + 2]$, but $Ff(x) \neq fF(x)$, therefore f and F are not weakly compatible.

In 1999, Pant [9] introduced the concept of reciprocally continuous single-valued mappings as a generalization of continuous mappings: f and g are reciprocally continuous if and only if $\lim_{n \rightarrow \infty} fgx_n = ft$ and $\lim_{n \rightarrow \infty} gfx_n = gt$ whenever $\{x_n\} \subset \mathcal{X}$ is such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = t \in \mathcal{X}$.

In 2002, Singh and Mishra [12] introduced the concept of reciprocal continuity for single and multivalued mappings as follows.

Definition 2. The mappings $F : \mathcal{X} \rightarrow CL(\mathcal{X})$ and $f : \mathcal{X} \rightarrow \mathcal{X}$ are reciprocally continuous on \mathcal{X} (resp., at $t \in \mathcal{X}$) if and only if $fFx \in CL(\mathcal{X})$ for each $x \in \mathcal{X}$ (resp., $fFt \in CL(\mathcal{X})$) and $\lim_{n \rightarrow \infty} fFx_n = fM$, $\lim_{n \rightarrow \infty} Ffx_n = Ft$ whenever $\{x_n\}$ is a sequence in \mathcal{X} such that $\lim_{n \rightarrow \infty} Fx_n = M \in CL(\mathcal{X})$, $\lim_{n \rightarrow \infty} fx_n = t \in M$.

Motivated by Pant, Singh and Mishra, we give the following notion of reciprocally continuous single and multivalued mappings which is different from the above definition and represents our second objective.

Definition 3. Mappings $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ are **reciprocally continuous** if and only if $\lim_{n \rightarrow \infty} fFx_n = \{ft\}$ and $\lim_{n \rightarrow \infty} Ffx_n = Ft$ whenever $\{x_n\} \subset \mathcal{X}$ is a sequence such that $\lim_{n \rightarrow \infty} fx_n = t$, $\lim_{n \rightarrow \infty} Fx_n = \{t\}$ for some $t \in \mathcal{X}$.

Our third objective here is to extend the concept of reciprocally continuous mappings of Pant and the above one to the setting of single and multivalued mappings.

Definition 4. Mappings $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ are **strictly subreciprocally continuous (shortly ssrc)** if and only if there exists a sequence $\{x_n\}$ in \mathcal{X} such that $\lim_{n \rightarrow \infty} fx_n = t$, $\lim_{n \rightarrow \infty} Fx_n = \{t\}$ for some $t \in \mathcal{X}$, $\lim_{n \rightarrow \infty} fFx_n = \{ft\}$ and $\lim_{n \rightarrow \infty} Ffx_n = Ft$.

The next example shows that there exist ssrc mappings which are not continuous.

Example 2. Let $\mathcal{X} = \mathbb{R}$. Define $f : \mathcal{X} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ by

$$fx = \begin{cases} x - 1, & \text{if } x < 0, \\ x, & \text{if } x \geq 0, \end{cases} \quad Fx = \begin{cases} [x - 1, -1], & \text{if } x < 0, \\ [0, x], & \text{if } x \geq 0. \end{cases}$$

It is clear to see that f and F are discontinuous at $x = 0$.

Consider the sequence $\{x_n\}$ in \mathcal{X} defined by $x_n = \frac{1}{n}$ for $n = 1, 2, \dots$. We have

$$\begin{aligned} \lim_{n \rightarrow \infty} fx_n &= \lim_{n \rightarrow \infty} x_n = 0 = t, \\ \lim_{n \rightarrow \infty} Fx_n &= \lim_{n \rightarrow \infty} [0, x_n] = \{0\} = \{t\}, \\ \lim_{n \rightarrow \infty} fFx_n &= \lim_{n \rightarrow \infty} [0, x_n] = \{0\} = \{f(0)\} = \{f(t)\}, \\ \lim_{n \rightarrow \infty} Ffx_n &= \lim_{n \rightarrow \infty} [0, x_n] = \{0\} = F(0) = F(t), \end{aligned}$$

therefore f and F are ssrc.

Now, we are ready to present and prove our main result.

Theorem 1. Let (\mathcal{X}, d) be a metric space. Let $f, g : \mathcal{X} \rightarrow \mathcal{X}$ and $F, G : \mathcal{X} \rightarrow CB(\mathcal{X})$ be single and multivalued mappings respectively such that f and F as well as g and G are reciprocally continuous and sswc or ssrc and compatible. Let $\varphi : \mathbb{R}_+^6 \rightarrow \mathbb{R}$ be a lower semi continuous function satisfying:

(φ_1): φ is nonincreasing in variables t_5 and t_6 ,

(φ_2): $\varphi(u, u, 0, 0, u, u) > 0$ for all $u > 0$ and the inequality

$$(1) \quad \varphi\left(H(Fx, Gy), d(fx, gy), d(fx, Fx), d(gy, Gy), d(fx, Gy), d(gy, Fx)\right) \leq 0,$$

for all x and y in \mathcal{X} , then, f, g, F and G have a strict common fixed point in \mathcal{X} .

Proof. Since f and F as well as g and G are reciprocally continuous and sswc or ssrc and compatible then, there exist two sequences $\{x_n\}$ and $\{y_n\}$ in \mathcal{X} such that $\lim_{n \rightarrow \infty} fx_n = t$, $\lim_{n \rightarrow \infty} Fx_n = \{t\}$ for some $t \in \mathcal{X}$, $\lim_{n \rightarrow \infty} fFx_n = \{ft\}$ and $\lim_{n \rightarrow \infty} Ffx_n = Ft$; $\lim_{n \rightarrow \infty} gy_n = z$, $\lim_{n \rightarrow \infty} Gy_n = \{z\}$ for some $z \in \mathcal{X}$, $\lim_{n \rightarrow \infty} gGy_n = \{gz\}$ and $\lim_{n \rightarrow \infty} Ggy_n = Gz$.

First we prove that $ft = gz$. In fact, by (1) we have

$$\begin{aligned} \varphi\left(H(Ft, Gz), d(ft, gz), d(ft, Ft), d(gz, Gz), d(ft, Gz), d(gz, Ft)\right) \leq 0. \end{aligned}$$

Since $Ft = \{ft\}$, $Gz = \{gz\}$ and φ is nonincreasing in t_5 and t_6 , we get

$$\varphi\left(d(ft, gz), d(ft, gz), 0, 0, d(ft, gz), d(gz, ft)\right) \leq 0,$$

which is a contradiction with φ_2 . Then $ft = gz$.

Now, we claim that $t = ft$, by (1) we have

$$\varphi\left(H(Fx_n, Gz), d(fx_n, gz), d(fx_n, Fx_n),\right. \\ \left. d(gz, Gz), d(fx_n, Gz), d(gz, Fx_n)\right) \leq 0.$$

Letting n tends to infinity and taking in account that φ is lower semi continuous, we get

$$\varphi(d(t, ft), d(t, ft), 0, 0, d(t, ft), d(ft, t)) \leq 0,$$

which contradicts φ_2 . Then $ft = t$.

Next, we prove that $z = t$. Indeed, by (1) we have

$$\varphi\left(H(Fx_n, Gy_n), d(fx_n, gy_n), d(fx_n, Fx_n),\right. \\ \left. d(gy_n, Gy_n), d(fx_n, Gy_n), d(gy_n, Fx_n)\right) \leq 0.$$

When n tends to infinity, we get

$$\varphi\left(d(t, z), d(t, z), 0, 0, d(t, z), d(z, t)\right) \leq 0,$$

which contradicts (φ_2) , therefore $z = t$. Consequently, t is a strict common fixed point of f, g, F and G . \square

Corollary 1. *Let (\mathcal{X}, d) be a metric space. Let $f, g : \mathcal{X} \rightarrow \mathcal{X}$ and $F, G : \mathcal{X} \rightarrow CB(\mathcal{X})$ be single and multivalued mappings respectively such that f and F as well as g and G are reciprocally continuous and sswc or ssrc and compatible. Assume that*

$$(2) \quad H(Fx, Gy) \leq m \max\left\{d(fx, gy), d(fx, Fx), d(gy, Gy),\right. \\ \left. \frac{1}{2}((d(fx, Gy) + d(gy, Fx)))\right\},$$

where $m \in (0, 1)$, or

$$(3) \quad H^2(Fx, Gy) \leq m^2 \max\left\{d^2(fx, gy), d(fx, Fx)d(gy, Gy), d(fx, Gy)d(gy, Fx),\right. \\ \left. d(fx, Gy)d(fx, Fx), d(gy, Fx)d(gy, Gy)\right\},$$

where $m^2 \in (0, 1)$, or

$$(4) \quad H^2(Fx, Gy) + \frac{H(Fx, Gy)}{1 + d(fx, Gy)d(gy, Fx)} - \\ - [ad^2(fx, gy) + bd^2(fx, Fx) + cd^2(gy, Gy)] \leq 0,$$

where $a, b, c > 0$ and $a + b + c < 1$.

Then f, g, F and G have a strict common fixed point.

- Remark 1.** (1) Our main result improves the main result of Popa [10].
 (2) By the above corollary and (2) for $f = g$ and $F = G$, we obtain an extension of the main result of Kaneko and Sessa [8].
 (3) The main result of Azam and Beg [3] follows from the above corollary and (2), because

$$d(fx, gy) \leq \max \left\{ d(fx, gy), d(fx, Fx), d(gy, Gy), \frac{1}{2}[d(fx, Gy) + d(gy, Fx)] \right\}.$$

- (4) Also, our main result improves the main results of Aliouche and Popa [2], Deshpande and Pathak [4] because, in our work, we have not continuity, neither completeness nor inclusion, and we did not impose a lot of conditions on the four mappings.

Example 3. Let $\mathcal{X} = [0, 2]$ endowed with the Euclidean metric, define mappings f, g, F and G as follows:

$$fx = \begin{cases} \frac{1}{2}(x+1), & 0 \leq x \leq 1, \\ 0, & 1 < x \leq 2, \end{cases} \quad gx = \begin{cases} x + \frac{1}{2}, & 0 \leq x < 1, \\ 1, & x = 1, \\ 2, & 1 < x \leq 2, \end{cases}$$

$$Fx = \begin{cases} [1, 2-x], & 0 \leq x \leq 1, \\ \{\frac{3}{2}\}, & 1 < x \leq 2, \end{cases} \quad Gx = \begin{cases} \{1\}, & 0 \leq x \leq 1 \\ [0, \frac{1}{2}], & 1 < x \leq 2. \end{cases}$$

We consider a sequence $\{x_n\}$ defined for each $n \geq 1$ by $x_n = 1 - \frac{1}{n}$, clearly $\lim_{n \rightarrow \infty} Fx_n = \{1\}$ and $\lim_{n \rightarrow \infty} fx_n = 1$, also we have

$$\lim_{n \rightarrow \infty} fFx_n = \{1\} = f\{1\},$$

$$\lim_{n \rightarrow \infty} Ffx_n = F1 = \{1\}.$$

Also, $\lim_{n \rightarrow \infty} H(fFx_n, Ffx_n) = 0$. Hence f and F are ssrc and compatible.

For g and G , consider a sequence $\{y_n\}$ defined by $y_n = \frac{1}{2}(1 - e^{-n})$, for all $n \geq 1$. It is clear that

$$\lim_{n \rightarrow \infty} gy_n = 1, \quad \lim_{n \rightarrow \infty} Gy_n = \{1\},$$

$$\lim_{n \rightarrow \infty} gGy_n = g\{1\} = \{1\}, \quad \lim_{n \rightarrow \infty} Ggy_n = G1 = \{1\},$$

$$\lim_{n \rightarrow \infty} H(gGy_n, Ggy_n) = 0,$$

i.e., g and G are ssrc and compatible.

By taking $m = \frac{4}{5}$ in corollary 1, we show that inequality (2) is satisfied. We have the following cases:

- (1) For $x, y \in [0, 1]$, we have $H(Fx, Gy) = 0$, obviously inequality (2) is satisfied.
 (2) For $x \in [0, 1]$ and $y \in (1, 2]$, we have

$$H(Fx, Gy) = 1 \leq \frac{6}{5} = \frac{4}{5}d(gy, Gy).$$

(3) For $x \in (1, 2]$ and $y \in [0, 1]$, we have

$$H(Fx, Gy) = \frac{1}{2} \leq \frac{6}{5} = \frac{4}{5}d(fx, Fx).$$

(4) For $x, y \in (1, 2]$, we have

$$H(Fx, Gy) = 1 \leq \frac{8}{5} = \frac{4}{5}d(fx, gy),$$

then all hypotheses of corollary 1 are satisfied and the point 1 is a strict common fixed point for f, g, F and G .

Corollary 2. *Let $f, g : \mathcal{X} \rightarrow \mathcal{X}$ be single valued mappings and let $F, G : \mathcal{X} \rightarrow CB(\mathcal{X})$ be multivalued mappings on a metric space (\mathcal{X}, d) such that the pairs f and F as well as g and G are ssrc and compatible for all $x, y \in \mathcal{X}$. Then the pair of mappings (f, F) and (g, G) has a strict coincidence point. Moreover f, g, F and G have a strict common fixed point in \mathcal{X} provided that mappings satisfy*

$$H(Fx, Gy) \leq \phi\left(d(fx, gy), d(fx, Fx), d(gy, Gy), d(fx, Gy), d(gy, Fx)\right),$$

where $\phi : \mathbb{R}_+^5 \rightarrow \mathbb{R}_+$ is an upper semi continuous function such that $\phi(0) = 0$ and $\phi(t, 0, 0, t, t) < t$ for each $t > 0$.

Proof. The proof follows immediately on taking

$$\varphi(t_1, t_2, t_3, t_4, t_5, t_6) = t_1 - \phi(t_2, t_3, t_4, t_5, t_6)$$

in Theorem 1, where $\phi : \mathbb{R}_+^5 \rightarrow \mathbb{R}_+$ is an upper semi continuous function such that $\phi(0) = 0$ and $\phi(t, 0, 0, t, t) < t$ for each $t > 0$. \square

3. ACKNOWLEDGMENT

The authors are highly thankful to the anonymous referee for his/her appreciation, valuable comments, and suggestions.

REFERENCES

- [1] R. P. Agarwal, R. K. Bisht, N. Shahzad, *A comparison of various noncommuting conditions in metric fixed point theory and their applications*, Fixed Point Theory and Applications, 38 (2014), 1–33.
- [2] A. Aliouche, V. Popa, *Coincidence and common fixed point theorems for hybrid mappings*, Mathematica Moravica, 12 (1) (2008), 1–13.
- [3] A. Azam, I. Beg, *Coincidence points of compatible multivalued mappings*, Demonstratio Mathematica, 29 (1) (1996), 17–22.
- [4] R. Deshpande, R. Pathak, *Common fixed point theorems for hybrid pairs of mappings using implicit relations*, Mathematica Moravica, 18 (1) (2014), 9–20.
- [5] G. Jungck, *Compatible mappings and common fixed points*, International Journal of Mathematics and Mathematical Sciences, 9 (4) (1986), 771–779.

- [6] G. Jungck, *Common fixed points for noncontinuous nonself maps on nonmetric spaces*, Far East Journal of Mathematical Sciences, 4 (2) (1996), 199–215.
- [7] G. Jungck, B. E. Rhoades, *Fixed points for set valued functions without continuity*, Indian Journal of Pure and Applied Mathematics, 29 (3) (1998), 227–238.
- [8] H. Kaneko, S. Sessa, *Fixed point theorems for compatible multi-valued and single-valued mappings*, International Journal of Mathematics and Mathematical Sciences, 12 (2) (1989), 257–262.
- [9] R. P. Pant, *A common fixed point theorem under a new condition*, Indian Journal of Pure and Applied Mathematics, 30 (2) (1999), 147–152.
- [10] V. Popa, *Some fixed point theorems for compatible mappings satisfying an implicit relation*, Demonstratio Mathematica, 32 (1) (1999), 157–163.
- [11] S. Sessa, *On a weak commutativity condition of mappings in fixed point considerations*, Publications de l'Institut Mathématique (N.S.), 32 (46) (1982), 149–153.
- [12] S. L. Singh, S. N. Mishra, *Coincidences and fixed points of reciprocally continuous and compatible hybrid maps*, International Journal of Mathematics and Mathematical Sciences, 30 (10) (2002), 627–635.
- [13] S. L. Singh, A. Tomar, *Weaker forms of commuting maps and existence of fixed points*, Journal of the Korean Society of Mathematical Education, Series B: Pure and Applied Mathematics, 10 (3) (2003), 145–161.

HAKIMA BOUHADJERA

LABORATORY OF APPLIED MATHEMATICS
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES
BADJI MOKHTAR-ANNABA UNIVERSITY
P.O. Box 12, 23000 ANNABA
ALGERIA
E-mail address: b_hakima2000@yahoo.fr

SAID BELOUL

EXACT SCIENCE FACULTY
EL OUED UNIVERSITY
P.O. Box 789, 39000 EL OUED
ALGERIA
E-mail address: beloulsaid@gmail.com

ACHREF EDDINE TABET

LABORATORY OF APPLIED MATHEMATICS
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES
BADJI MOKHTAR-ANNABA UNIVERSITY
P.O. Box 12, 23000 ANNABA
ALGERIA
E-mail address: achrefcte@gmail.com