On relationships between q -products identities, R_α, R_β and R_m functions related to Jacobi's triple-product identity

M. P. CHAUDHARY, SANGEETA CHAUDHARY

ABSTRACT. The authors establish a set of two new relationships involving q -product identities, R_α, R_β , and R_m ($m = 1, 2, 3, \dots$) functions; and answer a open question of Srivastava *et al.* [18]. The present work is motivated and based upon recent findings of Chaudhary *et al.* [8].

1. INTRODUCTION

The q -Pochhammer symbol or q -shifted factorial $(a; q)_n$ is defined (for $|q| < 1$) (see, for example, [3], [5, Chapter 3, Section 3.2.1], [19, Chapter 6] and [20, pp. 346 *et seq.*]), as

$$(a; q)_n := \begin{cases} 1, & (n = 0), \\ \prod_{k=0}^{n-1} (1 - aq^k), & (n \in \mathbb{N}), \end{cases} \quad (1)$$

where $a, q \in \mathbb{C}$ and it is assumed *tacitly* that $a \neq q^{-m}$ ($m \in \mathbb{N}_0$). We also write

$$(a; q)_\infty := \prod_{k=0}^{\infty} (1 - aq^k) = \prod_{k=1}^{\infty} (1 - aq^{k-1}), \quad (a, q \in \mathbb{C}; |q| < 1). \quad (2)$$

When $a \neq 0$ and $|q| \geq 1$, the infinite product in the equation (2) diverges. So, whenever $(a; q)_\infty$ is involved in a given formula, the constraint $|q| < 1$ will be *tacitly* assumed to be satisfied.

The following notations are also frequently used in our investigation:

$$(a_1, a_2, \dots, a_m; q)_n := (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n \quad (3)$$

2010 *Mathematics Subject Classification.* Primary: 11F27, 11P83; Secondary: 05A17, 05A30.

Key words and phrases. Theta-function identities; R_α, R_β , and R_m functions; Jacobi's triple-product identity; q -Product identities; Rogers-Ramanujan continued fraction; Rogers-Ramanujan identities.

Full paper. Received 2 June 2020, revised 11 August 2020, accepted 1 September 2020, available online 16 September 2020.

and

$$(a_1, a_2, \dots, a_m; q)_\infty := (a_1; q)_\infty (a_2; q)_\infty \cdots (a_m; q)_\infty. \quad (4)$$

Ramanujan (see [12] and [13]) defined the general theta function $f(a, b)$ as follows (see, for details, [6, p. 31, Eq. (18.1)] and [15]):

$$\begin{aligned} f(a, b) &= 1 + \sum_{n=1}^{\infty} (ab)^{\frac{n(n-1)}{2}} (a^n + b^n) \\ &= \sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}} = f(b, a), \quad (|ab| < 1), \end{aligned} \quad (5)$$

which is further expressed as

$$f(a, b) = a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}} f(a(ab)^n, b(ab)^{-n}) = f(b, a) \quad (n \in \mathbb{Z}). \quad (6)$$

Equation (6) holds true as stated only if n is any integer. Moreover, in case n is not an integer, this result (6) is only approximately true (see, for details, [12, Vol. 2, Chapter XVI, p. 193, Entry 18 (iv)]).

In fact, Ramanujan (see [12] and [13]) also rediscovered Jacobi's famous triple-product identity which, in Ramanujan's notation, is given by (see [6, p. 35, Entry 19]):

$$f(a, b) = (-a; ab)_\infty (-b; ab)_\infty (ab; ab)_\infty, \quad (7)$$

or, equivalently, by (see [10])

$$\begin{aligned} \sum_{n=-\infty}^{\infty} q^{n^2} z^n &= \prod_{n=1}^{\infty} (1 - q^{2n}) (1 + zq^{2n-1}) \left(1 + \frac{1}{z} q^{2n-1}\right) \\ &= (q^2; q^2)_\infty (-zq; q^2)_\infty \left(-\frac{q}{z}; q^2\right)_\infty, \quad (|q| < 1; z \neq 0). \end{aligned}$$

The q -series identity (7) or its above-mentioned equivalent form was first proved by Carl Friedrich Gauss (1777–1855).

Several q -series identities, which emerge naturally from Jacobi's triple-product identity (7), are worthy of note here (see, for details, [6, pp. 36–37, Entry 22]):

$$\begin{aligned} \varphi(q) &:= \sum_{n=-\infty}^{\infty} q^{n^2} = 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \\ &= \{(-q; q^2)_\infty\}^2 (q^2; q^2)_\infty = \frac{(-q; q^2)_\infty (q^2; q^2)_\infty}{(q; q^2)_\infty (-q^2; q^2)_\infty}; \end{aligned} \quad (8)$$

$$\psi(q) := f(q, q^3) = \sum_{n=0}^{\infty} q^{\frac{n(n+1)}{2}} = \frac{(q^2; q^2)_\infty}{(q; q^2)_\infty}; \quad (9)$$

and

$$\begin{aligned} f(-q) &:= \mathfrak{f}(-q, -q^2) = \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{n(3n-1)}{2}} \\ &= \sum_{n=0}^{\infty} (-1)^n q^{\frac{n(3n-1)}{2}} + \sum_{n=1}^{\infty} (-1)^n q^{\frac{n(3n+1)}{2}} = (q; q)_{\infty}. \end{aligned} \quad (10)$$

Equation (10) is known as Euler's *Pentagonal Number Theorem*. Remarkably, the following q -series identity:

$$(-q; q)_{\infty} = \frac{1}{(q; q^2)_{\infty}} = \frac{1}{\chi(-q)} \quad (11)$$

provides the analytic equivalent form of Euler's famous theorem (see, for details, [3] and [5]).

Theorem 1 (Euler's Pentagonal Number Theorem). *The number of partitions of a given positive integer n into distinct parts is equal to the number of partitions of n into odd parts.*

We also recall the Rogers-Ramanujan continued fraction $R(q)$ given by

$$\begin{aligned} R(q) &:= q^{\frac{1}{5}} \frac{H(q)}{G(q)} = q^{\frac{1}{5}} \frac{\mathfrak{f}(-q, -q^4)}{\mathfrak{f}(-q^2, -q^3)} = q^{\frac{1}{5}} \frac{(q; q^5)_{\infty} (q^4; q^5)_{\infty}}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}} \\ &= \frac{q^{\frac{1}{5}}}{1+} \frac{q}{1+} \frac{q^2}{1+} \frac{q^3}{1+}, \quad (|q| < 1). \end{aligned} \quad (12)$$

Here $G(q)$ and $H(q)$, which are associated with the widely-investigated Roger-Ramanujan identities, are defined as follows:

$$\begin{aligned} G(q) &= \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} = \frac{f(-q^5)}{\mathfrak{f}(-q, -q^4)} = \frac{1}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}} = \\ &= \frac{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty} (q^5; q^5)_{\infty}}{(q; q)_{\infty}} \end{aligned} \quad (13)$$

and

$$\begin{aligned} H(q) &:= \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q; q)_n} = \frac{f(-q^5)}{\mathfrak{f}(-q^2, -q^3)} \\ &= \frac{1}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}} = \frac{(q; q^5)_{\infty} (q^4; q^5)_{\infty} (q^5; q^5)_{\infty}}{(q; q)_{\infty}}, \end{aligned} \quad (14)$$

and the functions $\mathfrak{f}(a, b)$ and $f(-q)$ are given by the equations (5) and (10), respectively.

For a detailed historical account of (and for various related developments stemming from) the Rogers-Ramanujan continued fraction (12) as well as the Rogers-Ramanujan identities (13) and (14), the interested reader may refer to the monumental work [6, p. 77 *et seq.*] (see also [15] and [19]).

Theorem 2. Suppose that $|q| < 1$. Then

$$(q^2; q^2)_\infty (-q; q)_\infty = \frac{1}{1-} \frac{q}{1+} \frac{q(1-q)}{1-} \frac{q^3}{1+} \frac{q^2(1-q^2)}{1-} \frac{q^5}{1+} \frac{q^3(1-q^3)}{1-} \dots, \quad (15)$$

$$\frac{(q; q^5)_\infty (q^4; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty} = \frac{1}{1+} \frac{q}{1+} \frac{q^2}{1+} \frac{q^3}{1+} \frac{q^4}{1+} \frac{q^5}{1+} \frac{q^6}{1+} \dots, \quad (16)$$

and

$$C(q) = \frac{(q^2; q^5)_\infty (q^3; q^5)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} = 1 + \frac{q}{1+} \frac{q^2}{1+} \frac{q^3}{1+} \frac{q^4}{1+} \frac{q^5}{1+} \frac{q^6}{1+} \dots. \quad (17)$$

In 2015, Andrews *et al.* [4] investigated a number of interesting double-summation hypergeometric q -series representations for several families of partitions and further explored the rôle of double series in combinatorial-partition identities by introducing the general family $R(s, t, l, u, v, w)$ and also suggested its three special cases in term of multivariate R -functions. Thereafter, several new advancements and generalizations of the existing results were made in regard to combinatorial partition-theoretic identities (see, for example, [15] to [17]). An interesting recent investigation on the subject of combinatorial partition-theoretic identities by Hahn *et al.* [9] is also worth mentioning in this connection. In this paper, our main objective is to establish a set of two new identities which depict the inter-relationships in terms of R_α, R_β and R_m functions along with q -product identities.

Recently, Srivastva *et al.* (see [18]) has introduced three notations:

$$\begin{aligned} R_\alpha &= R(2, 1, 1, 1, 2, 2); & R_\beta &= R(2, 2, 1, 1, 2, 2); \\ R_m &= R(m, m, 1, 1, 1, 2). \end{aligned} \quad (18)$$

where $m = 1, 2, 3, \dots$

Each of the following preliminary results will be needed for the demonstration of our main results in this paper (see [11]):

For

$$A = \frac{f(q)f(-q^2)}{q^{\frac{1}{4}}f(q^3)f(-q^6)}, \quad C_n = \frac{f(-q^n)f(-q^{2n})}{q^{\frac{n}{4}}f(-q^{3n})f(-q^{6n})}; \quad n = 1, 2, 3, \dots$$

I. If

$$P = AC_1, \quad Q = \frac{A}{C_1},$$

then

$$\left(Q^8 + \frac{1}{Q^8}\right) + 7\left(Q^4 + \frac{1}{Q^4}\right) + \left(P^2 - \frac{9^2}{P^2}\right)\left(Q^2 - \frac{1}{Q^2}\right) - 24 = 0. \quad (19)$$

II. If

$$P = C_1C_4, \quad Q = \frac{C_1}{C_4},$$

then

$$\begin{aligned}
& \left(Q^{16} + \frac{1}{Q^{16}} \right) - 200 \left(Q^{12} + \frac{1}{Q^{12}} \right) - 6848 \left(Q^8 + \frac{1}{Q^8} \right) - 53255 \left(Q^4 + \frac{1}{Q^4} \right) \\
& - \left\{ 9320 \left(Q^2 + \frac{1}{Q^2} \right) + 2016 \left(Q^6 + \frac{1}{Q^6} \right) + 103 \left(Q^{10} + \frac{1}{Q^{10}} \right) \right\} \cdot \left(P^2 + \frac{9^2}{P^2} \right) \\
& - \left\{ 280 \left(Q^4 + \frac{1}{Q^4} \right) + 16 \left(Q^8 + \frac{1}{Q^8} \right) + (29)^2 \right\} \cdot \left(P^4 + \frac{9^4}{P^4} \right) \\
& - \left\{ 24 \left(Q^2 + \frac{1}{Q^2} \right) + \left(Q^6 + \frac{1}{Q^6} \right) \right\} \cdot \left(P^6 + \frac{9^6}{P^6} \right) - \left(P^8 + \frac{9^8}{P^8} \right) - 135032 = 0.
\end{aligned} \tag{20}$$

2. A SET OF MAIN RESULTS

In this section, we state and prove a set of two identities which depict inter-relationships among q -product identities; R_α , R_β , and R_m functions.

Theorem 3. *Each of the following relationships holds true:*

$$\begin{aligned}
& \left\{ \frac{(-q; -q)_\infty (-q^3; -q^3)_\infty}{q^{\frac{1}{2}} R_1 R_6 (-q; -q)_\infty} \right\}^2 \cdot \left\{ \frac{(q, q^2, q^2, q^2, q^2, q^2; q^2)_\infty}{(-q^3; -q^3)_\infty (q^3, q^3, q^6, q^6, q^6; q^6)_\infty (q^6, q^6; q^{12})_\infty} \right\}^2 \\
& + \left\{ \frac{9q^{\frac{1}{2}} R_1 R_6}{(-q^3; -q^3)_\infty} \right\}^2 \cdot \left\{ \frac{(-q^3; -q^3)_\infty (q^3, q^3, q^6, q^6, q^6; q^6)_\infty (q^6, q^6; q^{12})_\infty}{(q, q^2, q^2, q^2, q^2, q^2; q^2)_\infty} \right\}^2 \\
& = \{R_1 R_6 R_{12}\}^8 \cdot \left\{ \frac{(-q; -q)_\infty (q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}; q^{24})_\infty}{(-q^3; -q^3)_\infty (q^2, q^2; q^2)_\infty (q^{24}; q^{24})_\infty} \right\}^8 \\
& + \left\{ \frac{(-q^3; -q^3)_\infty (q^2, q^2; q^2)_\infty}{R_1 R_6 (-q; -q)_\infty (q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty} \right\}^8 + 7\{R_1 R_6 R_{12}\}^4 \\
& \cdot \left\{ \frac{(-q; -q)_\infty (q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}; q^{24})_\infty}{(-q^3; -q^3)_\infty (q^2, q^2; q^2)_\infty (q^{24}; q^{24})_\infty} \right\}^4 + \\
& + 7 \left\{ \frac{(-q^3; -q^3)_\infty (q^2, q^2; q^2)_\infty}{R_1 R_6 (-q; -q)_\infty (q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty} \right\}^4 + \{R_1 R_4 R_6 [R_{12}]^3\}^2 \\
& \cdot \left\{ \frac{(-q, -q; -q)_\infty (q; q^2)_\infty (q^2; q^4)_\infty^3 (q^3; q^6)_\infty (q^6; q^{12})_\infty^2 (q^{12}; q^{24})_\infty}{q^{\frac{1}{2}} (-q^3, -q^3; -q^3)_\infty (q^3, q^6, q^6, q^6; q^6)_\infty (q^{24}; q^{24})_\infty (q^8, q^8; q^8)_\infty} \right\}^2 \\
& + \left\{ \frac{9q^{\frac{1}{2}} (-q^3, -q^3; -q^3)_\infty (q^3, q^6, q^6, q^6; q^6)_\infty}{R_1 R_6 [R_{12}]^2} \right\}^2 \\
& \cdot \left\{ \frac{(q^4, q^8; q^8)_\infty}{(-q, -q; -q)_\infty (q; q^2)_\infty (q^2, q^2, q^2; q^4)_\infty (q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty} \right\}^2 - 24, \tag{21}
\end{aligned}$$

and

$$\begin{aligned}
& \{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^{16} \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^8; q^{16})_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^8, q^{16}, q^{16}; q^{16})_\infty} \right\}^{16} \\
& + \left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^{16} \\
& = 200\{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^{12} \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^{16}, q^{16}; q^{16})_\infty} \right\}^{12} \\
& + 200\left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^{12} + 6848\{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^8 \\
& \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^{16}, q^{16}; q^{16})_\infty} \right\}^8 + \\
& + 6848\left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^8 + 53255\{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^4 \\
& \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^{16}, q^{16}; q^{16})_\infty} \right\}^4 \\
& + 53255\left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^4 \\
& + 9320\left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24}; q^{24})_\infty} \right\}^2 \\
& \cdot \{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^2 \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^{16}, q^{16}; q^{16})_\infty} \right\}^2 \\
& + 9320\left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24}; q^{24})_\infty} \right\}^2 \\
& \cdot \left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^2 \\
& + 2016\left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24}; q^{24})_\infty} \right\}^2 \\
& \cdot \{q^{\frac{3}{4}}R_4R_8[R_{12}]^2\}^6 \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}; q^{24})_\infty (q^{16}, q^{16}; q^{16})_\infty} \right\}^6 \\
& + 2016\left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24}; q^{24})_\infty} \right\}^2 \\
& \cdot \left\{ \frac{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24})_\infty (q^8, q^{16}; q^{16})_\infty}{q^{\frac{3}{4}}R_4R_{12}(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty} \right\}^6 \\
& + 103\left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24}; q^{24})_\infty} \right\}^2
\end{aligned}$$

[illegible]

[illegible]

(22)

Proof. First of all, in order to prove our assertion (21), apply the identity (10) into (19), we obtain:

$$P = \{AC_1\} = \left\{ \frac{(-q; -q)_\infty (q, q^2, q^2, q^2; q^2)_\infty}{q^{\frac{1}{2}}(-q^3; -q^3)_\infty (q^3, q^6, q^6, q^6; q^{12})_\infty} \right\}, \quad (23)$$

$$Q = \left\{ \frac{A}{C_1} \right\} = \{R_1 R_6 R_{12}\} \cdot \left\{ \frac{(-q; -q)_\infty (q^3; q^6)_\infty (q^6; q^{12})_\infty^2 (q^{12}; q^{24})_\infty}{(-q^3; -q^3)_\infty (q^2, q^2, q^2)_\infty (q^{24}; q^{24})_\infty} \right\}. \quad (24)$$

combining (23) and (24), as precondition given in (19), by arranging the powers of suitable terms, we are led to the first assertion (21).

Finally. we attempt to prove our second identity (22), apply identity (10) into (20), we obtain:

$$\begin{aligned} P &= \{C_1 C_4\} = \\ &= \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4, q^4, q^8, q^8, q^8, q^8; q^8)_\infty}{q^{\frac{5}{4}}(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{12}, q^{12}, q^{12}, q^{24}, q^{24}, q^{24}, q^{24})_\infty} \right\}, \end{aligned} \quad (25)$$

and

$$Q = \left\{ \frac{C_1}{C_4} \right\} = \{q^{\frac{3}{4}} R_4 R_8 [R_{12}]^2\} \cdot \left\{ \frac{(q; q^2)_\infty (q^2, q^2; q^4)_\infty (q^4, q^4; q^8)_\infty (q^8; q^{16})_\infty (q^{12}; q^{24})_\infty}{(q^3; q^6)_\infty (q^6, q^6; q^{12})_\infty (q^{24}, q^{24}, q^{24})_\infty (q^8, q^{16}, q^{16}, q^{16})_\infty} \right\}. \quad (26)$$

combining (25) and (26), as precondition given in (20), by arranging the powers of suitable terms, we are led to the second assertion (22).

We thus have completed our proof of the above Theorem. $\square$

3. Acknowledgements

The research work of the second named author (Sangeeta Chaudhary) has been sponsored by the UGC Dr. D. S. Kothari Post Doctoral Fellowship Scheme by its letter ref No. F.4-2/2006 (BSR)/MA/18-19/0022. The authors would like to thank the referee(s) for their valuable comments and corrections which have improved the quality of this paper.

4. Conflicts of Interest

Both the authors declare that they have no conflict of interest.

REFERENCES

- [1] C. Adiga, N. A. S. Bulkhali, D. Ranganatha and H. M. Srivastava, *Some new modular relations for the Rogers-Ramanujan type functions of order eleven with applications to partitions*, Journal of Number Theory, 158 (2016), 281-297.
- [2] C. Adiga, N. A. S. Bulkhali, Y. Simsek and H. M. Srivastava, *A continued fraction of Ramanujan and some Ramanujan-Weber class invariants*, Filomat, 31 (2017), 3975-3997.
- [3] G. E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, London and New York, 1998.
- [4] G. E. Andrews, K. Bringman and K. Mahlburg, *Double series representations for Schur's partition function and related identities*, Journal of Combinatorial Theory, Series A, 132 (2015), 102-119.
- [5] T. M. Apostol, *Introduction to Analytic Number Theory*, Undergraduate Texts in Mathematics, Springer-Verlag, Berlin, New York and Heidelberg, 1976.
- [6] B. C. Berndt, *Ramanujan's Notebooks*, Part III, Springer-Verlag, Berlin, Heidelberg and New York, 1991.
- [7] J. Cao, H. M. Srivastava and Z.-G. Luo, *Some iterated fractional q -integrals and their applications*, Fractional Calculus and Applied Analysis, 21 (2018), 672-695.
- [8] M. P. Chaudhary, Sangeeta Chaudhary and Sonajharia Minz, *On relationships between q -products identities and combinatorial partition identities*, Mathematica Moravica, 24 (2020), 83-91.
- [9] H.-Y. Hahn, J.-S. Huh, E.-S. Lim and J.-B. Sohn, *From partition identities to a combinatorial approach to explicit Satake inversion*, Annals of Combinatorics, 22 (2018), 543-562.
- [10] C. G. J. Jacobi, *Fundamenta Nova Theoriae Functionum Ellipticarum*, Regiomonti, Sumtibus Fratrum Bornträger, Königsberg, Germany, 1829; Reprinted in *Gesammelte Mathematische Werke*, 1 (1829), 497-538; American Mathematical Society, Providence, Rhode Island, 1969, 97-239.
- [11] M. S. M. Naika, K. S. Bairy and N. P. Suman, *Modular relations for remarkable product of theta-functions and evaluations*, Recent Advances in Mathematics, RMS-Lecture Notes Series, 21 (2015), 133-145.
- [12] S. Ramanujan, *Notebooks*, Vols. 1 and 2, Tata Institute of Fundamental Research, Bombay, 1957.
- [13] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa Publishing House, New Delhi, 1988.
- [14] H. M. Srivastava, *Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis*, Iranian Journal of Science and Technology, Transaction A: Science, 44 (2020), 327-344.
- [15] H. M. Srivastava and M. P. Chaudhary, *Some relationships between q -product identities, combinatorial partition identities and continued-fraction identities*, Advanced Studies in Contemporary Mathematics, 25 (2015), 265-272.
- [16] H. M. Srivastava, M. P. Chaudhary and S. Chaudhary, *Some theta-function identities related to Jacobi's triple-product identity*, European Journal of Pure and Applied Mathematics, 11 (1) (2018), 1-9.

- [17] H. M. Srivastava, M. P. Chaudhary and S. Chaudhary, *A family of theta-function identities related to Jacobi's triple-product identity*, Russian Journal of Mathematical Physics, 27 (2020), 139-144.
- [18] H. M. Srivastava, R. Srivastava, M.P. Chaudhary and S. Uddin, *A family of theta function identities based upon combinatorial partition identities and related to Jacobi's triple-product identity*, Mathematics, 8 (6) (2020), Article ID: 918, 1-14.
- [19] H. M. Srivastava and J. Choi, *Zeta and q -Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.
- [20] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.
- [21] H. M. Srivastava and C.-H. Zhang, *A certain class of identities of the Rogers-Ramanujan type*, Pan-American Mathematical Journal, 19 (2009), 89-102.
- [22] A.-J. Yee, *Combinatorial proofs of generating function identities for F -partitions*, Journal of Combinatorial Theory, Series A, 102 (2003), 217-228.
- [23] J.-H. Yi, *Theta-function identities and the explicit formulas for theta-function and their applications*, Journal of Mathematical Analysis and Applications, 292 (2004), 381-400.

M. P. CHAUDHARY

DEPARTMENT OF MATHEMATICS

NETAJI SUBHAS UNIVERSITY OF TECHNOLOGY

NEW DELHI 110078

INDIA

E-mail address: dr.m.p.chaudhary@gmail.com

SANGEETA CHAUDHARY

SCHOOL OF COMPUTATER AND SYSTEMS SCIENCES

JAWAHARLAL NEHRU UNIVERSITY

NEW DELHI 110067

INDIA

E-mail address: sangeeta.ch289@gmail.com