

IDEALS OF IS-ALGEBRAS BASED ON \mathcal{N} -STRUCTURES

HASHEM BORDBAR^{1*}, MOHAMMAD MEHDI ZAHEDI², AND YOUNG BAE JUN³

ABSTRACT. The notion of a left (resp., right) \mathcal{N}_j -ideal is introduced, and related properties are investigated. Characterizations of a left (resp., right) \mathcal{N}_j -ideal are considered. Translations of a left (resp., right) \mathcal{N}_j -ideal are studied. We show that the homomorphic image (preimage) of a left (resp., right) \mathcal{N}_j -ideal is a left (resp., right) \mathcal{N}_j -ideal. The notion of retrenched left (resp., right) \mathcal{N}_j -ideals is introduced, and their properties are investigated.

1. INTRODUCTION

Most of the generalization of the crisp set have been conducted on the unit interval $[0, 1]$ and they are consistent with the asymmetry observation because a (crisp) set A in a universe X can be defined in the form of its characteristic function $\mu_A : X \rightarrow \{0, 1\}$ yielding the value 1 for elements belonging to the set A and the value 0 for elements excluded from the set A . In other words, the generalization of the crisp set to fuzzy sets relied on spreading positive information that fit the crisp point $\{1\}$ into the interval $[0, 1]$. Because no negative meaning of information is suggested, we now feel a need to deal with negative information. To do so, we also feel a need to supply mathematical tool. To attain such object, Jun et al. [3] introduced a new function which is called negative-valued function, and constructed \mathcal{N} -structures. They applied \mathcal{N} -structures to BCK/BCI-algebras, and discussed \mathcal{N} -subalgebras and \mathcal{N} -ideals in BCK/BCI-algebras. The \mathcal{N} -structures are applied to BE -algebras and subtraction algebras (see [1] and [5]).

In this paper, using the \mathcal{N} -structures, we introduce the notion of a left (resp., right) \mathcal{N}_j -ideal, and investigate related properties. We consider characterizations of a left

Key words and phrases. Left (resp., right) \mathcal{N}_j -ideal, translation, closed support, retrenched left (resp., right) \mathcal{N}_j -ideal.

2010 *Mathematics Subject Classification.* Primary: 06F35. Secondary: 03G25, 03B52.

Received: November 06, 2016.

Accepted: August 26, 2017.

(resp., right) \mathcal{N}_j -ideal, and study translations of a left (resp., right) \mathcal{N}_j -ideal. We show that the homomorphic image (preimage) of a left (resp., right) \mathcal{N}_j -ideal is a left (resp., right) \mathcal{N}_j -ideal. We also introduction the notion of retrenched left (resp., right) \mathcal{N}_j -ideals and investigate their properties.

2. PRELIMINARIES

Let $K(\tau)$ be the class of all algebras with type $\tau = (2, 0)$. By a *BCI-algebra* we mean a system $X := (X, *, \theta) \in K(\tau)$ in which the following axioms hold:

- (i) $((x * y) * (x * z)) * (z * y) = \theta$;
- (ii) $(x * (x * y)) * y = \theta$;
- (iii) $x * x = \theta$;
- (iv) $x * y = y * x = \theta \Rightarrow x = y$;

for all $x, y, z \in X$. If a BCI-algebra X satisfies $\theta * x = \theta$ for all $x \in X$, then we say that X is a *BCK-algebra*. We can define a partial ordering \preceq by

$$(\forall x, y \in X) (x \preceq y \Rightarrow x * y = \theta).$$

In a BCK/BCI-algebra X , the following hold:

$$(2.1) \quad (\forall x \in X) (x * \theta = x),$$

$$(2.2) \quad (\forall x, y, z \in X) ((x * y) * z = (x * z) * y).$$

A subset I of a BCK/BCI-algebra X is called an *ideal* of X if it satisfies

$$(I1) \quad 0 \in I;$$

$$(I2) \quad (\forall x, y \in X) (x * y \in I, y \in I \Rightarrow x \in I).$$

We refer the reader to the books [2] and [6] for further information regarding BCK/BCI-algebras.

An **IS-algebra** (see [4]) is a non-empty set X with two binary operations “ $*$ ” and “ \cdot ” and constant θ satisfying the conditions:

- $I(X) := (X, *, \theta)$ is a BCI-algebra;
- $S(X) := (X, \cdot)$ is a semigroup;
- the operation “ \cdot ” is distributive (on both sides) over the operation “ $*$ ”, that is,

$$x \cdot (y * z) = (x \cdot y) * (x \cdot z) \quad \text{and} \quad (x * y) \cdot z = (x \cdot z) * (y \cdot z),$$

for all $x, y, z \in X$.

In an **IS-algebra** X , the following hold:

$$(2.3) \quad (\forall x \in X) (\theta x = x\theta = \theta);$$

$$(2.4) \quad (\forall x, y, z \in X) (x \preceq y \Rightarrow xz \preceq yz, zx \preceq zy).$$

In what follows we use the notation xy instead of $x \cdot y$.

A nonempty subset A of an **IS-algebra** X is called a *left* (resp., *right*) *\mathcal{I} -ideal* of X (see [4]) if

- (i) A is a left (resp., right) stable subset of $S(X)$, that is, $xa \in A$ (resp., $ax \in A$) whenever $x \in S(X)$ and $a \in A$;
- (ii) $(\forall x, y \in I(X)) (x * y \in A, y \in A \Rightarrow x \in A)$.

For any family $\{a_i \mid i \in \Lambda\}$ of real numbers, we define

$$\bigvee \{a_i \mid i \in \Lambda\} := \begin{cases} \max\{a_i \mid i \in \Lambda\}, & \text{if } \Lambda \text{ is finite,} \\ \sup\{a_i \mid i \in \Lambda\}, & \text{otherwise.} \end{cases}$$

$$\bigwedge \{a_i \mid i \in \Lambda\} := \begin{cases} \min\{a_i \mid i \in \Lambda\}, & \text{if } \Lambda \text{ is finite,} \\ \inf\{a_i \mid i \in \Lambda\}, & \text{otherwise.} \end{cases}$$

3. IDEALS BASED ON \mathcal{N} -STRUCTURES

Denote by $\mathcal{F}(X, [-1, 0])$ the collection of functions from a set X to $[-1, 0]$. We say that an element of $\mathcal{F}(X, [-1, 0])$ is a *negative-valued function* from X to $[-1, 0]$ (briefly, \mathcal{N} -function on X). By an \mathcal{N} -structure we mean an ordered pair (X, f) of X and an \mathcal{N} -function f on X . In what follows, let X denote an **IS**-algebra unless otherwise specified.

Definition 3.1. An \mathcal{N} -structure (X, f) is said to satisfy the *left* (resp., *right*) *condition* in $S(X)$ if $f(xy) \leq f(y)$ (resp., $f(xy) \leq f(x)$) for all x and y in $S(X)$.

Definition 3.2. An \mathcal{N} -structure (X, f) is called a *left* (resp., *right*) \mathcal{N}_j -ideal of X if (X, f) satisfies the left (resp., right) condition in $S(X)$ and

$$(3.1) \quad (\forall x, y \in X) \left(f(\theta) \leq f(x) \leq \bigvee \{f(x * y), f(y)\} \right).$$

Example 3.1. Define two binary operations “ $*$ ” and “ \cdot ” on a set $X = \{\theta, a, b, c\}$ as follows:

$*$	θ	a	b	c
θ	θ	θ	c	b
a	a	θ	c	b
b	b	b	θ	c
c	c	c	b	θ

\cdot	θ	a	b	c
θ	θ	θ	θ	θ
a	θ	θ	θ	θ
b	θ	θ	b	c
c	θ	θ	c	b

Then X is an **IS**-algebra (see [4]). Let (X, f) be an \mathcal{N} -structure in which f is given as follows:

$$f = \begin{pmatrix} \theta & a & b & c \\ -0.8 & -0.6 & -0.3 & -0.3 \end{pmatrix}.$$

It is routine to verify that (X, f) is both a left and a right \mathcal{N}_j -ideal of X .

We provide characterizations of a left (resp., right) \mathcal{N}_j -ideal.

Theorem 3.1. *An \mathcal{N} -structure (X, f) is a left \mathcal{N}_j -ideal of X if and only if the following assertions are valid*

$$(3.2) \quad (\forall x, y \in X) (f(xy) \leq f(y)),$$

$$(3.3) \quad (\forall x, y \in X) \left(f(x) \leq \bigvee \{f(x * y), f(y)\} \right).$$

Proof. The necessity is clear. Assume that (X, f) satisfies two conditions (3.2) and (3.3). Using (2.3) and (3.2) induce $f(\theta) = f(\theta y) \leq f(y)$ for all $y \in X$. Hence (X, f) is a left \mathcal{N}_j -ideal of X . \square

Similarly we have the following theorem.

Theorem 3.2. *An \mathcal{N} -structure (X, f) is a right \mathcal{N}_j -ideal of X if and only if f satisfies the condition (3.3) and*

$$(3.4) \quad (\forall x, y \in X) \quad (f(xy) \leq f(x)).$$

For any \mathcal{N} -structure (X, f) and $t \in [-1, 0)$, the set

$$C(f; t) := \{x \in X \mid f(x) \leq t\}$$

is called a *closed t -support* of (X, f) (see [3]).

Theorem 3.3. *If an \mathcal{N} -structure (X, f) is a left \mathcal{N}_j -ideal of X , then the closed t -support of (X, f) is a left \mathcal{J} -ideal of X for all $t \in [f(\theta), 0]$.*

Proof. Let $x \in S(X)$ and $a \in C(f; t)$ for $t \in [f(\theta), 0]$. Then $f(a) \leq t$, and so $f(xa) \leq f(a) \leq t$ which shows that $xa \in C(f; t)$. It follows from (2.3) that $\theta = \theta a \in C(f; t)$. Let $x, y \in X$ be such that $x * y \in C(f; t)$ and $y \in C(f; t)$ for $t \in [f(\theta), 0]$. Then $f(x * y) \leq t$ and $f(y) \leq t$. It follows from (3.3) that

$$f(x) \leq \bigvee \{f(x * y), f(y)\} \leq t$$

and so that $x \in C(f; t)$. Therefore $C(f; t)$ is an \mathcal{J} -ideal of X for all $t \in [f(\theta), 0]$. \square

Theorem 3.4. *If an \mathcal{N} -structure (X, f) is a right \mathcal{N}_j -ideal of X , then the closed t -support of (X, f) is a right \mathcal{J} -ideal of X for all $t \in [f(\theta), 0]$.*

Proof. It is similar to the proof of Theorem 3.3. \square

Theorem 3.5. *Given an \mathcal{N} -structure (X, f) , if the nonempty closed t -support of (X, f) is a left \mathcal{J} -ideal of X for all $t \in [-1, 0)$, then (X, f) is a left \mathcal{N}_j -ideal of X .*

Proof. Assume that $C(f; t)$ is a left \mathcal{J} -ideal of X for all $t \in [-1, 0)$ with $C(f; t) \neq \emptyset$. If $f(ab) > f(b)$ for some $a, b \in X$, then there exists $t \in [-1, 0)$ such that $f(ab) > t \geq f(b)$. It follows that $b \in C(f; t)$ and $ab \notin C(f; t)$, which is a contradiction. Hence (3.2) is valid. Now suppose that (3.3) is false. Then there exists $a, b \in X$ such that

$$f(a) > \bigvee \{f(a * b), f(b)\}.$$

Taking $t := \frac{1}{2}(f(a) + \bigvee \{f(a * b), f(b)\})$ implies that $a * b \in C(f; t)$, $b \in C(f; t)$ and $a \notin C(f; t)$. This is a contradiction, and so (3.3) is valid. Therefore (X, f) is a left \mathcal{N}_j -ideal of X by Theorem 3.1. \square

Similarly we have the following theorem.

Theorem 3.6. *Given an \mathcal{N} -structure (X, f) , if the nonempty closed t -support of (X, f) is a right \mathcal{J} -ideal of X for all $t \in [-1, 0)$, then (X, f) is a right \mathcal{N}_j -ideal of X .*

Theorem 3.7. *For any left \mathcal{J} -ideal A of X and any fixed number t in an open interval $(-1, 0)$, there exists a left $\mathcal{N}_{\mathcal{J}}$ -ideal (X, f) of X on which A is the closed t -support of (X, f) .*

Proof. Let (X, f) be an \mathcal{N} -structure on which f is given as follows:

$$f(x) = \begin{cases} t, & \text{if } x \in A, \\ 0, & \text{if } x \notin A. \end{cases}$$

Let $x, y \in X$. If $y \notin A$, then $f(y) = 0$ and thus

$$f(x) \leq 0 = \vee\{f(x * y), f(y)\}.$$

Assume that $y \in A$. If $x \in A$, then $x * y$ may or may not belong to A . In any case, we have

$$f(x) \leq \vee\{f(x * y), f(y)\}.$$

If $x \notin A$, then $x * y \notin A$ and hence

$$f(x) = 0 = \vee\{f(x * y), f(y)\}.$$

For any $x, y \in X$, if $y \in A$ then $xy \in A$. Hence $f(xy) = t = f(y)$. If $y \notin A$, then $f(y) = 0$ and so $f(xy) \leq 0 = f(y)$. It follows from Theorem 3.1 that (X, f) is a left $\mathcal{N}_{\mathcal{J}}$ -ideal of X . Obviously, $A = C(f; t)$. \square

Similarly, we have the following theorem.

Theorem 3.8. *For any right \mathcal{J} -ideal A of X and any fixed number t in an open interval $(-1, 0)$, there exists a right $\mathcal{N}_{\mathcal{J}}$ -ideal (X, f) of X on which A is the closed t -support of (X, f) .*

Theorem 3.9. *For any nonempty subset A of X and $t \in [-1, 0)$, let (X, f) be an \mathcal{N} -structure on which f is given as follows:*

$$f(x) = \begin{cases} t, & \text{if } x \in A, \\ 0, & \text{if } x \notin A. \end{cases}$$

If A is a left (resp., right) \mathcal{J} -ideal of X , then (X, f) is a left (resp., right) $\mathcal{N}_{\mathcal{J}}$ -ideal of X .

Proof. Suppose that A is a left \mathcal{J} -ideal of X . Let $x, y \in X$. If $y \in A$, then $xy \in A$, and

(i) $x * y$ may or may not belong to A whenever $x \in A$;

(ii) $x * y \notin A$ whenever $x \notin A$.

Hence $f(xy) = t = f(y)$ and $f(x * y) \leq \vee\{f(x * y), f(y)\}$. If $y \notin A$, then $f(xy) \leq 0 = f(y)$ and $f(x * y) \leq 0 = \vee\{f(x * y), f(y)\}$. Therefore (X, f) is a left $\mathcal{N}_{\mathcal{J}}$ -ideal of X by Theorem 3.1. Similarly we can prove it for the right case. \square

Corollary 3.1. *For any nonempty subset A of X and an \mathcal{N} -structure (X, f) with $\text{Im}(f) = \{-1, 0\}$, the following assertions are equivalent.*

- (1) A is a left (resp., right) \mathcal{J} -ideal of X .
 (2) (X, f) is a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal of X .

Theorem 3.10. *If an \mathcal{N} -structure (X, f) is a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal of X , then the set*

$$X_f := \{x \in X \mid f(x) = f(\theta)\}$$

is a left (resp., right) \mathcal{J} -ideal of X .

Proof. Assume that (X, f) is a left $\mathcal{N}_\mathcal{J}$ -ideal of X and let $x, y \in X$. If $y \in X_f$, then $f(xy) \leq f(y) = f(\theta)$ and so $f(xy) = f(\theta)$, that is, $xy \in X_f$. Obviously, $\theta \in X_f$. Suppose that $x * y \in X_f$ and $y \in X_f$. Then

$$f(x) \leq \bigvee \{f(x * y), f(y)\} = f(\theta),$$

and so $f(x) = f(\theta)$, i.e., $x \in X_f$. Therefore X_f is a left \mathcal{J} -ideal of X . Similarly, we can prove it for the right case. \square

Given an \mathcal{N} -structure (X, f) , we denote

$$\perp := -1 - \bigwedge \{f(x) \mid x \in X\}.$$

For any $\alpha \in [\perp, 0]$, the α -translation of (X, f) is defined to be the new \mathcal{N} -structure (X, f_α) on which f_α is defined by $f_\alpha(x) = f(x) + \alpha$ for all $x \in X$.

Theorem 3.11. *For every $\alpha \in [\perp, 0]$, the α -translation of a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal is a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal of X .*

Proof. Let $\alpha \in [\perp, 0]$ and let (X, f) be a left $\mathcal{N}_\mathcal{J}$ -ideal of X . For any $x, y \in X$, we have $f_\alpha(xy) = f(xy) + \alpha \leq f(y) + \alpha = f_\alpha(y)$ and

$$\begin{aligned} f_\alpha(x) &= f(x) + \alpha \leq \bigvee \{f(x * y), f(y)\} + \alpha \\ &= \bigvee \{f(x * y) + \alpha, f(y) + \alpha\} \\ &= \bigvee \{f_\alpha(x * y), f_\alpha(y)\}. \end{aligned}$$

It follows from Theorem 3.1 that (X, f_α) is a left $\mathcal{N}_\mathcal{J}$ -ideal of X . For the right case, it is similar. \square

Theorem 3.12. *For \mathcal{N} -structure (X, f) , if there exists $\alpha \in [\perp, 0]$ such that every α -translation of (X, f) is a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal, then (X, f) is a left (resp., right) $\mathcal{N}_\mathcal{J}$ -ideal of X .*

Proof. Assume that the α -translation (X, f_α) of (X, f) is a left $\mathcal{N}_\mathcal{J}$ -ideal of X . For any $x, y \in X$, we have $f(xy) + \alpha = f_\alpha(xy) \leq f_\alpha(y) = f(y) + \alpha$ and

$$\begin{aligned} f(x) + \alpha &= f_\alpha(x) \leq \bigvee \{f_\alpha(x * y), f_\alpha(y)\} \\ &= \bigvee \{f(x * y) + \alpha, f(y) + \alpha\} \\ &= \bigvee \{f(x * y), f(y)\} + \alpha. \end{aligned}$$

It follows that $f(xy) \leq f(y)$ and $f(x) \leq \bigvee\{f(x * y), f(y)\}$. Therefore (X, f) is a left \mathcal{N}_J -ideal of X by Theorem 3.1. \square

For any \mathcal{N} -structure (X, f) , $\alpha \in [\perp, 0]$ and $t \in [-1, \alpha)$, the set

$$C_\alpha(f; t) := \{x \in X \mid f(x) \leq t - \alpha\}$$

is called the α -translation of closed t -support of (X, f)

Theorem 3.13. *Let (X, f) be an \mathcal{N} -structure and $\alpha \in [\perp, 0]$. If (X, f) is a left (resp., right) \mathcal{N}_J -ideal of X , then the α -translation of closed t -support of (X, f) is a left (resp., right) \mathcal{J} -ideal of X for all $t \in [-1, \alpha)$.*

Proof. Let $x, y \in X$. If $y \in C_\alpha(f; t)$, then $f(y) \leq t - \alpha$ and so

$$(3.5) \quad f(xy) \leq f(y) \leq t - \alpha.$$

Thus $xy \in C_\alpha(f; t)$. Suppose that $x * y \in C_\alpha(f; t)$ and $y \in C_\alpha(f; t)$. Then

$$f(\theta) \leq f(x) \leq \bigvee\{f(x * y), f(y)\} \leq t - \alpha$$

by (3.1). Thus $\theta \in C_\alpha(f; t)$ and $x \in C_\alpha(f; t)$. Consequently, $C_\alpha(f; t)$ is a left \mathcal{J} -ideal of X for all $t \in [-1, \alpha)$. Similarly we can prove it for the right case. \square

Theorem 3.14. *For any \mathcal{N} -structure (X, f) and $\alpha \in [\perp, 0]$, the following assertions are equivalent.*

- (1) *The α -translation of closed t -support of (X, f) is a left (resp., right) \mathcal{J} -ideal of X for all $t \in [-1, \alpha)$.*
- (2) *The α -translation of (X, f) is a left (resp., right) \mathcal{N}_J -ideal of X .*

Proof. Suppose that (X, f_α) is a left \mathcal{N}_J -ideal of X for $\alpha \in [\perp, 0]$ and let $t \in [-1, \alpha)$. For any $x, y \in X$, if $x * y \in C_\alpha(f; t)$ and $y \in C_\alpha(f; t)$, then

$$\begin{aligned} f(x) + \alpha &= f_\alpha(x) \leq \bigvee\{f_\alpha(x * y), f_\alpha(y)\} \\ &= \bigvee\{f(x * y) + \alpha, f(y) + \alpha\} \\ &= \bigvee\{f(x * y), f(y)\} + \alpha \\ &\leq t - \alpha + \alpha = t \end{aligned}$$

and so $f(x) \leq t - \alpha$. Thus $x \in C_\alpha(f; t)$. Since

$$f(\theta) + \alpha = f_\alpha(\theta) \leq f_\alpha(x) = f(x) + \alpha \leq t - \alpha + \alpha = t,$$

for any $x \in C_\alpha(f; t)$, we have $f(\theta) \leq t - \alpha$, i.e., $\theta \in C_\alpha(f; t)$. Now if $y \in C_\alpha(f; t)$, then $f(y) \leq t - \alpha$ which implies that

$$f(xy) + \alpha = f_\alpha(xy) \leq f_\alpha(y) = f(y) + \alpha \leq t,$$

that is, $f(xy) \leq t - \alpha$ for all $x \in X$. Hence $xy \in C_\alpha(f; t)$, and therefore $C_\alpha(f; t)$ is a left \mathcal{J} -ideal of X .

Conversely, assume that the α -translation of closed t -support of (X, f) is a left \mathcal{J} -ideal of X for all $t \in [-1, \alpha)$. Suppose that there exist $a, b \in X$ and $t_0 \in [-1, \alpha)$

such that $f_\alpha(ab) > t_0 \geq f_\alpha(b)$. Then $f(ab) + \alpha > t_0$ and $f(b) + \alpha \leq t_0$, which imply that $b \in C_\alpha(f; t_0)$ and $ab \notin C_\alpha(f; t_0)$. This is a contradiction, and thus $f_\alpha(xy) \leq f_\alpha(y)$ for all $x, y \in X$. If

$$f_\alpha(a) > \bigvee\{f_\alpha(a * b), f_\alpha(b)\},$$

for some $a, b \in X$, then there exists $t_1 \in [-1, \alpha)$ such that

$$f_\alpha(a) > t_1 \geq \bigvee\{f_\alpha(a * b), f_\alpha(b)\},$$

which implies that $f(a) > t_1 - \alpha$, $f(a * b) \leq t_1 - \alpha$ and $f(b) \leq t_1 - \alpha$. Hence $a * b \in C_\alpha(f; t_1)$ and $b \in C_\alpha(f; t_1)$, but $a \notin C_\alpha(f; t_1)$, which is a contradiction. Hence $f_\alpha(x) \leq \bigvee\{f_\alpha(x * y), f_\alpha(y)\}$ for all $x, y \in X$. Therefore (X, f_α) is a left \mathcal{N}_j -ideal of X by Theorem 3.1. \square

Given two \mathcal{N} -structures (X, f) and (X, g) , we say that (X, f) is a *retrenchment* of (X, g) if $f \subseteq g$, that is, $f(x) \leq g(x)$ for all $x \in X$.

Definition 3.3. Given two \mathcal{N} -structures (X, f) and (X, g) , we say that (X, f) is a *retrenched left (resp., right) \mathcal{N}_j -ideal* of (X, g) , denoted by

$$(X, f) \tilde{\subseteq}_l (X, g) \text{ (resp., } (X, f) \tilde{\subseteq}_r (X, g)),$$

if (X, f) is a retrenchment of (X, g) , and (X, f) is a left (resp., right) \mathcal{N}_j -ideal of X whenever (X, g) is a left (resp., right) \mathcal{N}_j -ideal of X .

Theorem 3.15. *Let (X, g) be a left (resp., right) \mathcal{N}_j -ideal of X . For every $\alpha \in [\perp, 0]$, the α -translation (X, g_α) of (X, g) is a retrenched left (resp., right) \mathcal{N}_j -ideal of X .*

Proof. For any $x \in X$, we have $g_\alpha(x) = g(x) + \alpha \leq g(x)$. Thus (X, g_α) is a retrenchment of (X, g) . If (X, g) is a left \mathcal{N}_j -ideal of X , then Theorem 3.11 shows that (X, g_α) is a left \mathcal{N}_j -ideal of X . Therefore (X, g_α) is a retrenched left \mathcal{N}_j -ideal of X . Similarly, we can prove it for the right case. \square

Theorem 3.16. *Let (X, g) be a left (resp., right) \mathcal{N}_j -ideal of X . If (X, f_1) and (X, f_2) are retrenched left (resp., right) \mathcal{N}_j -ideals of (X, g) , then so is $(X, f_1 \cup f_2)$, where $(f_1 \cup f_2)(x) = \bigvee\{f_1(x), f_2(x)\}$ for all $x \in X$.*

Proof. Assume that (X, f_1) and (X, f_2) are retrenched left \mathcal{N}_j -ideals of a left \mathcal{N}_j -ideal (X, g) of X . Then $f_1(x) \leq g(x)$ and $f_2(x) \leq g(x)$, for all $x \in X$. Thus $(f_1 \cup f_2)(x) = \bigvee\{f_1(x), f_2(x)\} \leq g(x)$ for all $x \in X$, and so $(X, f_1 \cup f_2)$ is a retrenchment of (X, g) . For any $x, y \in X$, we have

$$\begin{aligned} (f_1 \cup f_2)(xy) &= \bigvee\{f_1(xy), f_2(xy)\} \\ &\leq \bigvee\{f_1(y), f_2(y)\} \\ &= (f_1 \cup f_2)(y) \end{aligned}$$

and

$$\begin{aligned} (f_1 \cup f_2)(x) &= \bigvee \{f_1(x), f_2(x)\} \\ &\leq \bigvee \left\{ \bigvee \{f_1(x * y), f_1(y)\}, \bigvee \{f_2(x * y), f_2(y)\} \right\} \\ &= \bigvee \left\{ \bigvee \{f_1(x * y), f_2(x * y)\}, \bigvee \{f_1(y), f_2(y)\} \right\} \\ &= \bigvee \{(f_1 \cup f_2)(x * y), (f_1 \cup f_2)(y)\}. \end{aligned}$$

It follows from Theorem 3.1 that $(X, f_1 \cup f_2)$ is a left \mathcal{N}_j -ideal of X . Therefore $(X, f_1 \cup f_2)$ is a retrenched left \mathcal{N}_j -ideal of (X, g) . The proof is similar for the right case. \square

Theorem 3.17. *Let (X, g) be a left \mathcal{N}_j -ideal of X and let $\alpha, \beta \in [\perp, 0]$. If $\alpha \leq \beta$, then the α -translation (X, g_α) of (X, g) is a retrenched left \mathcal{N}_j -ideal of the β -translation (X, g_β) of (X, g) .*

Proof. Note that the α -translation (X, g_α) and the β -translation (X, g_β) of (X, g) are left \mathcal{N}_j -ideal of X by Theorem 3.11. If $\alpha \leq \beta$, then

$$g_\alpha(x) = g(x) + \alpha \leq g(x) + \beta = g_\beta(x),$$

for all $x \in X$. Hence (X, g_α) is a retrenchment of (X, g_β) . Therefore (X, g_α) is a retrenched left \mathcal{N}_j -ideal of (X, g_β) . \square

Similarly we have the following theorem for the right case.

Theorem 3.18. *Let (X, g) be a right \mathcal{N}_j -ideal of X and let $\alpha, \beta \in [\perp, 0]$. If $\alpha \leq \beta$, then the α -translation (X, g_α) of (X, g) is a retrenched right \mathcal{N}_j -ideal of the β -translation (X, g_β) of (X, g) .*

Theorem 3.19. *Let (X, g) be a left (resp., right) \mathcal{N}_j -ideal of X and let $\beta \in [\perp, 0]$. For every retrenched left (resp., right) \mathcal{N}_j -ideal (X, f) of the β -translation (X, g_β) of (X, g) , there exists $\alpha \in [\perp, 0]$ such that $\alpha \leq \beta$ and (X, f) is a retrenched left (resp., right) \mathcal{N}_j -ideal of the α -translation (X, g_α) of (X, g) .*

Proof. It is straightforward. \square

A mapping $\varphi : X \rightarrow Y$ is called a *homomorphism* of **IS**-algebras if $\varphi(x * y) = \varphi(x) * \varphi(y)$ and $\varphi(xy) = \varphi(x)\varphi(y)$, for all $x, y \in X$.

Let $\varphi : X \rightarrow Y$ be an onto mapping. Given an \mathcal{N} -structure (Y, g) , the \mathcal{N} -structure (X, f) , where $f = g \circ \varphi$, is called the *preimage* of (Y, g) under φ . Given an \mathcal{N} -structure (X, f) , the *image* of (X, f) under φ is defined to be the \mathcal{N} -structure (Y, g) on which g is denoted by $\varphi(f)$ and is given by

$$g(y) = \bigwedge_{x \in \varphi^{-1}(y)} f(x),$$

for all $y \in Y$.

Theorem 3.20. *Every preimage of a left (resp., right) \mathcal{N}_j -ideal under onto homomorphism is a left (resp., right) \mathcal{N}_j -ideal.*

Proof. Let $\varphi : X \rightarrow Y$ be an onto homomorphism of **IS**-algebras and let an \mathcal{N} -structure (X, f) is the preimage of a left \mathcal{N}_j -ideal (Y, g) of Y . For any $x, y \in X$, we have

$$\begin{aligned} f(xy) &= (g \circ \varphi)(xy) = g(\varphi(xy)) \\ &= g(\varphi(x)\varphi(y)) \leq g(\varphi(y)) \\ &= (g \circ \varphi)(y) = f(y) \end{aligned}$$

and

$$\begin{aligned} f(x) &= (g \circ \varphi)(x) = g(\varphi(x)) \\ &\leq \bigvee \{g(\varphi(x) * y'), g(y')\} \text{ for all } y' \in Y \\ &= \bigvee \{g(\varphi(x) * \varphi(y)), g(\varphi(y))\} \\ &= \bigvee \{g(\varphi(x * y)), g(\varphi(y))\} \\ &= \bigvee \{(g \circ \varphi)(x * y), (g \circ \varphi)(y)\} \\ &= \bigvee \{f(x * y), f(y)\}. \end{aligned}$$

It follows from Theorem 3.1 that (X, f) is a left \mathcal{N}_j -ideal of X . Similarly we can verify it for the right case. \square

Lemma 3.1. *Let $\varphi : X \rightarrow Y$ be an onto mapping. Given an \mathcal{N} -structure (X, f) and $t \in [-1, 0)$, we have*

$$C(\varphi(f); t) = \bigcap_{t < s < 0} \varphi(C(f; t - s)).$$

Proof. For any $y = f(x) \in Y$, if $y \in C(\varphi(f); t)$, then

$$\bigwedge_{z \in \varphi^{-1}(\varphi(x))} f(z) = \varphi(f)(\varphi(x)) = \varphi(f)(y) \leq t.$$

Hence, for every $s \in (t, 0)$, there exists $x_0 \in \varphi^{-1}(y)$ such that $f(x_0) \leq t - s$. Thus $y = \varphi(x_0) \in \varphi(C(f; t - s))$, and so $y \in \bigcap_{t < s < 0} \varphi(C(f; t - s))$.

Conversely, let $y \in \bigcap_{t < s < 0} \varphi(C(f; t - s))$. Then $y \in \varphi(C(f; t - s))$ for every $s \in (t, 0)$, and hence there exists $x_0 \in C(f; t - s)$ such that $y = \varphi(x_0)$. It follows that $f(x_0) \leq t - s$ and $x_0 \in \varphi^{-1}(y)$. Therefore

$$\varphi(f)(y) = \bigwedge_{x \in \varphi^{-1}(y)} f(x) \leq \bigwedge_{t < s < 0} \{t - s\} = t,$$

and thus $y \in C(\varphi(f); t)$. \square

Theorem 3.21. *Every image of a left (resp., right) \mathcal{N}_j -ideal under onto homomorphism is a left (resp., right) \mathcal{N}_j -ideal.*

Proof. Let $\varphi : X \rightarrow Y$ be an onto homomorphism of **IS**-algebras and let an \mathcal{N} -structure (Y, g) is the image of a left \mathcal{N}_J -ideal (X, f) of X . Let $t \in [-1, 0)$ be such that $C(\varphi(f); t) \neq \emptyset$. Then

$$C(\varphi(f); t) = \bigcap_{t < s < 0} \varphi(C(f; t - s)),$$

by Lemma 3.1, and so $\varphi(C(f; t - s))$ is nonempty for all $s \in (t, 0)$. Since (X, f) is a left \mathcal{N}_J -ideal of X , $C(f; t - s)$ is a left \mathcal{J} -ideal of X and so the onto homomorphic image $\varphi(C(f; t - s))$ of $C(f; t - s)$ under φ is a left \mathcal{J} -ideal of Y . Hence $C(\varphi(f); t)$ is a left \mathcal{J} -ideal of Y . It follows from Theorem 3.5 that (Y, g) is a left \mathcal{N}_J -ideal of Y . \square

Acknowledgements. The research is supported by the Iranian National Science Foundation (Grant No. 96008529)

REFERENCES

- [1] S. Abdullah and A. F. Ali, *Applications of \mathcal{N} -structures in implicative filters of BE-algebras*, Journal of Intelligent & Fuzzy Systems **29**(2) (2015), 517–524.
- [2] Y. S. Huang, *BCI-algebra*, Science Press, Beijing, 2006.
- [3] Y. B. Jun, K. J. Lee and S. Z. Song, *\mathcal{N} -ideals of BCK/BCI-algebras*, J. Chungcheong Math. Soc. **22** (2009), 417–437.
- [4] Y. B. Jun, X. L. Xin and E. H. Roh, *A class of algebras related to BCI-algebras and semigroups*, Soochow Journal of Mathematics **24**(4) (1998), 309–321.
- [5] K. J. Lee and Y. B. Jun, *The essence of subtraction algebras based on \mathcal{N} -structures*, Commun. Korean Math. Soc. **27**(1) (2012), 15–22.
- [6] J. Meng and Y. B. Jun, *BCK-algebras*, Kyungmoon Sa Co. Seoul, 1994.

¹DEPARTMENT OF MATHEMATICS,
SHAHID BEHESHTI UNIVERSITY, TEHRAN, IRAN
Email address: Bordbar.amirh@gmail.com

²DEPARTMENT OF MATHEMATICS,
GRADUATE UNIVERSITY OF ADVANCED TECHNOLOGY, MAHAN-KERMAN, IRAN
Email address: zahedi_mm@kgut.ac.ir

³DEPARTMENT OF MATHEMATICS EDUCATION,
GYEONGSANG NATIONAL UNIVERSITY, JINJU 52828, KOREA
Email address: skywine@gmail.com

*CORRESPONDING AUTHOR