

## IMPORTANT BIOMARKERS THAT PLAY A ROLE IN THE CHRONIC OBSTRUCTIVE PULMONARY DISEASE PROCESS

### VAŽNI BIOMARKERI OD ZNAČAJA ZA PROCES HRONIČNE OPSTRUKTIVNE BOLESTI PLUĆA

Emre Avci<sup>1</sup>, Gulcin Alp Avci<sup>2</sup>

<sup>1</sup>Faculty of Science and Arts, Department of Molecular Biology and Genetics, Biochemistry, Hitit University, Corum, Turkey

<sup>2</sup>Faculty of Science and Arts, Department of Molecular Biology and Genetics, Molecular Microbiology, Hitit University, Corum, Turkey

#### Summary

**Background:** Chronic obstructive pulmonary disease (COPD) that includes multiple mechanisms such as inflammation, infection, smoking, hypoxia, and lack of antioxidant response can cause oxidative stress. In our study, we aimed to determine the changes in some oxidative stress [malondialdehyde and glutathione] and some cellular immunity markers (neopterin and TGF- $\beta$ ) in patients diagnosed with COPD and determine the damage to the organism.

**Methods:** While the high-performance liquid chromatography (HPLC) method (Immuchrom kit, Germany) was utilized to determine MDA, GSH and NP levels, the ELISA method was used for TGF- $\beta$  levels.

**Results:** All obtained data regarding each parameter were compared with both COPD and healthy individuals and between parameters. There was a statistically significant difference between the control group of healthy subjects and COPD group in all parameters ( $p < 0.05$ ). A negative and correlation between oxidant MDA and antioxidant GSH parameters was determined ( $p = -0.394$ ).

**Conclusions:** As a result, it was seen that oxidative balance changed in the patient group and cellular immunity increased. When the obtained data and literature are taken into account, these changes occurring in oxidative balance and cellular immunity are of importance in determining the development in the pathogenesis of COPD, treatment options and their risks for heart disease in advance.

**Keywords:** chronic obstructive pulmonary disease, oxidative stress, cellular immunity

#### Kratak sadržaj

**Uvod:** Hronična opstruktivna bolest pluća (HOBP) koja uključuje više mehanizama kao što su inflamacija, infekcija, pušenje, hipoksija i odsustvo antioksidantnog odgovora može izazvati oksidativni stres. U našoj studiji, cilj je bio da se utvrde promene u nekim markerima oksidativnog stresa (malondialdehid i glutation) i ćelijskog imuniteta (neopterin i TGF- $\beta$ ) kod pacijenata sa dijagnozom HOBP i procene oštećenja organizma.

**Metode:** Dok je za određivanje nivoa MDA, GSH i NP korišćena metoda HPLC (Immuchrom kit, Nemačka), za nivoje TGF- $\beta$  upotrebljen je metod ELISA.

**Rezultati:** Svi dobijeni podaci vezani za svaki parametar upoređeni su kod obolelih od HOBP i zdravih subjekata kao i međusobno. Pronađena je statistički značajna razlika između kontrolne grupe zdravih osoba i grupe sa HOBP za sve parametre ( $p < 0,05$ ). Negativna i jaka korelacija utvrđena je između oksidantnog parametra MDA i antioksidantnog parametra GSH ( $p = -0,394$ ).

**Zaključak:** Kao rezultat, uočeno je da je u grupi obolelih izmenjena oksidativna ravnoteža i da je pojačan ćelijski imunitet. Kad se uzmu u obzir dobijeni podaci i literatura, ove promene koje nastaju u oksidativnoj ravnoteži i ćelijskom imunitetu od značaja su za određivanje razvoja u patogenezi HOBP, opcija lečenja i budućih rizika od srčanih bolesti.

**Ključne reči:** hronična opstruktivna bolest pluća, oksidativni stres, ćelijski imunitet

Address for correspondence:

Emre Avci  
Hitit University, Faculty of Science and Arts,  
Department of Molecular Biology and Genetics, Molecular  
Biology and Biochemistry, Hitit University, Corum, Turkey  
e-mail: avci.emre@yahoo.com; emreavci@hitit.edu.tr,  
Business Phone: +90364 227 7000/1647,  
Fax: +90 364 227 7005

## Introduction

Chronic obstructive pulmonary disease (COPD) is a multifactorial condition which is the third leading cause of mortality after the cardiovascular diseases and affects millions of people around the world (1–4). COPD has a chronic nature with genetic predisposition characterized by progressive and irreversible air-flow restriction occurring as a result of anti-oxidant and anti-protease systems impaired by smoking or several occupational-environmental factors (1, 5, 6). It produces a clinical course affecting severity and prognosis of the disease by features starting with the progressive airway obstruction and pulmonary involvement and then continuing with extra-pulmonary involvement. It is of importance, however, that it may be prevented by early treatment and it is a treatable condition (1, 6). Additionally, data from several studies show that one of the leading causes of death in COPD is ischemic heart disease (7, 8). Large-sized studies indicate that the COPD patients are at 2 to 3 times higher risk of cardiovascular mortality (8–10). Atherogenesis and the cardiovascular disease process are accelerated in patients with COPD. Key mechanisms in the pathogenesis of COPD include oxidative stress, decreased level of anti-oxidant protection, pulmonary and systemic inflammation, and endothelial dysfunction (4).

Oxidative stress has been defined as increased level of oxidants or decreased level of anti-oxidants. The oxidants are electron-acceptor molecules in the biological systems with the lungs being the most commonly involved (11). Several studies suggest that many mechanisms lead to oxidative stress. Especially, inflammation, several infections, hypoxia, exposure to tobacco smoke, decreased level of anti-oxidants may lead to oxidative stress (12, 13). Being an indirect marker that can detect the degree of oxidative damage in the systemic circulation and being a marker of oxidative stress, malondialdehyde (MDA) is formed as the result of oxidation of polyunsaturated fatty acids in the membrane structures impacted by free radicals (14).

Meanwhile, MDA is an end-product of lipid peroxidation and is used as a marker of tissue reaction chain rate and in detecting reactive oxygen species (ROS). Plasma concentration of MDA is due to oxidative destruction of the lipid-peroxides (15). Glutathione is present in the cells in oxidized (GSSG) and reduced (GSH) forms. Its anti-oxidant effect occurs during the cycle between these two forms. Glutathione is one of the anti-oxidant substances protecting the cells from the destructive effects of free oxygen species forming during metabolic processes. Oxidative stress in the organism may be increased by many factors including environmental, metabolic, and individual ones and thus the formation of free radicals may increase. Thus, the level of GSH protecting cells from the destructive effects of these free

radicals also varies. GSH level decreases and on the other hand the level of GSSG increases with increasing oxidative stress in the cell (16, 17).

Neopterin (NP) is one of the markers that may be used to detect activation of the cellular immune system (18). NP is an important marker for the coronary heart diseases and has been defined as a risk factor for acute respiratory infections in the patients with COPD (19). It is also a useful marker of oxidative stress, with its level responding to immunological stimulation rapidly. It may be detected more rapidly than erythrocyte sedimentation rate or soluble cytokine receptor concentration especially in the patients with inflammatory conditions (18). According to several studies, a strong correlation exists between inflammation and the extracellular matrix modulating cytokines (20–22). On the other hand, transforming growth factor-beta (TGF- $\beta$ ) plays an important role in re-inducing the airway in such chronic pulmonary diseases as COPD, asthma, and idiopathic pulmonary fibrosis (21). It is important for the airway remodeling. It may act as a regulatory key in many cellular and molecular processes (22, 23). Furthermore, TGF- $\beta$  is an important polypeptide regulated by autocrine intracellular processes (23). It also has important effects including stimulation of chemotaxis of the inflammatory cells in wound healing (24). Increased expression of TGF- $\beta$ 1 has been reported in some cardiovascular diseases. Thus, TGF- $\beta$ 1 expression may contribute to progression of some cardiac conditions (25).

The present study aimed to detect damage in the organism by determining the changes in some markers of oxidative stress, anti-oxidant levels, and cellular immunity in patients with the diagnosis of COPD and emphasize its effects on mortality and morbidity of cardiac conditions.

## Material and Methods

Our study includes patients diagnosed with COPD and being treated, with their stage of disease determined, in Corum Chest Diseases Hospital. According to the power analysis conducted before the study, 80 patients diagnosed with COPD and 30 healthy subjects (control group) were included in the study.

### *Storage of samples*

Serum samples were obtained from the related center under the supervision of the project manager, protected from light and stored in a freezer at -86 °C until the work was done.

### Determination of malondialdehyde (MDA) and glutathione (GSH) levels

Immuchrom kit in an HPLC system was utilized in the determination of MDA and GSH levels (HPLC Agilent Technologies 1200 series, USA; kit: Immuchrom GmbH, Germany) (Table I).

### Neopterin levels

Neopterin levels were determined using the HPLC method (High Performance Liquid Chromatography) used by Avci (2008) and Cakır (2010) (18, 26) (Table I).

### Transforming growth factor-beta (TGF- $\beta$ ) levels

In the determination of TGF- $\beta$  levels in serums, high-sensitivity TGF- $\beta$  ELISA (Bendermed System

Diagnostics, eBioscience, Austria, kit No. BMS249 / 4) kit was used (measurement wavelength: 450 nm, reference wavelength: 620 nm).

### Statistical analysis

Statistical analysis was performed using the SPSS 20.0 software (SPSS Inc., Chicago, IL). All measurements were taken in triplicates. The data were expressed as mean  $\pm$  SD. For assessing the normality of distribution and homogeneity of variances of the data, Shapiro-Wilk were used. According to Huck (27), the values of skewness and kurtosis must change between -1 and +1 in order for the data to show normal distribution. Accordingly, values obtained in this study have emerged that allow the condition of normality (Table II).

When all data were appropriate, parametric independent sample T test was used. To determine

**Table I** The working conditions and system parameters of the HPLC system used in the MDA, GSH and neopterin studies.

Working condition	System Parameters		
	MDA	GSH	Neopterin
HPLC Model	Agilent Technologies 1200 Series		
Sensor	Flouresans Sensor		
Mobile Phase	-	-	6.4 pH 0.015 mol/L phosphate buffer
Column material	Bischoff Prontosil Eurobond, 5 $\mu$ m; 125 mm $\times$ 4 mm	MZ Inertsil ODS, 5 $\mu$ m; 125 mm $\times$ 4 mm	Analytical Column 4.6 $\times$ 250 mm, Allsphere ODS- 2, C <sub>18</sub> 5 $\mu$ m reverse-phase Protector Column Spherisorb ODS-2, C <sub>18</sub> 5 mm reverse-phase
Wavelength	515 nm excitation and 553 nm emission	385 nm excitation and 515 nm emission	353 nm excitation and 438 nm emission
Flow Rate	0.8 to 1.0 mL/minute	0.75 to 1.0 mL/minute	0.8 mL/minute
Sample Volume	20 $\mu$ L		
Operation time	4 minute		20 minute
Temperature	30 $^{\circ}$ C		
Discrimination	-	-	Reverse phase ion exchange

**Table II** Skewness and kurtosis values of all parameters.

	Skewness	Skewness STD Error	R <sup>1</sup>	Kurtosis	Kurtosis STD Error	R <sup>2</sup>
MDA	0.753	0.23	0.523	-0.251	0.457	-0.708
GSH	0.379	0.23	0.149	-0.439	0.457	-0.896
NP	-0.956	0.23	-1.186	-0.944	0.457	-1.401
TGFB	-0.972	0.23	-1.202	-0.876	0.457	-1.333

R<sup>1</sup> = Skewness - Skewness STD Error,

R<sup>2</sup> = Kurtosis - Kurtosis STD Error, R<sup>2</sup> < | Kurtosis | = normal distribution

the association between MDA and GSH tested parameters, Pearson correlation coefficient was used. The critical significance level for the statistical tests performed was set to be 0.05.

### Ethics

This study was approved by the local ethical committee (Protocol 01-01/2014).

### Results

The serums of 80 COPD patients and of 30 healthy individuals in the control group were included in the study. The mean age for COPD patients enrolled in our study was  $54 \pm 7$ , while the mean age for the control group was detected to be  $48 \pm 4$ . The gender and age data of all patients are presented in Table III.

When, as markers of oxidative stress, oxidant MDA and antioxidant GSH data were evaluated, statistically significant changes were observed compared to the healthy control group. While MDA level in the control group was  $1.38 \pm 0.16$  mmol/L, it was deter-

mined to be  $1.93 \pm 0.48$  mmol/L in COPD patients. When it comes to the antioxidant indicator GSH, control group data determined to be  $394.77 \pm 30.65$  mmol/L was seen to decrease in COPD group to  $275.21 \pm 41.19$  mmol/L. Statistically significant difference between the groups was observed in both parameters ( $p < 0.05$ ) (Table IV, Figure 1). In addition, since MDA and GSH parameters of the data were parametric, negative and strong correlations between the two parameters were determined following Pearson correlation test ( $p = -0.394$ ).

In our study, when the immunity markers NP and TGF- $\beta$  data were studied, both indicators in COPD patients were detected to have increased compared to healthy subjects making up the control group (Table IV, Figure 2). In addition, statistically significant differences were found between the groups in both parameters ( $p < 0.05$ ).

### Discussion

Chronic obstructive pulmonary disease (COPD) is a chronic condition which develops with inflammation caused in response to harmful gases and particles and which forms progressive and irreversible air-flow limitation (2, 28). Since oxidative stress plays a central role in the pathogenesis of COPD, it must always be determined and in addition to smoking and childhood exposure to such harmful gas particles may also lead to the development of COPD (28, 29). Following the literature search results, the investigations of such parameters as antioxidant status, total oxidative status, various interleukins, C-reactive protein (CRP), tumor necrosis factor  $\alpha$  were generally given weight in COPD patients and the importance of


**Table III** Demographic data of groups, both healthy-control and patients with COPD.

	Control (n=30)	COPD (n=80)
Gender (Male/Female)	17/13	35-65
Age range	48/32	45-73


**Table IV** Values of oxidative stress markers (MDA and GSH) and immune markers (NP and TGF- $\beta$ ) of patients with COPD and control group and their 95% confidence intervals.

Parameters	Groups	n	<sup>a</sup> Mean $\pm$ SD	95 % CI
				Min-Max
MDA, $\mu\text{mol/L}^*$	Control	30	$1.38 \pm 0.16$	1.31-1.44
	COPD	80	$1.93 \pm 0.48$	1.82-2.03
GSH, $\mu\text{mol/L}^*$	Control	30	$394.77 \pm 30.65$	383.32-406.21
	COPD	80	$275.21 \pm 41.19$	266.04-284.37
NP, $\mu\text{mol/L}^*$	Control	30	$16.28 \pm 1.06$	15.88-16.68
	COPD	80	$38.15 \pm 1.86$	37.73-38.56
TGF-beta, pg/mL*	Control	30	$4.28 \pm 0.40$	4.13-4.43
	COPD	80	$8.74 \pm 0.41$	8.64-8.83

MDA: Malondialdehyde, GSH: Glutathione, NP: Neopterin; TGF-beta: Transforming growth factor-beta; CI: Confidence interval for mean; <sup>a</sup>Each of groups is mean value and standard deviation. \* $p < 0.005$ ; statistically significant differences were found between the groups.



**Figure 1** Oxidative stress markers (MDA and GSH) of patients with COPD and control group.



**Figure 2** Immune markers (NP and TGF-beta) of patients with COPD and control group.

better nutrition was emphasized (3, 19, 28, 30). In our study, investigating such parameters as oxidant (MDA), antioxidant (GSH) and cellular immunity markers (neopterin) and cytokine levels (TGF- $\beta$ ) of the patients who received COPD treatment, the data obtained were compared with a healthy control group and an assessment was made to decrease mortality and morbidity in terms of some diseases.

According to our study, MDA levels as a marker of lipid peroxidation in the COPD patients were determined to have increased compared to the healthy control group. Similarly, in studies conducted by many researchers related to oxidative stress, it was also seen that MDA level increased among patients compared to the control group (28, 31, 32). In our study, GSH, an important antioxidant and protector of

cells from the destructive effects of free oxygen radicals, was seen to have decreased in COPD patients compared with the healthy control group.

Different from our study, Drost et al. (33) found that GSH levels increased in the patient group compared to the control group. However, what was different in their study was that they studied the GSH parameter in bronchoalveolar lavage fluid. The results of studies reveal that the bronchoalveolar lavage technique is not suitable for diagnosis and is reported to have very low sensitivity for oxidative stress studies. In a study by Maury et al. (34, 35) in which they investigated the heterogeneity of systemic oxidative stress profiles in COPD patients, it was reported that GSH levels were quite low in COPD patients compared to the control group (36). Liu et al. (37) also reported in

a study in which they investigated the effect of cigarette smoke extracts on skeletal muscle cell aging by causing oxidative stress that they particularly observed a significant decrease in SOD and GSH activities.

When the formation mechanism of the important and precise marker of cellular immunity, neopterin, is examined, it is seen that the stimulated lymphocytes produce IFN-gamma, which activates macrophages and NP is synthesized from activated macrophages (38). NP both regulates the intracellular redox state and allows translocation of subunits of nuclear factor kB to the nucleus (39, 40). NP is regulated by a pro-inflammatory gene, so the clinical use of NP as an early inflammatory marker is important. Rising concentrations of NP are seen in patients with condensed monocyte/macrophage activity (41). Monocyte/macrophage increase is often encountered in people with heart diseases. Therefore, a relationship between NP and atherosclerosis was determined in the studies done.

Also, especially NP concentrations reflect the oxidative stress caused by the activation of the immune system (42–45). In the studies we have done so far, NP levels in COPD patients were found to increase compared with healthy individuals. Similarly, in a study by Rbyka et al. (46) in which they investigated the link between COPD and depression, to further investigate the inflammatory status of the patient, they assessed the cellular immune activation by measuring serum neopterin levels. As a result, they found a statistically significant increase of neopterin levels in patients with COPD compared to the control group. TGF- $\beta$  is one of the most important growth factors in the pathogenesis of COPD (47). When these studies are investigated, it can be seen that the inflammatory cytokine TGF- $\beta$  level is higher compared to healthy individuals (48, 49).

In a study by Zhang et al. (49) in which they identified several markers in patients with COPD, it was also seen that TGF- $\beta$  level increased significantly compared to the control group. In our study, TGF- $\beta$

levels of COPD patients were determined to have increased approximately two times as compared to healthy subjects. When the present studies were examined, we encountered quite a few studies that investigated serum neopterin and TGF- $\beta$  levels. It was also seen that there are not enough studies conducted to determine the role of oxidative stress markers. The data we obtained and literature search show that oxidant balance varies in this group of patients and that cellular immunity increases. When all the parameters are considered together, by referring to the burden on the development of COPD and the organism, we could determine that it created significant changes in several parameters.

In conclusion, it was also seen that the oxidant-antioxidant balance varies in COPD patients and that the levels of cellular immunity are directly or indirectly associated. It is also thought that the molecular changes in the levels of the oxidants and immunological agents caused by COPD might be risk factors for the development of cardiovascular disease in these patients. At the same time, our study is believed to shed light to the future studies to be conducted to determine both oxidative conditions and immunity in this patient group. Also, the effects of treatment protocols to be applied on oxidative stress antioxidants and on immune mediators are also believed to play a major role in the prognosis of the disease in these patients.

*Acknowledgment.* We would like to thank Dr. Mesut Arslan, MD, Corum Chest Diseases Hospital, Clinic of Chest Diseases, Corum, Turkey, for the clinical evaluation of the patients. This work was supported by the University of Hitit, Department of Scientific Research Projects through project no: FEF04.14.001

### **Conflict of interest statement**

The authors declare that they have no conflicts of interest for this work.

## References

- Decramer M, Janssens W, Miravittles M. Chronic obstructive pulmonary disease. *Lancet* 2012; 379: 1341–51.
- Bose S, Hansel NN, Tonorezos ES, Williams DL, Bilderback A, Breyse PN, Diette GB, McCormack MC. Indoor Particulate Matter Associated with Systemic Inflammation in COPD. *Journal of Environmental Protection* 2015; 6: 566–72.
- Puig-Vilanova E, Rodriguez DA, Lloreta J, Ausin P, Pascual-Guardia S, Broquetas J, et al. Oxidative stress, redox signaling pathways, and autophagy in cachectic muscles of male patients with advanced COPD and lung cancer. *Free Radical Biology and Medicine* 2015; 79: 91–108.
- Samorukova EI, Adasheva TV, Zadionchenko VS, Rosliakova GA. Pleiotropic Effects of Rosuvastatin in Chronic Obstructive Pulmonary Disease Patients With High Cardiovascular Risk. *Kardiologia* 2015; 55 (7): 32–7.
- Al-Azzawi AM, Ghoneim HA, Elmadbouh I. Evaluation of vitamin D, vitamin D binding protein gene polymorphism with oxidant – antioxidant profiles in chronic obstructive pulmonary disease. *J Med Biochem* 2017; 36: DOI: 10.1515/jomb-2017-0012.
- Yıldırım N, Demir T, Gemicioğlu B, Kıyan E, Oğuzülgen K, Polatlı M, et al. Acute exacerbation in COPD and asthma. *Tüberk Toraks* 2015; 63 (2): 111–31.
- Hansell AL, Walk JA, Soriano JB. What do chronic obstructive pulmonary disease patients die from? A multiple cause coding analysis. *Eur Respir J* 2003; 22: 809–14.
- Sin DD, Paul Man SF. Chronic Obstructive Pulmonary Disease as a Risk Factor for Cardiovascular Morbidity and Mortality. *Proc Am Thorac Soc* 2005; 2: 8–11.
- Jousilahti P, Vartiainen E, Tuomilehto J, Puska P. Symptoms of chronic bronchitis and the risk of coronary disease. *Lancet* 1996; 348: 567–72.
- Janssens JP, Herrmann F, MacGee W, Michel JP. Cause of death in older patients with anatomic-pathological evidence of chronic bronchitis or emphysema: a case-control study based on autopsy findings. *J Am Geriatr Soc* 2001; 49: 571–6.
- Umut S. KOAH patogenezinde oksidatif stress. In: Umut S, ed. KOAH seminer notları. 1997; İstanbul: Küre Basım.
- Yang W, Block ER. Effect of hypoxia and reoxygenation on the formation and release of reactive oxygen species by porcine pulmonary artery endothelial cells. *J Cell Physiol* 1995; 164: 414–23.
- Repine EJ, Bast A, Lankhorst I. Oxidative stress in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997; 156: 341–57.
- Koca T, Berber A, Koca HB, Demir TA, Koken T. Effects of hemodialysis period on levels of blood trace elements and oxidative stress. *Clin Exp Nephrol* 2010; 14 (5): 463–8.
- Repetto M, Jimena S, Alberto B. Lipid Peroxidation: Chemical Mechanism, Biological Implications and Analytical Determination. Ed. Angel Catala. *Biochemistry, Genetics and Molecular Biology*, Chapter 1, 2012; Intech.
- Meister A, Anderson ME. Glutathione. *Ann Rev Biochem* 1983; 52: 711–60.
- Kamiloğlu NN, Beytut E. Effects of Vitamin A and  $\beta$ -Carotene Injection on Levels of Vitamin E and on Glutathione Peroxidase Activity in Pregnant Tuj Sheep. *Turkish Journal of Veterinary & Animal Sciences* 2005; 29: 1033–8.
- Avci E, Coşkun Ş, Çakır E, Kurt Y, Özgür Akgül E, Bilgi C. Relations Between Concentrations of Asymmetric Dimethylarginine and Neopterin as Potential Risk Factors for Cardiovascular Diseases in Haemodialysis-Treated Patients. *Renal Failure* 2008; 30(8): 784–90.
- Garrod R, Marshall J, Barley E, Fredericks S, Hagan G. The relationship between inflammatory markers and disability in chronic obstructive pulmonary disease (COPD). *Prim Care Respir J* 2007; 16 (4): 236–40.
- Takizawa H, Tanaka M, Takami K, Ohtoshi T, Ito K, Satoh M, Okada Y, Yamasawa F, Nakahara K, Umeda A. Increased expression of transforming growth factor- $\beta$ 1 in small airway epithelium from tobacco smokers and patients with chronic obstructive pulmonary disease (COPD). *Am J Respir Crit Care Med* 2001; 163 (6): 1476–83.
- Fernandez IE, Eickelberg O. The impact of TGF- $\beta$  on lung fibrosis: from targeting to biomarkers. *Proc Am Thorac Soc* 2012; 9 (3): 111–16.
- Shahriary A, Seyedzadeh MH, Ahmadi A, Salimian J. The footprint of TGF- $\beta$  in airway remodeling of the mustard lung. *Inhal Toxicol* 2015; 27 (14): 745–53.
- Halwani R, Al-Muhsen S, Al-Jahdali H, Hamid Q. Role of transforming growth factor- $\beta$  in airway remodeling in asthma. *Am J Respir Cell Mol Biol* 2011; 44: 127–33.
- Karami A, Ghanei M, Alaeddini F, et al. Vascular endothelial growth factor in bronchoalveolar lavage fluid in sulfur mustard exposed lung patients. *Oman Med J* 2011; 26: 118–21.
- H Lim, YZ Zhu. Role of transforming growth factor- $\beta$  in the progression of heart failure. *Cell Mol Life Sci* 2006; 63: 2584–96.
- Cakir E, Akgul OE, Aydin I, Cayci T, Kurt YG, Onguru O, et al. The association between neopterin and acetaminophen-induced nephrotoxicity. *Renal Failure* 2010; 32 (6): 740–6.
- Huck, SW. *Reading statistics and research* (6th ed). Boston: Pearson, 2012.
- Dhokal N, Lamsal M, Baral N, Shrestha S, Dhokal SS, Bhatta N, Dubey RK. Oxidative stress and nutritional status in chronic obstructive pulmonary disease. *J Clin Diagn Res* 2015; 9 (2): BC01–4.
- Shaw JG, Vaughan A, Dent AG, O'Hare PE, Goh F, Bowman RV, Fong KM, Yang IA. Biomarkers of progression of chronic obstructive pulmonary disease (COPD). *J Thorac Dis* 2014; 6 (11): 1532–47.

30. Hsieh Meng-Jer, Yang Tsung-Ming, Tsai Ying-Huang. Nutritional supplementation in patients with chronic obstructive pulmonary disease. *Journal of the Fromosan Medical Association* 2016; 115 (8): 595–601.
31. Vibhuti A, Arif E, Deepak D, Singh B, Qadar-Pasha MA. Correlation of oxidative status with BMI and lung function in COPD. *Clin Biochem* 2007; 40: 958–63.
32. Bartoli ML, Novelli F, Costa F, Malagrino L, Melosini L, Bacci E, et al. Malondialdehyde in exhaled breath condensate as a marker of oxidative stress in different pulmonary diseases. *Mediators Inflamm* 2011; 2011; 891752.
33. Drost EM, Skwarski KM, Saulea J, Soler N, Roca J, Agusti A, et al. Oxidative stress and airway inflammation in severe exacerbations of COPD. *Thorax* 2005; 60(4): 293–300.
34. Domej W, Oettl K, Renner W. Oxidative stress and free radicals in COPD – implications and relevance for treatment. *International Journal of COPD* 2014; 9; 1207–24.
35. Turner RD, Bothamley GH. N-acetylcysteine for COPD: the evidence remains inconclusive. *Lancet Respir Med* 2014; 2(4): e3.
36. Maury J, Gouzi F, De Rigal P, Heraud N, Pincemail J, Molinari N, et al. Heterogeneity of Systemic Oxidative Stress Profiles in COPD: A Potential Role of Gender. *Oxid Med Cell Longev* 2015; 2015; 201843.
37. Liu W, Su W, Yang X, Bai J, Zhong X, He Z. Cigarette smoke extract induces senescence of murine skeletal muscle cells by oxidative stress-induced down-regulation of histone deacetylase. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi = Chinese Journal of Cellular and Molecular Immunology* 2015; 31(5): 630–3, 638.
38. Ursavaş A, Karadag M, Oral AY, Demirdogen E, Oral HB, Ege E. Association between serum neopterin, obesity and daytime sleepiness in patients with obstructive sleep apnea. *Respir Med* 2008; 102 (8): 1193–7.
39. Wirleitner B, Baier-Bitterlich G, Hoffmann G, et al. Neopterin derivatives to activate NF-kappa B. *Free Radic Biol Med* 1997; 23: 177–9.
40. Barnes PJ, Karin M. Nuclear factor-kappa B: a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med* 1997; 336: 1066–71.
41. Weiss G, Widner B, Zoller H, Schobersberger W, Fuchs D. Immune response and iron metabolism. *Br J Anaesth* 1998; 81 (Suppl. 1): 6–9.
42. Gupta S, Fredericks S, Schwartzman RA, Holt DW, Kaski JC. Serum neopterin in acute coronary syndromes. *Lancet* 1997; 349: 1252e3.
43. Schumacher M, Halwachs G, Tatzber F, et al. Increased neopterin in patients with chronic and acute coronary syndromes. *J Am Coll Cardiol* 1997; 30: 703–7.
44. Garcia-Moll X, Cole D, Zouridakis E, Kaski JC. Increased serum neopterin: a marker of coronary artery disease activity in women. *Heart* 2000; 83: 346–50.
45. Smith DA, Zouridakis EG, Mariani M, Fredericks S, Cole D, Kaski JC. Neopterin levels in patients with coronary artery disease are independent of Chlamydia pneumoniae seropositivity. *Am Heart J* 2003; 146: 69–74.
46. Rybka J, Korte SM, Czajkowska-Malinowska M, Wiese M, Kędziora-Kornatowska K, Kędziora J. The links between chronic obstructive pulmonary disease and comorbid depressive symptoms: role of IL-2 and IFN- $\gamma$ . *Clin Exp Med* 2015; 1–10.
47. Gorka K, Soja J, Jakiela B, Plutecka H, Gross-Sondej I, Cmiel A, et al. Relationship between the thickness of bronchial wall layers, emphysema score, and markers of remodeling in bronchoalveolar lavage fluid in patients with chronic obstructive pulmonary disease. *Pol Arch Med Wewn* 2016 Jun 30; 126 (6): 402–10.
48. Hodge SJ, Hodge GL, Reynolds PN, Scicchitano R, Holmes M. Increased production of TGF- $\beta$  and apoptosis of T lymphocytes isolated from peripheral blood in COPD. *Am J Physiol Lung Cell Mol Physiol* 2003; 285: L492–L499.
49. Zhang S, Xie JG, Su BT, Li JL, Hu N, Chen J, et al. MFG-E8, a clearance glycoprotein of apoptotic cells, as a new marker of disease severity in chronic obstructive pulmonary disease. *Braz J Med Biol Res* 2015; 48 (11): 1032–8.

Received: April 24, 2017

Accepted: May 31, 2017