

EFFECTS OF ZINC SUPPLEMENTATION ON SERUM COPPER TO ZINC AND CRP TO ALBUMIN RATIOS IN HEMODIALYSIS PATIENTS**EFEKTI SUPLEMENTACIJE CINKOM NA ODNOS SERUMSKOG BAKRA I CINKA I ODNOS CRP PREMA ALBUMINU KOD PACIJENATA NA HEMODIJALIZI**

Marwa Hajji^{1,2}, Rania Khedher^{1,3}, Mehdi Mrad^{1,4}, Mohamed Bassem Hammami^{1,2}, Nawel Rafrafi⁵, Salma Chouchi⁶, Moncef Feki^{1,2}, Afef Bahlous^{1,4}, Karim Zouaghi^{1,3}, Hayet Fellah^{1,2}

¹University of Tunis El Manar, Faculty of Medicine of Tunis, 1007 Tunis, Tunisia

²Rabta University Hospital, Laboratory of Biochemistry, LR99ES 11, 1007 Tunis, Tunisia

³Rabta University Hospital, Department of Nephrology, 1007 Tunis, Tunisia

⁴Pasteur Institute, Laboratory of Biochemistry, 1002 Tunis, Tunisia

⁵Radial Hemodialysis Clinic, 2010 Manouba, Tunisia

⁶Udial Hemodialysis Clinic, 1007 Tunis, Tunisia

Summary

Background: Zinc (Zn) deficiency is a common condition and could contribute to poor outcomes in hemodialysis (HD) patients. The aim of this study was to evaluate the effects of Zn supplementation on serum copper (Cu) to Zn and C-reactive protein (CRP) to albumin ratios (CAR) in HD patients.

Methods: Seventy-seven HD patients were enrolled in a multicentre simple-blind randomized clinical trial. Only 37 HD patients completed the study; they were randomly divided into two groups and supplemented with zinc sulphate (n=17) or placebo (n=20) for two months. Serum Zn and Cu were measured by atomic absorption spectrophotometry. Serum albumin and hypersensitive-CRP were assessed by colorimetric and immunoturbidimetric method, respectively. Determinations were performed before and after supplementation.

Results: After two months of supplementation, serum Zn significantly increased, and Cu to Zn ratio decreased in Zn supplemented group, but remained unchanged in the placebo group. In parallel, serum albumin concentrations significantly increased, and CAR decreased in Zn supplemented group only.

Kratak sadržaj

Uvod: Manjak cinka (Zn) je uobičajeno stanje i može doprineti lošim ishodima kod pacijenata na hemodijalizi (HD). Cilj ove studije je bio da proceni efekte suplementacije Zn na odnos serumskog bakra (Cu) i Zn i odnos C-reaktivnog proteina (CRP) i albumina (CAR) kod pacijenata na hemodijalizi.

Metode: Sedamdeset i sedam pacijenata na hemodijalizi je uključeno u jednostruko – slepo i randomizovano kliničko ispitivanje u više centara. Samo 37 pacijenata na hemodijalizi je završilo studiju. U nasumično podeljene dve grupe počelo se sa suplementacijom u ishrani cink sulfatom (n = 17) ili placebo (n = 20) tokom dva meseca. Serum Zn i Cu su mereni atomskom apsorpcionom spektrofotometrijom. Albuminski serum i hipersenzitivni CRP procenjeni su kolorimetrijskim i imunoturbidimetrijskim metodom, svaki posebno. Merenja su vršena i pre i posle suplementacije.

Rezultati: Nakon dva meseca suplementacije Zn u serumu se značajno povećao, a odnos Cu i Zn smanjio u grupi gde je sprovedena suplementacija Zn, ali je ostao nepromenjen u grupi sa placebo. Paralelno sa tim, koncentracije albumina u serumu značajno su porasle, a CAR se smanjio samo u grupi sa suplementacijom Zn.

Address for correspondence:

Marwa Hajji
Rabta Hospital, Laboratory of Biochemistry, 1007 Jebbari,
Tunis, Tunisia
Tel/Fax: 216 71 561 912; Mobile: 216 22 761880;
e-mail: marouahajji1@gmail.com

List of abbreviations: CAR, CRP to albumin ratio; CRP, C-reactive protein; HD, hemodialysis.

Conclusions: Zn supplementation reduces Cu to Zn and CRP to albumin ratios in HD patients. These changes point towards an improvement in nutritional, oxidative and inflammatory status. The study findings suggest that correcting Zn deficiency reduces poor outcomes in HD patients.

Keywords: copper to zinc ratio, C-reactive protein to albumin ratio, hemodialysis, zinc supplement

Introduction

HD is associated with a high burden of poor cardiovascular outcomes and increased mortality (1). This has been associated with conditions such as malnutrition, inflammation and oxidative stress (2, 3). Zn is a trace element that is essential for human health. It regulates numerous enzyme activities and immune function and exerts antioxidant and anti-inflammatory effects (4, 5). It is involved in proteins and lipids metabolism, nucleic acids synthesis and oxygen transportation (6, 7). Zn deficiency, a common condition in HD patients, is associated with malnutrition, inflammation and oxidative stress (6–9) and thus could be an additional contributor to poor outcomes in HD patients (10). Cu, another trace element, has a role in haemoglobin synthesis, immune function and defence against oxidants (11). It was shown that Cu and Zn interact antagonistically at high levels (12). Serum Cu to Zn ratio is considered a better marker for inflammation, oxidative stress and cardiovascular risk than serum Zn or Cu alone (3, 13). Elevated serum CRP and hypoalbuminemia are common in HD patients and were associated with adverse health effects in these patients (14–16). CAR is a biomarker of inflammation and nutrition. Elevated CAR reflects increased inflammation superimposed with malnutrition. It was shown to predict poor outcomes and increased mortality risk in patients with severe sepsis (17), cancer (18), acute coronary syndrome (19) as well as patients undergoing HD (20).

Zn supplementation was proved to ameliorate nutritional, inflammatory and oxidant status (9, 21, 22). A previous study showed a decrease in Cu to Zn ratio in patients under dialysis following Zn supplementation (9). However, the impact on CAR has never been investigated. The present study aimed to examine the effect of Zn supplementation on serum Cu to Zn and CRP to albumin ratios.

Material and Methods

A multicentre randomized single-blind clinical trial was conducted among 77 HD patients (45 men and 32 women) from February 2017 to March 2018. The patients were recruited from five centres: Rabta Hospital Nephrology department, Matri Hospital HD unit, Manouba HD Radial centre, Manouba HD public centre as well as HD Udial centre. An individual

Zaključak: Suplementacija Zn smanjuje odnos Cu u odnosu na Zn, i CRP u odnosu na albumin kod pacijenata na hemodijalizi. Ove promene ukazuju na poboljšanje statusa u vezi sa ishranom, oksidacijom i zapaljenjima. Nalazi studije sugerišu da smanjenje deficita Zn umanjuje šanse za loše ishode kod pacijenata na hemodijalizi.

Ključne reči: odnos bakra prema cinku, odnos C-reaktivnog proteina i albumina, hemodijaliza, suplement cinka

clinical card was established on which personal data were collected. The eligibility criteria for patients to be enrolled were over 18 years of age, HD treatment for at least six months, HD performed three times per week (each for 4 hours) through a polysulfone membrane against a dialysis liquid containing the following ions: Na^+ : 138 mmol/L; K^+ : 2 mmol/L; Ca^{++} : 1.5 mmol/L; Mg^{++} : 0.5 mmol/L; Cl^- : 109 mmol/L; CH_3CO_2^- : 3 mmol/L; HCO_3^- : 35 mmol/L. The clearance of urea evaluated by the ratio Kt/V was 1.2. The exclusion criteria were infection, gastrointestinal and liver diseases, congestive heart failure, cancer, psychiatric illness, pregnancy, use of immunosuppressants, corticoids, estrogens or contraceptives, as well as active smoking and alcoholism.

The patients were randomly divided into two groups; 43 HD patients received one oral capsule containing 220 mg of Zn sulphate (100 mg of elemental Zn) (Zn group), and 34 HD patients received one oral capsule of similar appearance containing 220 mg of maltodextrin. Capsules were taken daily after dinner during 60 consecutive days. Randomization was performed while stratifying on gender, 5-year age class and duration of HD. Predialysis blood samples were obtained from all patients after a fast of 12 hours at inclusion (day 1) and after 60 days of supplementation (day 61). Serum Zn and Cu concentrations were determined by atomic absorption spectrophotometry (Perkin Elmer, Waltham, USA). Serum hypersensitive-CRP concentrations were determined by the immunoturbidimetric method using the Architect CI8200 auto analyser (Abbott Diagnostics, Chicago, USA). Serum albumin was assessed by colorimetric method using Cobas Integra 400 auto-analyser (Roche Diagnostics, Meylan, France). The study protocol was approved by the Ethics committee of Rabta Hospital and all patients gave their informed and signed consent to participate in the study. A total of 37 HD patients (17 in the supplemented group and 20 in the placebo group) completed the trial (Figure 1).

Statistical analysis

Statistical analyses were performed using the software package SPSS 22.0 for windows (SPSS Inc, Chicago, USA). Depending on the distribution of variables, data were reported as mean \pm standard devia-




Figure 1 Flow chart for the process of hemodialysis patients selection.

tion (SD) or median (interquartile range). Between-group comparisons were made using independent t-test or Mann Whitney test, as appropriate. Within-group comparisons of variables before and after supplementation were performed using paired t-test or Wilcoxon rank test, as appropriate. A p-value <0.05 based on two-sided calculation was considered significant.

Results


Baseline clinical and biochemical characteristics, including serum Zn concentrations and prevalence of

hypo zincemia (serum Zn < 800 µg/L), were comparable in Zn and placebo groups (Table I). After two months of supplementation, serum Zn concentrations increased and serum Cu concentrations and Cu to Zn ratio decreased in the Zn group but remained unchanged in the placebo group. Zn supplementation also resulted in a significant increase in serum albumin and decrease in CAR. Figure 2 shows an individual variation in Cu to Zn and CRP to albumin ratios under supplementation in Zn and placebo groups.

Table 1 Baseline clinical and biochemical characteristics in zinc supplemented and placebo groups of hemodialysis patients.

	Zinc group (n=17)	Placebo group (n=20)	p value
Mean age, years	52.9±13.5	53.6±15.8	0.89
Male gender, %	70.6	55.0	0.33
Duration of	6.65±3.93	6.55±3.73	0.93
Body mass index, kg/m ²	25.2±4.34	26.4±5.74	0.53
Serum creatinine, μmol/L	967±264	896±361	0.50
Serum albumin, g/L	37.1±2.17	37.2±3.62	0.90
Serum CRP, mg/L	3.90 (14.9)	5.90 (13.95)	0.85
Serum CRP to albumin ratio	0.09 (0.41)	0.15 (0.35)	0.88
Serum zinc, μg/L	768±166	812±168	0.42
Zinc deficiency *, %	64.7	55.0	0.55
Serum copper, μg/L	1160±332	1170±281	0.91
Serum copper to zinc ratio	1.52±0.38	1.47±0.35	0.62

Values are expressed as mean ± SD or median (interquartile range); CRP, C-reactive protein; *, serum Zn <800 μg/L

**Figure 2** Changes in serum copper to zinc and C-reactive protein to albumin ratios in hemodialysis patients after two months of zinc supplementation.

Discussion

Our study showed that Zn deficiency is common in Tunisian HD patients with a prevalence of 59.5%. This result is consistent with the findings of previous studies which reported Zn deficiency in more than 50% of HD patients (23–25). Hypozincemia may result from Zn removal during HD treatment, reduced gastrointestinal absorption, low dietary intake and increased urinary excretion as well as protein restriction, increased expression of intracellular metallothioneins, and multi-infections (3, 21, 26).

Zn supplementation in these HD patients resulted in an increase in serum Zn and a decrease in Cu to Zn ratio, which is also consistent with literature data (9, 21). If the improvement in Zn status following Zn supplementation is expected and comprehensible, the decrease in serum Cu could be explained by competition between the two trace elements at the phases of intestinal absorption and cellular trafficking. Indeed, Zn and Cu share the same enterocyte membranes' transporters and intracellular trafficking proteins (11, 12). The competition may explain the decrease in Cu to Zn ratio following Zn supplementation. A high Cu to Zn ratio reflects oxidative stress, increased inflammation and immune dysfunction (12, 13, 27). Thus, its reduction in HD patients is considered beneficial, which supports the usefulness of Zn supplementation in these patients.

The study also showed an increase in serum albumin and a reduction in CAR following Zn supplementation. Zn and Cu are known to influence the function of immune cells and the secretion of cytokines (26). Zn exerts anti-inflammatory effects through downregulation of NF- κ B, leading to a reduction in proinflammatory cytokines (28). Zn deficiency was shown to induce a decrease in CD4: CD8 ratio and the number of B-cells (29). However, Zn supplementation was demonstrated to increase cells expressing CD19 (9). Cu excess is considered potentially proinflammatory by inducing oxidative stress (30, 31). CAR combines levels of CRP and albumin, a positive and a negative protein of acute phase inflammation, respectively. The reduction in the inflammatory response associated with Cu to Zn ratio

decrease may inhibit the synthesis of positive proteins of acute phase inflammation including CRP and stimulate the synthesis of negative proteins of acute phase inflammation such as albumin. Our results are consistent with the findings of previous studies which reported a reduction in serum CRP concentrations (9, 21) and an increase in serum albumin after Zn supplementation (9, 32). However, no previous study has investigated the impact on the ratio in HD patients.

The study has some limitations. First, the small sample size might have lowered the statistical power of the study. In fact, a number of patients in both Zn and placebo groups dropped out from the trial for lack of compliance with the study design or due to adverse health effects. HD patients were supplemented without taking into account their baseline serum Zn concentrations. This might explain some side effects that occurred in some patients. Further, trials with large sample size in Zn deficient patients are needed to confirm our results.

In conclusion, Zn supplementation in HD patients resulted in a decrease in Cu to Zn and CRP to albumin ratios. Such modifications point towards an improvement of nutritional, inflammatory and oxidative status. The study findings suggest that Zn supplementation is beneficial in HD patients and may contribute to reduce poor outcomes and improve the survival and quality of life in these patients. Therefore, patients on HD should be screened for Zn deficiency, and any deficiency should be adequately corrected.

Acknowledgements. The authors gratefully acknowledge the patients and administrative, nursing and technical staff for their valuable help. Special thanks to «Vital Laboratory» for providing Zn sulphate and placebo.

The study was supported by funds of Research Laboratory LR99ES11, Ministry of Higher Education and Scientific Research of Tunisia.

Conflict of interest statement

The authors declare that they have no conflicts of interest in this work.

References

1. Wright Alastair Hutchison J. Cardiovascular disease in patients with chronic kidney disease. *Vas Heal and Risk Manag* 2009; 5: 713–22.
2. Fung F, Sherrard DJ, Gillen DL, et al. Increased risk for cardiovascular mortality among malnourished end-stage renal disease patients. *Am J Kidney Dis* 2002; 40: 307–14.
3. Guo CH, Chen PC, Yeh MS, Hsiung DY, Wang CL. Cu/Zn ratios are associated with nutritional status, oxidative stress, inflammation, and immune abnormalities in patients on peritoneal dialysis. *Clin Biochem* 2011; 44: 275–80.
4. Sun JY, Jing MY, Wang JF, et al. Effect of zinc on biochemical parameters and changes in related gene expression assessed by cDNA microarrays in pituitary of growing rats. *Nutrition* 2011; 22: 187–96.
5. Prasad AS. Clinical, Immunological, anti-inflammatory and antioxidant roles of zinc. *Exp Gerontol* 2008; 43: 370–7.

6. Jern NA, VanBeber AD, Gorman MA, Weber CG, Liepa GU, Cochran CC. The effects of zinc supplementation on serum zinc concentration and protein catabolic rate in hemodialysis patients. *J Ren Nutr* 2000; 10: 148–53.
7. Ribeiro RC, Sales VS, Neves FAR, Draibe S, Brandao-Neto J. Effects of Zinc on Cell-Mediated Immunity in Chronic Hemodialysis Patients. *Biol Trace Elem Res* 2004; 98: 209–18.
8. Fellah H, Feki M, Soussi M, et al. Oxidative stress in end-stage renal disease: evidence and association with cardiovascular events in Tunisian patients. *Tunis Med* 2006; 84: 724–9.
9. Guo CH, Wang CL. Effects of zinc supplementation on plasma copper/zinc ratios, oxidative stress, and immunological status in hemodialysis patients. *Int J Med Sci* 2013; 10: 79.
10. Miao X, Wang Y, Sun J, et al. Zinc protects against diabetes-induced pathogenic changes in the aorta: Roles of metallothionein and nuclear factor (erythroid-derived 2)-like 2. *Cardiovasc Diabetol* 2013; 12: 54.
11. Maggini S, Wintergerst ES, Beveridge S, Hornig DH. Selected vitamins and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *Br J Nutr* 2007; 98: 29–35.
12. Sudha R, Ponsuganthi K, Jones R. Serum zinc and copper levels in maintenance haemodialysis patients and its relationship with depression and anxiety. *Global Journal of Medicine and Public Health* 2015; 4.
13. Karahan SC, Deger O, Orem A, Ucar F, Erem C, Alver A. The effects of impaired trace element status on polymorphonuclear leukocyte activation in the development of vascular complications in type 2 diabetes mellitus. *Clin Chem Lab Med* 2001; 39: 109–15.
14. Yeun JY, Levine RA, Mantadilok V, Kaysen GA. C-reactive protein predicts all-cause and cardiovascular mortality in hemodialysis patients. *Am J Kidney Dis* 2000; 35: 469–76.
15. Kawaguchi T, Tong L, Robinson BM, et al. C-reactive protein and mortality in hemodialysis patients: the dialysis outcomes and practice patterns study (DOPPS). *Nephron Clin Pract* 2011; 117: 167–78.
16. Poon PYK, Szeto CC, Kwan BCH, Chow KM, Li PKT. Relationship between CRP polymorphism and cardiovascular events in Chinese peritoneal dialysis. *Clin J Am Soc Nephrol* 2012; 7: 304–9.
17. Kim MH, Ahn JY, Song JE, et al. The C-reactive protein/albumin ratio as an independent predictor of mortality in patients with severe sepsis or septic shock treated with early goal-directed therapy. *PLoS One* 2015; 10: e0132109.
18. Wu M, Guo J, Guo L, Zuo Q. The C-reactive protein/albumin ratio predicts overall survival of patients with advanced pancreatic cancer. *Tumour Biol* 2016; 37: 12525–33.
19. Çağdaş M, Rencüzoğulları I, Karakoyun S. Assessment of relationship between C-reactive protein to albumin ratio and coronary artery disease severity in patients with acute coronary syndrome. *Angiology* 2017; 70: 361–8.
20. Jyh-Chang H, Ming-Yan J, Yi-Hua L, Charn-Ting W. Precedent fluctuation of serum hs-CRP to albumin ratios and mortality risk of clinically stable hemodialysis patients. *PLoS One* 2015; 10: e0120266.
21. Rashidi AA, Salehi M, Piroozmand A, Sagheb MM. Effects of zinc supplementation on serum zinc and C-Reactive Protein concentrations in hemodialysis patients. *J Ren Nutr* 2014; 19: 475–8.
22. Argani H, Mahdavi R, Ghorbani-Haghjo A, Razzaghi E, Nikniaz L, Ghaemmaghami SJ. Effects of zinc supplementation on serum zinc and leptin levels, BMI, and body composition in hemodialysis patients. *J Trace Elem Med Biol* 2014; 28: 35–8.
23. Erten Y, Kayatas M, Sezer S, et al. Zinc deficiency: prevalence and causes in hemodialysis patients and effect on cellular immune response. *Trans Proc* 2010; 30: 850–2.
24. Dashti-Khavidaki S, Khalili H, Vahedi SM, Lessan-Pezeshki M. Serum zinc concentrations in patients on maintenance hemodialysis and its relationship with anemia, parathyroid hormone concentrations and pruritus severity. *Saudi J Kidney Dis Transpl* 2010; 21: 641–5.
25. Shiota J, Tagawa H, Izumi N, Higashikawa S, Kasahara H. Effect of zinc supplementation on bone formation in hemodialysis patients with normal or low turnover bone. *Ren Fail* 2015; 37: 57–60.
26. Bozalioglu S, Ozkann Y, Turan M, Simsek B. Prevalence of zinc deficiency and immune response in short-term hemodialysis. *Trace Elem Med Biol* 2005; 8: 243–49.
27. Malavolta M, Giacconi R, Piacenza F, et al. Plasma copper/zinc ratio: an inflammatory/nutritional biomarker as predictor of all-cause mortality in elderly population. *Biogerontology* 2010; 11: 309–19.
28. De Martin R, Hoeth M, Hofer-Warbinek R, Schmid JA. The transcription factor NF- κ B and the regulation of vascular cell function. *Arterioscler Thromb Vasc Biol* 2000; 20: 83–8.
29. Haase H, Rink L. The immune system and the impact of zinc during aging. *Immun Ageing* 2009; 6: 9.
30. Wintergerst ES, Maggini S, Hornig DH. Contribution of selected vitamins and trace elements to immune function. *Ann Nutr Metab* 2007; 51: 301–23.
31. Guo CH, Wang CL, Chen PC, Yang TC. Linkage of some trace elements, peripheral blood lymphocytes, inflammation, and oxidative stress in ESRD patients undergoing either hemodialysis or peritoneal dialysis. *Periton Dialysis* 2011; *Int* 31: 583–59.
32. Ghaemmaghami J, Mahdavi R, Faramarzi E, Mohamadpour N, Argani H. Does zinc supplementation improve dietary intake, symptoms of eating problems, and serum zinc levels in hemodialysis patients? *Dial Transplant* 2010; 39: 530–3.

Received: May 25, 2020

Accepted: July 1, 2020