

ASSOCIATION BETWEEN GENETIC VARIANTS IN HSA-MIR-27A AND HSA-MIR-146A GENES AND MALE INFERTILITY

ASOCIJACIJA VARIJANTI U GENIMA HSA-MIR-27A I HSA-MIR-146A SA MUŠKIM STERILITETOM

Srećko Rajovski¹, Suzana Matijašević Joković², Nikoleta Milanović², Nemanja Radovanović², Miloš Brkušanin², Dušanka Savić-Pavićević², Zorana Dobrijević^{3*}, Goran Brajušković^{2*}

¹Private Healthcare Institution Clinical Hospital »Acibadem Sistina«, Skopje, North Macedonia

²University of Belgrade, Faculty of Biology, Centre for Human Molecular Genetics, Belgrade, Serbia

³University of Belgrade, Institute for the Application of Nuclear Energy - INEP, Department for Metabolism, Belgrade, Serbia

Summary

Background: miRNAs have enormous potential to be used as diagnostic and prognostic markers as well as therapeutic targets in male infertility and diseases of the reproductive system. This study aimed to investigate the association between the two functional genetic variants in the *hsa-miR-27a* (rs2910164) and *hsa-miR-146a* gene (rs895819) and male infertility in North Macedonian population, as well as to test their association with the values of major seminal parameters.

Methods: The case group included in this study comprised 158 men initially diagnosed with idiopathic male infertility. The control group included 126 age-matched healthy male volunteers who fathered at least one child.

Results: We report the association of rs2910164 minor allele C for the first time with the increased susceptibility to asthenoteratozoospermia. Additionally, our results indicating the association of allele C with low sperm vitality are a novel finding. We did not demonstrate the association between genetic variant rs895819 and the risk of different types of male infertility. Still, the number of participants with CC genotype in subjects diagnosed with asthenoteratozoospermia was null, while in controls, it reached 7.2%. We further detected the rs895819 genotype-dependent difference in rapid progressive sperm motility.

Kratak sadržaj

Uvod: miRNA molekule odlikuje izuzetan potencijal za upotrebu u vidu dijagnostičkih i prognostičkih biomarkera i terapijskih targeta kod muškog steriliteta i bolesti reproduktivnog sistema. Cilj ove studije je bilo ispitivanje asocijacije dveju funkcionalnih genetičkih varijanti gena *hsa-miR-27a* (rs2910164) i *hsa-miR-146a* (rs895819) i idiopatskog muškog steriliteta u populaciji Severne Makedonije i testiranje njihove asocijacije sa vrednostima seminalnih parametara.

Metode: Grupu ispitanika činilo je 158 muškaraca sa inicijalnom dijagnozom idiopatskog muškog steriliteta. Kontrolna grupa je uključivala 126 muškaraca volontera podudarne starosti koji su imali barem jedno dete.

Rezultati: Po prvi put je pokazana asocijacija manje učestalosti alela C varijante rs2910164 sa povišenim rizikom za pojavu steriliteta kod muškaraca sa astenoteratozoospermijom. Takođe, rezultati koji ukazuju na asocijaciju alela C sa nižom stopom vijabilnosti spermatozoida predstavljaju novi nalaz. Nije pokazana asocijacija varijante rs895819 sa rizikom za pojavu muškog steriliteta. Ipak, procenat učesnika u studiji sa genotipom CC među muškarcima sa dijagnozom astenoteratozoospermije bio je 0, dok je u grupi kontrola iznosio 7,2%. Pored toga, ustanovljen je efekat varijante rs895819 na brzu progresivnu pokretljivost spermatozoida.

Address for correspondence:

Zorana Dobrijević, University of Belgrade – Institute for the Application of Nuclear Energy, Department for Metabolism, Belgrade, Serbia

Phone: +381 11 2618 666

e-mail: zorana.dobrijevic@inep.co.rs

Goran Brajušković

University of Belgrade – Faculty of Biology, Centre for Human Molecular Genetics, Belgrade, Serbia

Phone: +381 11 2639 100

e-mail: brajuskovic@bio.bg.ac.rs

List of abbreviations: AIC, Akaike information criterion; AZF, azoospermia factor; CI, confidence interval; EDTA, ethylenediaminetetraacetic acid; GWAS, Genome-Wide Association Studies; HWE, Hardy-Weinberg equilibrium; IQR, interquartile range; KDM3A, lysine-specific demethylase 3A; miR-146a, *hsa-miR-146a-5p*; miR-27a, *Hsa-miR-27a-3p*; miRNAs, microRNAs; ncRNAs, non-coding RNAs; NOA, non-obstructive azoospermia; OR, odds ratio; PCR, polymerase chain reaction; PRM1, protamine 1; SSCs, spermatogonial stem cells; TNP1, transition protein 1; WHO, World Health Organization.

Conclusions: The association of rs2910164 and rs895819 with idiopathic male infertility in general is unlikely. However, both of these variants show an association with certain types of male infertility and with sperm abnormalities, which need to be confirmed in later studies in different ethnic groups.

Keywords: male infertility, rs2910164, rs895819, polymorphism, miR-27a, miR-146a

Introduction

Infertility is a disease of the male or female reproductive system defined by the failure to achieve a pregnancy after 12 months or more of regular unprotected sexual intercourse. Infertility impacts millions of people worldwide, often with devastating consequences (1).

MicroRNAs (miRNAs), an important class of non-coding RNAs (ncRNAs), are major regulators of gene expression and have been recognised as important regulators of various biological processes, including spermatogenesis. The presence of miRNAs in testis, epididymis, sperm cells, seminal plasma and extracellular vesicles (i.e., exosomes and microvesicles) and the known functions in the post-transcriptional regulation of gene expression associated with these molecules suggest that deregulation of their expression may lead to alterations in spermatogenesis and embryogenesis (2). At the same time, miRNAs have enormous potential to be used as diagnostic and prognostic markers as well as therapeutic targets in male infertility and diseases of the reproductive system (3).

Numerous specific miRNAs have been discovered in spermatogonial stem cells (SSCs), with roles ranging from initiating or blocking the differentiation process to modulating the self-renewal mechanism. Hsa-miR-146a-5p (miR-146a) directly binds and inhibits a coregulator of retinoid receptors, known as the mediator complex subunit 1. High levels of retinoic acid were found to inhibit the expression of miR-146a in undifferentiated spermatogonia, while upregulation of miR-146a was found to be sufficient to antagonise the effects of retinoic acid on spermatogonia (4). The genetic variant rs2910164 within miR-146a gene (GRCh38 5:160485411, C>G) is a functional polymorphism located in the »seed« region of the passenger strand and has been shown to generate a false C:U in the stem region, affecting the biogenesis of mature miR-146a (5). Therefore, this genetic variant could influence the risk of developing male infertility through its complex effect on the intertwined processes of miR-146a biogenesis, mature strand selections and the function of the resulting mature miRNA, all potentially related to retinoic acid signalling. However, despite its experimentally well-supported functional properties, rs2910164 has

Zaključak: Asocijacija rs2910164 i rs895819 sa idiopatskim muškim sterilitetom uopšteno nije potvrđena. S druge strane, za obe navedene varijante pokazana je asocijacija sa rizikom za pojavu određenih tipova muškog steriliteta, kao i sa abnormalnostima spermatozoida. Navedene rezultate je potrebno potvrditi u budućim studijama u drugim etničkim grupama.

Ključne reči: muški sterilitet, rs2910164, rs895819, polimorfizam, miR-27a, miR-146a

never been evaluated as a male infertility-related genetic variant in a Slavic population.

Hsa-miR-27a-3p (miR-27a) is a member of the miR-27 family, which has two members (miR-27a and miR-27b) and has been identified as an oncogenic miRNA that plays an important role in cancer development (6). This miRNA was predicted to target and regulate a specific H3K9 demethylase (lysine-specific demethylase 3A: KDM3A) with a proven role in male infertility in animal models. Disruption of epigenetic modification by KDM3A leads to a lack of spermatid chromatin after meiosis. Without the function of KDM3A in round spermatids, excessive methylation would lead to transcriptional suppression of transition protein 1 (TNP1) and protamine 1 (PRM1) genes and the occurrence of infertility in animal models. During the evolution of germline cells, epigenetic modifications via ncRNAs are crucial for regulating gene expression, genome stability and genome imprinting. Several studies have been conducted on human abnormalities and their association with miRNAs, including miR-27a. Norioun et al. (7) have shown that miR-27a is overexpressed in men with non-obstructive azoospermia. The genetic variant rs895819 (GRCh38 19:13836478, T>C) is located within a gene cluster encoding a single pri-miRNA for miR-23a, miR-24a and miR-27a. It was identified as an SNP that affects the sequence of the pre-miRNA terminal loop of miR-27a, leading to a change in its secondary structure and affecting the processing of miR-27a precursors (8). Similarly, as for rs2910164, hypotheses on the potential involvement of rs895819 in the molecular pathogenesis of male infertility rely on its functional role in the biogenesis of mature miRNA and on the downstream effect of the disrupted miR-27a function on its target mRNAs indispensable for maintaining a normal spermatogenesis process.

Since current knowledge about the involvement of miRNA-related variants in the genetic basis of idiopathic male infertility is limited, the aim of this study was to investigate the association between the two functional genetic variants in the miR-27a (rs895819) and miR-146a genes (rs2910164) and male infertility in the North Macedonian population. Furthermore, we aimed at testing the association of these genetic variants with the values of major semi-

nal parameters, which is even less investigated to date, and these results could provide valuable insight into the involvement of rs895819 and rs2910164, as well as miR-27a and miR-146a, in various aspects of male reproductive disorders.

Materials and Methods

The case group included in this study comprised 158 men initially diagnosed with idiopathic male infertility in 2018 at Acibadem Sistina Hospital, Skopje, North Macedonia. This research was conducted with the approval of the Ethics Committee of this medical institution (08-5110/01, May 2nd, 2018, Acibadem Sistina Hospital, Skopje, North Macedonia) and following the Helsinki Declaration of 1975. Written informed consent was obtained from all participants before their inclusion in the study. The control group included 126 healthy male volunteers from the North Macedonian population age-matched with the patients' group who fathered at least one child.

Exclusion criteria for the case group were diagnosis of orchitis and obstruction of vas deferens. In all patients, cytogenetic analysis excluded chromosome abnormalities as a possible cause of infertility. Multiplex polymerase chain reaction (PCR) based screening detected no microdeletions of the azoospermia factor (AZF) region within the Y chromosome in the selected group. Semen analysis in infertile men was conducted according to World Health Organization (WHO) guidelines, and patients were stratified into subgroups based on the abnormalities in semen concentration, sperm motility or morphology. Patients' basic data on age, serum

testosterone levels and the results of semen analysis are described in detail in our previous publication (9) and are given in *Table 1*.

Peripheral blood samples from patients and healthy controls were collected into commercially available 3 ml tubes with ethylenediaminetetraacetic acid (EDTA). Genomic DNA was isolated from peripheral blood samples using magnetic bead technology following the manufacturer's protocol. Genotyping was performed by using TaqMan™ SNP Genotyping Assays (C__3056952_20 and C__15946974_10) (Applied Biosystems, Waltham, MA, USA), according to the manufacturer's protocol. Real-time PCR amplification was carried out by Step One Real-time PCR System (Applied Biosystems, Waltham, MA, USA).

The exact test implemented in SNPStats software (Institut Català d'Oncologia, Barcelona, Spain) was used to assess the potential deviations of genotype distributions from Hardy-Weinberg equilibrium (HWE). The same software was used to test the associations of rs2910164 and rs895819 with the risk of idiopathic male infertility and with sperm viability assessed according to the threshold of 60% recommended by WHO. Odds ratios (ORs) with the corresponding 95% confidence intervals (CIs) were used as effect size measures in association tests, while *P* values <0.05 were considered to be statistically significant. The best-fitting association model among the five tested (log-additive, codominant, dominant, recessive and overdominant), was determined using the Akaike information criterion (AIC). By employing AIC as an estimator of the relative quality of statistical models, the best-fitting model was selected among the alternative models according to the lowest corresponding value of AIC. For quantitative results of

Table 1 Clinical characteristics of study participants diagnosed with idiopathic male infertility. Results are presented as mean±SD. NOA – Nonobstructive azoospermia.

Clinical characteristic	NOA	Other types of infertility
N	14	144
Age (years)	35.57±5.91	39.79±8.30
Sperm parameters		
Ejaculate volume (mL)	4.06±2.04	3.02±1.55
Total sperm number (10 ⁶)	-	225.15±170.32
Sperm density (10 ⁶ /mL)	-	75.34±52.16
Progressive motility (%)	-	33.57±13.87
Rapid progressive motility (%)	-	17.78±8.30
Average progression grade (0–4)	-	2.59±0.76
Vitality (%)	-	47.44±17.59

Table II Association of genetic variants rs895819 and rs2910164 with male infertility. Abbreviations: OR – odds ratio; CI – confidence interval; AIC – Akaike information criteria.

SNP	Genetic model	No of controls (%)	No of cases (%)	cases vs controls		
				OR (95% CI) ^a	P value ^a	AIC
rs895819						
	Codominant					
	TT	68 (54.4)	76 (48.4)	1.00	0.58	393
	TC	48 (38.4)	68 (43.3)	1.29 (0.79–2.12)		
	CC	9 (7.2)	13 (8.3)	1.26 (0.51–3.14)		
	Dominant					
	TT	68 (54.4)	76 (48.4)	1.00	0.3	391
	TC+CC	57 (45.6)	81 (51.6)	1.29 (0.80–2.06)		
	Recessive					
	TT+TC	116 (92.8)	144 (91.7)	1.00	0.79	392
	CC	9 (7.2)	13 (8.3)	1.13 (0.46–2.74)		
	Overdominant					
	TT+CC	77 (61.6)	89 (56.7)	1.00	0.36	391.2
	TC	48 (38.4)	68 (43.3)	1.25 (0.77–2.03)		
	Log-additive					
	-	-	-	1.20 (0.82–1.74)	0.35	391.2
rs2910164						
	Codominant					
	GG	69 (55.2)	84 (53.5)	1.00	0.58	393
	GC	47 (37.6)	67 (42.7)	1.29 (0.79–2.12)		
	CC	9 (7.2)	6 (3.8)	1.26 (0.51–3.14)		
	Dominant					
	GG	69 (55.2)	84 (53.5)	1.00	0.3	391
	GC+CC	56 (44.8)	73 (46.5)	1.29 (0.80–2.06)		
	Recessive					
	GG+GC	116 (92.8)	151 (96.2)	1.00	0.79	392
	CC	9 (7.2)	6 (3.8)	1.13 (0.46–2.74)		
	Overdominant					
	GG+CC	78 (62.4)	90 (57.3)	1.00	0.36	391.2
	GC	47 (37.6)	67 (42.7)	1.25 (0.77–2.03)		
	Log-additive					
	-	-	-	1.20 (0.82–1.74)	0.35	391.2

^aAdjusted for age.

semen analysis, the Kolmogorov-Smirnov test was used to assess the normality of the distribution of results, while the F-test was employed to analyse the equality of variances in study groups. Two-tailed Student's T-test was used to compare normally distributed results, while in cases of significant deviations from the normal distribution, the Mann-Whitney U test was applied.

Results

Table I summarises the basic data and semen parameters of patients diagnosed with male infertility. Most of the infertile men engaged in the present study were diagnosed with infertility types other than NOA (91%), including oligozoospermia, oligoasthenozoospermia, oligoteratozoospermia, oligoasthenoteratozoospermia, asthenoteratozoospermia

Table III Comparison of genotype distributions of rs895819 and rs2910164 between subjects with asthenoteratozoospermia and fertile controls. Abbreviations: OR – odds ratio; CI – confidence interval; AIC – Akaike information criteria.

SNP	Genetic model	No of controls (%)	No of cases (%)	cases vs controls		
				OR (95% CI) ^a	P value ^a	AIC
rs895819	Codominant					
	TT	68 (54.4)	10 (45.5)	1.00	0.11	127.7
	TC	48 (38.4)	12 (54.5)	1.72 (0.68–4.32)		
	CC	9 (7.2)	0 (0)	0.00 (0.00–NA)		
	Dominant					
	TT	68 (54.4)	10 (45.5)	1.00	0.44	129.5
	TC+CC	57 (45.6)	12 (54.5)	1.43 (0.58–3.56)		
	Recessive					
	TT+TC	116 (92.8)	22 (100)	1.00	0.082 ^b	127.1
	CC	9 (7.2)	0 (0)	0.00 (0.00–NA)		
	Overdominant					
	TT+CC	77 (61.6)	10 (45.5)	1.00	0.16	128.1
	TC	48 (38.4)	12 (54.5)	1.94 (0.77–4.87)		
	Log-additive					
-	-	-	-	1.05 (0.50–2.19)	0.9	130.1
rs2910164	Codominant					
	GG	69 (55.2)	6 (27.3)	1.00	0.025	124.7
	GC	47 (37.6)	15 (68.2)	3.77 (1.35–10.51)*		
	CC	9 (7.2)	1 (4.5)	1.27 (0.14–11.80)		
	Dominant					
	GG	69 (55.2)	6 (27.3)	1.00	0.013	123.9
	GC+CC	56 (44.8)	16 (72.7)	3.35 (1.22–9.16)		
	Recessive					
	GG+GC	116 (92.8)	21 (95.5)	1.00	0.63	129.8
	CC	9 (7.2)	1 (4.5)	0.61 (0.07–5.08)		
	Overdominant					
	GG+CC	78 (62.4)	7 (31.8)	1.00	0.0068	122.7
	GC	47 (37.6)	15 (68.2)	3.66 (1.38–9.71)		
	Log-additive					
-	-	-	-	1.87 (0.93–3.78)	0.082 ^b	127.1

^a adjusted for age; ^b statistical trend of significance; * Statistically significant results are shown in bold.

Table IV Comparison of genotype distributions of rs895819 and rs2910164 between infertile subjects with sperm vitality of at least 60% and those with a lower percentage of live spermatozoa. Abbreviations: OR – odds ratio; CI – confidence interval; AIC – Akaike information criteria.

SNP	Genetic model	Vit. 60% (%)	Vit. <60% (%)	Vit. <60% vs Vit. 60%		
				OR (95% CI) ^a	P value ^a	AIC
rs895819						
	Codominant					
	TT	16 (47.1)	47 (46.1)	1.00	0.63	160
	TC	14 (41.2)	48 (47.1)	1.17 (0.51–2.65)		
	CC	4 (11.8)	7 (6.9)	0.59 (0.15–2.29)		
	Dominant					
	TT	16 (47.1)	47 (46.1)	1.00	0.93	158.9
	TC+CC	18 (52.9)	55 (53.9)	1.04 (0.48–2.26)		
	Recessive					
	TT+TC	30 (88.2)	95 (93.1)	1.00	0.37	158.1
	CC	4 (11.8)	7 (6.9)	0.55 (0.15–2.00)		
	Overdominant					
	TT+CC	20 (58.8)	54 (52.9)	1.00	0.55	158.5
	TC	14 (41.2)	48 (47.1)	1.27 (0.58–2.79)		
Log-additive						
-	-	-	-	0.90 (0.49–1.67)	0.75	158.8
rs2910164						
	Codominant					
	GG	24 (70.6)	50 (48.5)	1.00	0.066 ^b	156
	GC	9 (26.5)	50 (48.5)	2.69 (1.13–6.37)*		
	CC	1 (2.9)	3 (2.9)	1.39 (0.14–14.20)		
	Dominant					
	GG	24 (70.6)	50 (48.5)	1.00	0.023	154.3
	GC+CC	10 (29.4)	53 (51.5)	2.56 (1.11–5.88)		
	Recessive					
	GG+GC	33 (97.1)	100 (97.1)	1.00	0.98	159.5
	CC	1 (2.9)	3 (2.9)	0.97 (0.10–9.69)		
	Overdominant					
	GG+CC	25 (73.5)	53 (51.5)	1.00	0.02	154.1
	GC	9 (26.5)	50 (48.5)	2.65 (1.12–6.24)		
Log-additive						
-	-	-	-	2.19 (1.01–4.75)	0.039	155.2

^a adjusted for age

^b statistical trend of significance

* Statistically significant results are shown in bold.

and teratozoospermia. The values of semen parameters demonstrated high dispersion concerning the mean due to variable results associated with different diagnoses and characteristics of different forms of infertility. The average age of healthy fertile controls

(40.85±10.58, not shown in Table I) at sampling was almost matched to the age of infertile individuals.

Genotyping of rs895819 and rs2910164 was successful for 99.4% (157/158) of study participants




Figure 1 Sperm progressive motility (a) and rapid progressive motility (b) in infertile men across genotypes of rs95819. Data represent the median and interquartile range (IQR). $P < 0.05$ is indicated by an asterisk.

diagnosed with idiopathic infertility and 98.4% (125/127) of fertile controls. Genotype distributions, summarised in *Table II*, were not found to significantly deviate from HWE in control subjects ($P = 1$ and $P = 0.82$ for rs95819 and rs2910164, respectively).

According to the results of genetic association tests presented in *Table II*, neither rs95819 nor rs2910164 demonstrated an association with the risk of idiopathic infertility in our North Macedonian male cohort. The results for all genetic models tested remained statistically insignificant for comparing genotype distributions between infertile men and age-matched fertile controls (*Table II*).

When we stratified infertile subjects into groups according to the results of semen analysis in terms of abnormalities in semen parameters and the related diagnosis, infertile men with asthenoteratozoospermia showed a higher frequency of rs2910164 heterozygous genotype compared to controls (*Table III*). According to AIC score, the over-dominant genetic model was selected as the best-fitting model of association between rs2910164 and asthenoteratozoospermia, while the corresponding OR value for GC vs. homozygous genotypes was 3.66 (95%CI 1.38–9.71, $P = 0.0068$) (*Table III*), indicative of an increased risk of asthenoteratozoospermia associated with GC genotype. Additionally, statistically significant differences in genotype distributions between infertile men diagnosed with asthenoteratozoospermia and fertile controls was demonstrated for dominant and codominant genetic models ($P = 0.013$ and $P = 0.025$, respectively), with an increased risk of asthenoteratozoospermia associated with GC genotype or with a combination of genotypes which

includes GC. For rs95819, only a statistical trend of significance ($0.05 > P > 0.1$) was reached for a recessive genetic model of association with asthenoteratozoospermia. However, none of the subjects with asthenoteratozoospermia had rs95819 genotype CC, while the frequency of this genotype in the control group was 7.2% (*Table III*).

Both analysed variants showed no statistically significant differences in genotype distributions between subjects with other types of idiopathic male infertility and fertile controls (results not shown). We considered it statistically unjustified to compare genotype frequencies between patients diagnosed with non-obstructive azoospermia (NOA) and the control group since the number of participants in this group of infertile men was just 14 in our study cohort (*Table I*).

Non-azoospermic infertile patients with rs2910164 GC genotype presented with lower sperm viability, as defined by the threshold of 60% viable sperm (*Table IV*). Similarly, as for association with asthenoteratozoospermia, the over-dominant genetic model was selected as the best-fitting model of association between rs2910164 and low sperm viability and the calculated OR for GC vs. homozygous genotypes was 2.65 (95%CI 1.12–6.24, $P = 0.02$) (*Table IV*), suggesting that GC genotype associates with lower sperm viability. Although the AIC score was the lowest for the over-dominant model, statistical significance was reached for the other two genetic association models – dominant and log-additive ($P = 0.023$ and $P = 0.039$, respectively). For all the tested genetic models for which the statistical significance was reached, as well as for the comparison of GC vs GG

under the codominant model, the obtained ORs for GC genotypes and the genotype combinations that incorporate heterozygotes were over 2, indicating an increase in risk for lower sperm viability associated with tested compared to referent genotypes.

Even though rs895819 showed no evidence of association with male infertility diagnosis, as well as with progressive motility score (Figure 1A), carriers of TC genotype displayed a lower percentage of sperm with rapid progressive motility compared to infertile subjects with the most common TT genotype ($P=0.018$) (Figure 1B).

Discussion

Male infertility, which is a major contributing factor in around 40–50% of cases of a couple's inability to conceive, is a multifactorial disorder with a recognised causative role of genetic factors involved in its complex molecular basis (10, 11). Besides chromosome aberrations and single gene mutations with high penetrability, which account for a relatively small percentage of infertile cases, genetic variants with modest or low penetrability were identified as contributors to genetic susceptibility to develop male infertility (11). These risk-associated genetic variants were discovered through large genome-wide association studies (GWAS), high-throughput techniques, and candidate gene-based approaches (11). For such genetic association studies focused on single genes or groups of functionally related genes, candidates were selected based on their involvement in the molecular basis of spermatogenesis, sperm functionality, oxidative stress response and other major processes related to infertility. In the recent decade, miRNA molecules emerged as important regulators of the spermatogenesis process, while their dysregulation was associated with a reduced sperm count, lower sperm viability and/or motility, as well as abnormalities in sperm morphology (4, 12). Therefore, miRNA-related genes seem plausible candidates for genetic association studies on male infertility.

MiRNA variants commonly selected for case-control studies are predicted or experimentally confirmed as functional polymorphisms that affect multiple aspects of miRNA-based mechanisms. This includes miRNA biogenesis, their localisation, sequence of the mature miRNA molecules, strand selection, efficiency of post-transcriptional regulation of targeted genes, and the selection of mRNA targets (13). Given that these genetic variants could simultaneously affect multiple processes regulated by the same miRNA, most of them were evaluated in the context of various human pathological conditions, including malignant and immunological diseases, metabolic disorders, and other common health issues (14–18). Still, data on the effect of commonly evaluated genetic variants on the susceptibility to male

infertility are scarce or even lacking for certain aspects or types of infertility-related abnormalities.

We selected genetic variants within genes encoding miR-146a and miR-27a for our present study. Both of these miRNA molecules are expressed in germ cells and were previously found to have an altered expression level in biological samples obtained from infertile men compared to healthy fertile controls (4, 7, 19). Additionally, their role in regulating key cellular processes disrupted in disorders of sperm maturation was revealed, while several studies confirmed their involvement in oxidative stress response and/or inflammation (2, 4, 7, 20–22). It is worth noting that miR-27a is an androgen-regulated miRNA that stimulates androgen signalling via a positive feedback loop (23). We aimed to evaluate the effects of variants with functional properties, so we selected an extensively analysed genetic variant rs2910164 within the *hsa-miR-146a* gene. This genetic variant exhibits its functional effect on miR-146a function by altering the structure of the pre-miRNA hairpin, which affects the efficiency of mature miRNA biogenesis. Furthermore, rs2910164 could potentially affect the leader strand selection and change the sequence of the seed region of miR-146a-3p, which has consequences on the selection of the target gene pool (24). The second selected genetic variant is located within a host gene of a cluster of miRNAs miR-23a, miR-24-2 and miR-27a. This genetic variant alters the sequence of the terminal loop of pre-miR-27a and potentially affects the biogenesis of mature miR-27a by influencing the processing of primary miRNA by the Drosha enzyme (25, 26).

To date, rs2910164 was evaluated as a susceptibility variant in male infertility in two studies conducted in the Turkish population (27) and in Han Chinese (28). No evidence of such genetic association was found in both of these studies. However, a study by Tunçdemir et al. (27) evaluated the association of rs2910164 with idiopathic oligospermia without considering other types of infertility. Furthermore, their publication did not include results on the potential association between rs2910164 genotypes and the values of semen parameters, which we intended to evaluate in the present study. Lu et al. (28), on the other hand, evaluated their large study case-control study on the association of rs2910164 with idiopathic male infertility in general and with specific types of infertility diagnosed according to the results of semen analysis. Still, their criteria for patient selection differed from ours since their case group included men with normospermia, who, in fact, represented the largest percentage of their infertile subjects (28).

Our study was the first one to analyse the effects of rs2910164 on a Slavic population with a European Caucasian genetic background. This genetic variant shows a significant global variation in allele frequen-

cies. For this reason, we expected that our novel findings could differ from the previous and would be valuable for a more precise evaluation of the effects of rs2910164 in the context of the potential influence of ethnicity as a confounder. Regarding the results obtained for the association between this genetic variant and overall idiopathic male infertility, our findings match the previously published reports from Turkish and Han Chinese populations since our results did not support the supposed association (27, 28). However, these two studies did not evaluate the association of rs2910164 with asthenoteratozoospermia and provided no results on the potential effects of this genetic variant on abnormal sperm morphology or the combination of sperm motility and abnormal phenotype (27, 28). Therefore, our results suggest that the association of rs2910164 minor allele C with increased susceptibility to asthenoteratozoospermia could not be compared with any previous findings. Additionally, our results indicating the association of allele C with low sperm vitality are a novel finding (Table IV).

When it comes to rs895819 in a gene encoding miR-27a, a single previous association study related to male infertility was conducted in a Serbian cohort of infertile men (29). This previous case-control evaluation in a genetically close related population to the present one reported a lack of statistically significant association between rs895819 and idiopathic male infertility risk, consistent with the study's finding in the North Macedonian cohort. However, the number of infertile men and fertile controls in this previous study was relatively low, which warranted additional evaluation of the effects of rs895819 (29). Our present study did not demonstrate the association between this genetic variant and the risk of different types of male infertility. Still, the number of participants with CC genotype in a small group of subjects diagnosed with asthenoteratozoospermia was null, while in con-

trols, it reached 7.2%. Therefore, it is possible that the association of rs895819 under a recessive or allelic genetic model would be detected in a larger validation study. Our novel finding of the genotype-dependent difference in rapid progressive sperm motility in infertile men should also be evaluated in future more extensive studies. However, our finding is consistent with a previous report on the effect of miR-27a expression on sperm progressive motility (20).

A relatively small number of participants in both case and control groups is the main limitation of this study. Therefore, in order to provide a more precise estimation and to make further conclusions about the potential association between the analysed genetic variants and male infertility, including the effect on sperm parameters, the study size needs to be enlarged. The association of rs2910164 and rs895819 with idiopathic male infertility is generally unlikely, considering the matching results between the present and previous analyses. However, both of these variants show an association with certain types of male infertility and with sperm abnormalities, which need to be confirmed in later studies in different ethnic groups.

Acknowledgements. The research was financially supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Agreement no. 451-03-47/2023-01/200042). We owe gratitude to the Diagnostic laboratories and Assisted Reproduction (IVF) Department of Clinical Hospital »Acibadem Sistina« employees for their assistance in collecting and preparing samples.

Conflict of interest statement

All the authors declare that they have no conflict of interest in this work.

References

1. Cox CM, Thoma ME, Tchangalova N, Mburu G, Bornstein MJ, Johnson CL, et al. Infertility prevalence and the methods of estimation from 1990 to 2021: a systematic review and meta-analysis. *Hum Reprod Open* 2022; 2022: hoac051.
2. Salas-Huetos A, James ER, Aston KI, Carrell DT, Jenkins TG, Yeste M. The role of miRNAs in male human reproduction: a systematic review. *Andrology* 2020; 8: 7–26.
3. Khawar MB, Mehmood R, Roohi N. MicroRNAs: Recent insights towards their role in male infertility and reproductive cancers. *Bosn J Basic Med Sci* 2019; 19: 31–42.
4. Barbu MG, Thompson DC, Suci N, Voinea SC, Cretoiu D, Predescu DV. The Roles of MicroRNAs in Male Infertility. *Int J Mol Sci* 2021; 22: 2910.
5. Nikolić ZZ, Savić Pavićević DLj, Vukotić VD, Tomović SM, Cerović SJ, Filipović N, et al. Association between genetic variant in hsa-miR-146a gene and prostate cancer progression: evidence from Serbian population. *Cancer Causes Control* 2014; 25: 1571–5.
6. Bi W, Li J, Xiong M, Pan B, Zhang Z, Nasifu L, et al. The diagnostic and prognostic role of miR-27a in cancer. *Pathol Res Pract* 2023; 247: 154544.
7. Norioun H, Motovali-Bashi M, Javadirad SM. Hsa-miR-27a-3p overexpression in men with non-obstructive azoospermia: A case-control study. *Int J Reprod Biomed* 2020; 18: 961–8.
8. Shi D, Li P, Ma L, Zhong D, Chu H, Yan F, et al. A genetic variant in pre-miR-27a is associated with a reduced renal

- cell cancer risk in a Chinese population. *PLoS One* 2012; 7: e46566.
9. Rajovski S, Vučić N, Karanović J, Matijašević S, Savić-Pavićević D, Dobrijević Z, et al. Association of PRMT6, PEX10 and SOX5 genetic variants with idiopathic male infertility: Evidence from North Macedonian population and an updated meta-analysis. *Genetika* 2023; 55: 355–72.
 10. Kumar N, Singh AK. Trends of male factor infertility, an important cause of infertility: A review of literature. *J Hum Reprod Sci* 2015; 8: 191–6.
 11. Krausz C, Escamilla AR, Chianese C. Genetics of male infertility: from research to clinic. *Reproduction*. 2015; 150: R159–74.
 12. Klees C, Alexandri C, Demeestere I, Lybaert P. The Role of microRNA in Spermatogenesis: Is There a Place for Fertility Preservation Innovation? *Int J Mol Sci* 2023; 25: 460.
 13. Ryan BM, Robles AI, Harris CC. Genetic variation in microRNA networks: the implications for cancer research. *Nat Rev Cancer* 2010; 10: 389–402.
 14. Nikolić ZZ, Savić Pavićević DL, Vučić NL, Romac SP, Brajušković GN. Association between a Genetic Variant in the hsa-miR-146a Gene and Cancer Risk: An Updated Meta-Analysis. *Public Health Genomics* 2015; 18: 283–98.
 15. Dai J, Chen Y, Gong Y, Gu D, Chen J. Association of microRNA-27a rs895819 polymorphism with the risk of cancer: An updated meta-analysis. *Gene* 2020; 728: 144185.
 16. Li G, Liu B, Jiang Q, Zhang J, Xin S, Xu K. The association of two common polymorphisms in miRNAs with diabetes mellitus: A meta-analysis. *Medicine (Baltimore)* 2019; 98: e17414.
 17. Wang X, Xing Y, Wang Y, Du Z, Zhang C, Gao J. Association of microRNA gene polymorphisms with recurrent spontaneous abortion: An updated meta analysis. *Exp Ther Med* 2023; 25: 179.
 18. Zhang J, Tan H, Cao Q, Su G, Yang P. Meta-Analysis of miRNA Variants Associated with Susceptibility to Autoimmune Disease. *Dis Markers* 2021; 2021: 9978460.
 19. Abu-Halima M, Galata V, Backes C, Keller A, Hammadeh M, Meese E. MicroRNA signature in spermatozoa and seminal plasma of proven fertile men and in testicular tissue of men with obstructive azoospermia. *Andrologia* 2020; 52: e13503.
 20. Zhou JH, Zhou QZ, Yang JK, Lyu XM, Bian J, Guo WB, et al. MicroRNA-27a-mediated repression of cysteine-rich secretory protein 2 translation in asthenoteratozoospermic patients. *Asian J Androl* 2017; 19: 591–5. Erratum in: *Asian J Androl* 2023; 25: 758.
 21. Robertson SA, Zhang B, Chan H, Sharkey DJ, Barry SC, Fullston T, et al. MicroRNA regulation of immune events at conception. *Mol Reprod Dev* 2017; 84: 914–25.
 22. Zhao Y, Dong D, Reece EA, Wang AR, Yang P. Oxidative stress-induced miR-27a targets the redox gene nuclear factor erythroid 2-related factor 2 in diabetic embryopathy. *Am J Obstet Gynecol* 2018; 218: 136.e1–136.e10.
 23. Fletcher CE, Dart DA, Sita-Lumsden A, Cheng H, Rennie PS, Bevan CL. Androgen-regulated processing of the oncomir miR-27a, which targets Prohibitin in prostate cancer. *Hum Mol Genet* 2012; 21: 3112–27.
 24. Xu B, Feng NH, Li PC, Tao J, Wu D, Zhang ZD, et al. A functional polymorphism in Pre-miR-146a gene is associated with prostate cancer risk and mature miR-146a expression in vivo. *Prostate* 2010; 70: 467–72.
 25. Sun Q, Gu H, Zeng Y, Xia Y, Wang Y, Jing Y, et al. Hsa-mir-27a genetic variant contributes to gastric cancer susceptibility through affecting miR-27a and target gene expression. *Cancer Sci* 2010; 101: 2241–7.
 26. Nikolić Z, Savić Pavićević D, Vučić N, Cidilko S, Filipović N, Cerović S, et al. Assessment of association between genetic variants in microRNA genes hsa-miR-499, hsa-miR-196a2 and hsa-miR-27a and prostate cancer risk in Serbian population. *Exp Mol Pathol* 2015; 99: 145–50.
 27. Tunçdemir M, Yenmiş G, Tombultürk K, Arkan H, Soydaş T, Burak Tek R, et al. NFKB1 rs28362491 and pre-miRNA-146a rs2910164 SNPs on E-Cadherin expression in case of idiopathic oligospermia: A case-control study. *Int J Reprod Biomed* 2018; 16: 247–54.
 28. Lu J, Gu H, Tang Q, Wu W, Yuan B, Guo D, et al. Common SNP in hsa-miR-196a-2 increases hsa-miR-196a-5p expression and predisposes to idiopathic male infertility in Chinese Han population. *Sci Rep* 2016; 6: 19825.
 29. Vučić N, Nikolić Z, Savić Pavićević D, Kanazir S, Vukotić V, Romac S, et al. Assessment of possible association between genetic variant rs895819 in miR-27a gene and infertility in males diagnosed with non-obstructive azoospermia (NOA) from Serbian population [abstract]. *EAU 10th South Eastern European Meeting (SEEM); 2014 Oct 24-26; Belgrade, Serbia. Eur Urol Suppl* 2014; 13: e1475.

Received: May 17, 2024

Accepted: July 06, 2024