

# Cognitive aspects of schizophrenia - a narrative review

## Kognitivni aspekti šizofrenije – narativni pregled

Jelena Djordjevic<sup>1,2</sup>, Slađana Arsic<sup>3</sup>, Dragan Pavlovic<sup>4</sup>, Aleksandra Pavlovic<sup>5</sup>

1-Clinic for Neurology and Psychiatry for Children and Youth, Belgrade, Serbia.

2-University of Kragujevac, Faculty of Medical Sciences, Kragujevac, Serbia

3-The Academy of Applied Preschool Teaching and Health Studies, Department of Medical Studies, Cuprija, Serbia

4-University of Belgrade, Belgrade, Serbia

5-University of Belgrade, Faculty for Special Education and Rehabilitation, Belgrade, Serbia

RECEIVED 09.04.2022.

ACCEPTED 02.10.2022.

Jelena Đorđević<sup>1,2</sup>, Slađana Arsić<sup>4</sup>, Dragan Pavlović<sup>4</sup>, Aleksandra Pavlović<sup>5</sup>

1-Klinika za neurologiju i psihijatriju za decu i omladinu, Beograd

2-Univerzitet u Kragujevcu, Fakultet medicinskih nauka, Kragujevac

3-Akademija vaspitačko-medicinskih strukovnih studija, Odsek medicinskih studija, Čuprija

4-Univerzitet u Beogradu, Beograd, Srbija

5-Univerzitet u Beogradu, Fakultet za specijalnu edukaciju i rehabilitaciju, Beograd

PRIMLJEN 09.04.2022.

PRIHVAĆEN 02.10.2022.

### ABSTRACT

Cognitive dysfunction has been recognized as a key aspect of schizophrenia evident even during remission of symptoms. During the previous decade, interest in cognitive deficits has increased due to the recognition of their key importance for the functional outcome and degree of social adaptation. Prominent cognitive deficits are disorders of memory, attention and executive functions that manifest themselves as a disorder of verbal fluency, problems with serial learning, difficulty in problem solving, and disturbance of executive functions. The observed cognitive deficit is attributed to the dysfunction of cortical-cerebellar-thalamic circuits. Social cognition is the ability to construct an image of relationships between oneself and others and the ability to use flexible mental constructs as a guide to social interactions. The data from previous studies strongly support the association of neurocognitive and socio-cognitive deficits with the performance of independent functioning and quality of life, emphasizing the need for prevention and treatment of cognitive deficits.

Key words: schizophrenia; cognitive dysfunction; social cognition.

### APSTRAKT

Kognitivna disfunkcija je prepoznata kao ključni aspekt šizofrenije, očigledan čak i tokom remisije simptoma. Tokom prethodne decenije povećano je interesovanje za kognitivne deficite zbog prepoznavanja njihovog ključnog značaja za funkcionalni ishod i stepen socijalne adaptacije. Izraženi kognitivni deficiti su poremećaji pamćenja, pažnje i izvršnih funkcija koji se manifestuju kao poremećaj verbalne fluentnosti, problemi sa serijskim učenjem, teškoće u rešavanju problema i poremećaj izvršnih funkcija. Uočeni kognitivni deficit se pripisuje disfunkciji kortikalno – cerebelarno - talamičkih kortikalnih kola. Socijalna kognicija je sposobnost da se konstruiše slika odnosa između sebe i drugih i sposobnost korišćenja fleksibilnih mentalnih konstrukata kao vodiča za društvene interakcije. Podaci iz prethodnih studija snažno podržavaju povezanost neurokognitivnih i sociokognitivnih deficita sa performansama samostalnog funkcionisanja i kvalitetom života, naglašavajući potrebu za prevencijom i lečenjem kognitivnih deficita.

Ključne reči: šizofrenija; kognitivna disfunkcija; socijalna kognicija.

### CORRESPONDENCE / KORESPONDENCIJA

Jelena Djordjevic, Clinic of Mental Disorders "Dr Laza Lazarević", Višegradska 26, 11000, Belgrade, Serbia, Phone +381 60 3232103, e-mail: jelenadjordjevic2000@yahoo.com

Jelena Đorđević, Klinika za psihijatrijske bolesti "Dr Laza Lazarević", Višegradska 26, 11000 Beograd, Tel. 060 3232103, e-mail: jelenadjordjevic2000@yahoo.com

## INTRODUCTION

Cognitive deficits have been recognized as a key feature of schizophrenia since the time of Krepelin<sup>1</sup>. Krepelin emphasized his concept of key characteristics of the disorder by the name he chose for it: it is a disease that begins at an early age ("precox") and has a relatively chronic course characterized by significant cognitive and social damage ("dementia")<sup>1</sup>. The neuropsychological theory begins with the assumption that psychotic disorders are a consequence of cerebral dysfunction and that clinical symptomatology has neurofunctional correlates, that is, it is possible to link a psychological deficit with a site of brain damage<sup>2</sup>.

Cognitive processes are the fundamental basis of psychological, behavioral and interpersonal aspects of human interactions<sup>3</sup>. Cognition includes every mental process that can be described as an experience of knowing, observing, recognizing, reasoning, and concluding<sup>3</sup>. Within the cognitive functioning, two domains are taken into consideration: the domain of neurology and social cognitive domains. Conceptually, neuroscience includes the acquisition of general information and processing functions such as attention, memory, information processing speed, language, speech and executive functions<sup>4</sup>.

Social cognition is the ability to construct an image of relationships between oneself and others and the ability to use flexible mental constructs as a guide to social interactions, with the aim of solving adaptive problems within complex social behavior<sup>5</sup>. The capacity for presenting one's own mental state and the mental state of another person is a key feature of social competence<sup>6</sup>. Social cognition is presented as a multidimensional construct which includes different subcomponents: 1) Theory of the mind (ToM), 2) Social perception, 3) Social knowledge, 4) The recognition of emotions, and 5) Attributional style<sup>7,8</sup>. Although these domains are generally accepted and represent a construct of social cognition, their boundaries can not be considered absolute and there is a significant overlap between them<sup>8</sup>.

In patients with psychiatric disorders, cognitive errors of internal processes and their interference with external stimuli were observed i.e. the misrecognition of internal content as external content, early conclusions, intolerance towards ambiguity, attention and perception disturbances, delusional beliefs, errors of conclusion<sup>9</sup>, have been observed. The processing of socially relevant information relies on neurocognitive capacities, and therefore it is not clear to which degree these two cognitive domains are independent, given the neuronal correlations of both cognitive constructs. The results of factor analysis suggest that neurocognition and social cognition are two separate fields of sensitivity in psychosis<sup>10-12</sup>.

## COGNITIVE DEFICITS IN SCHIZOPHRENIA

Cognitive dysfunction has been recognized as a key aspect of schizophrenia evident even during symptom remission<sup>13,14</sup>. During the previous decade, interest in cognitive deficits had been increased due to the recognition of their key importance for the functional outcome and degree of social adaptation<sup>15</sup>. Prominent cognitive deficits are disorders of memory, attention and executive functions that manifest themselves as a disorder of verbal fluency, problems with serial learning, issues with problem solving, and dysfunction of executive functions (problems with maintaining and focusing attention, concentration, prioritization, and modulating behavior in relation on social hints)<sup>16,17</sup>. The observed cognitive deficit is attributed to the dysfunction of cortical-cerebellar-thalamic circuits<sup>18</sup>. Attention deficit is considered to be the primary cognitive deficit in schizophrenia, also observed in the pre-morbid period<sup>19</sup>. Schizophrenia is a neurodevelopmental disorder, but the etiopathogenesis of cognitive deficits is still unknown. The assumption is that changes in the brain represent a cumulative effect of neurodevelopmental abnormalities, changes in neuroplasticity and changes in neural maturing<sup>20</sup>. Changes in the neuroplasticity of brain-mediated neurotrophic growth hormone (BDNF) hippocampal cells contribute to the formation of cognitive deficits<sup>21</sup>.

Schizophrenia is to some extent characterized by cognitive heterogeneity, which is confirmed by the coexistence of a smaller group of schizophrenic patients with preserved cognitions<sup>22,23</sup>. The existence of a group of neuropsychologically normal individuals with schizophrenia is incompatible with the idea that the damage to neurocognition is the main feature of schizophrenia. Therefore, the existence of such a subgroup could have important implications for efforts to understand not only the neuropathology of this disorder, but also the etiological heterogeneity of this disorder and the possible implications for treatment<sup>24</sup>. In other words, if cognitive impairment is understood as a fundamental feature, then it should have occurred in every patient, and only the degree of its severity may vary<sup>25</sup>.

Considering the existence of cognitive heterogeneity, stratification into four cognitive subgroups within schizophrenia is proposed; two extreme subgroups (one with a nearly normal cognitive profile and one with extremely worsened cognitive performance) and two transitional categories with moderate cognitive-domain dysfunctions<sup>26</sup>. Only a few studied the prevalence of otherwise normal neuropsychologically status of patients with schizophrenia, and the results of these studies were inconsistent and estimates of prevalence vary between 55 and 2%<sup>22, 24</sup>. However, Kremen and associates point out that even neuropsychologi-

cally normal patients have poorer neuropsychological performance compared to the expected or premorbid level of functioning<sup>22</sup>.

Studies that did research of cognitive deficits in different stages of illness (first episode, repeated episodes, chronic stage) confirm the existence of cognitive deficits in all phases, including healthy relatives. Within the neurodevelopmental model, it is considered that these deficits exist even before the onset of full psychosis, while the persistence of deficits that is more prominent in chronic groups confirms current neurodevelopmental and neurodegenerative changes<sup>27</sup>.

Magnetic resonance functional studies have documented that patients with schizophrenia have reduced blood flow and glucose utilization in a prefrontal cortex, a concept known as hypofrontality. Hypofrontality is functionally expressed in the deficiencies of working memory and exquisite functions<sup>28, 29</sup>. Despite extensive literature on hypofrontality in schizophrenia, neuronal mechanisms that are at the core of this pathophysiological trait remain unknown. Possible causes include damaged synaptic association and neurotransmission resulting from neurodevelopmental and/or genetic factors. These disorders can, at least partially, explain the dysregulation of dopamine activity in the limbic striatum and mediodorsal thalamus, which have reciprocal relationships with the prefrontal cortex (cortical-cerebellar-thalamic circuits). The activity in this neural circuit is regulated by gabaergic neurons by producing synapses with glutamatergic pyramidal neurons in the prefrontal cortex<sup>30</sup>. Gabaergic interneurons containing parvalbumin as markers of these neurons are particularly relevant in schizophrenia studies<sup>31</sup>. In addition, this network activity is associated with EEG gamma rhythms and cognitive processing. Indeed, patients with schizophrenia show deficiency in gamma rhythms<sup>32</sup>, which increases the possibility that dysfunction of gabaergic neurons (especially parvalbumin-containing cells) is an important contributor to the dysfunction of the prefrontal cortex in the disease and the onset of cognitive deficits.

The relationship between cognitive abilities and positive and negative psychopathology in schizophrenia has been the subject of several authors' studies. There is a general consensus that positive psychotic symptoms are not associated with cognitive performance<sup>33,34</sup>. However, the relationship between negative symptoms and cognition is less clear. Individually, the authors of the studies recorded inconsistent conclusions as to the possible relationship between negative symptoms and cognitive abilities. The relationship between them is described by some authors<sup>35-38</sup>, but not by all<sup>39-41</sup>. The results of meta-analytic studies have shown the

existence of a small to moderate correlation<sup>33,34</sup>. Considering the complex relationship, models have been suggested in which it is proposed that negative symptoms and cognitive dysfunction have a common etiology<sup>42</sup>.

Most of the social cognitive studies in schizophrenic patients were conducted in relation to the three most commonly investigated domains: Theory of the Mind (ToM), the emotional perception / recognition and attributional style<sup>43</sup>, and therefore the research does not provide complete and consistent answers. Studies that dealt with ToM in schizophrenia have almost the same conclusions that people with schizophrenia show clear deficiencies in ToM in relation to healthy control subjects and patients with other psychiatric disorders<sup>44,46</sup>. Given that first-degree sufferers of schizophrenia also report disturbances in ToM, this finding suggests that ToM deficiency could be a potential endophenotype for schizophrenia<sup>47</sup>. By examining deficits through different stages of the disease, the existence of deficits throughout all stages of the disease was confirmed, and that the deficit at the onset of the disease was comparable to the deficit observed in the chronic stage of the disease<sup>48</sup>. However, studies still can not give a precise answer as to whether these deficits are "straight" or "trait" markers<sup>49,50</sup>. Also, meta-analytical studies of socio-cognitive deficits in schizophrenia confirm the existence of deficits<sup>51</sup> and emphasize the existence of a direct relationship between a sociocognitive deficit and a functional outcome<sup>11,52</sup>.

Although recent modeling studies suggest that social cognition is independent of negative symptoms<sup>53,54</sup>, other studies show the association of negative symptoms with the results of social cognitive tests<sup>55</sup>. Relationships with positive symptoms are similarly inconsistent and unclear<sup>56</sup>. Somewhat higher consistency in literature was found for associations between attributive style and delusion or paranoid belief<sup>57,58</sup>. Deficiencies identified during the symptomatic remission period confirm the independence of the clinical condition deficit, while on the other hand there are findings confirming the deterioration of the performances during the acute disease<sup>59</sup> and indicate the partial correlation of the specific characteristics of the disease, such as the negative symptoms<sup>60</sup> and disorganization<sup>61</sup>.

The data from previous studies strongly supports the correlation of neurocognitive and sociocognitive deficits with the performance of independent functioning and quality of life, emphasizing the need for the prevention and treatment of cognitive deficits<sup>62,63</sup>. Lam et al.<sup>64</sup> in the study of the relationship between neurology and social cognition in schizophrenia attempted to check the hypothesis that the theory of mind and empathy could mediate between neurocognitive abilities (executive function or perception)

and clinical symptomatology (negative, positive and general symptoms). In an attempt to clarify the relationship between neurocognitive abilities and negative or positive symptoms in schizophrenia, it has been found that ToM (Theory of Mind) is a complete intermediate<sup>11</sup>. In particular, the findings of the research indicate that impairments in perception or executive functioning worsen the interpretation and understanding of mental states or intentions of others. Such a deficit, according to Fet and associates, can condition the emergence of negative symptoms<sup>11</sup>. In other words, deficits in the organization, identification and interpretation of sensory information in presenting and understanding of the environment, as well as the skills of problem solving, influence the ability to detect social cues of "Faux pas" in interindividual interactions. On the other hand, there is a backward impact on social functioning, disabling pleasure in social interactions. In addition, the deficit of executive functions may predispose the emergence of positive symptoms by increasing ToM deficiency in schizophrenia<sup>11</sup>. In general, the conclusion would be that the lack of control and regulation of cognitive processes, such as work memory, reasoning, flexibility in solving tasks in people with schizophrenia, aggravates the ability to detect social indications and "Faux pas". This further leads to several psychopathological outcomes such as hallucinations, delusional beliefs, and bizarre behavior<sup>64</sup>.

## NEUROANATOMIC BASES OF COGNITIVE DEFICITS

In studies of diffuse structural abnormalities, the expansion of the third chamber and lateral chambers of the cerebrospinal fluid system is the most consistent of the abnormalities in schizophrenia<sup>65</sup>. Although this is a non-specific marker, ventricular enlargement is associated with impaired performances in neuropsychological tasks and possibly followed by negative symptoms<sup>66</sup>. Findings indicate that abnormalities during neurological development affect neuronal migration, survival and interconnection<sup>67</sup>. In functional neuroimaging studies, there was no localized center for hallucinations or delusional beliefs, but a contribution has been observed of the cortex, frontal areas, cingulate gyros, and subcortical structures<sup>68</sup>.

From the standpoint of cognitive neuropsychology, the occurrence of auditory hallucinations is explained as a monitoring disorder internal speech<sup>68</sup>. The source of hallucinations themselves is not clear, but it is assumed that they are externalized thoughts due to the disturbance of internal speech monitoring mediated by normal premotor cortex, cerebellar cortex, lentiform cores, thalamus, hippocampus, temporal cortex<sup>4</sup>. Patients with schizophrenia and auditory verbal hallucinations have, according to measurements of

magnetic resonance, reduced volume of the upper temporal gyrus and wider cerebral chambers, where those with more intense hallucinations have a lower volume of the upper temporal convolution<sup>69</sup>. Voxel MR volumetry shows that the reduction of the left Heschl gyrus is associated with hallucinations, while the reduction of the left temporal planum associated with delusional ideas<sup>70</sup>. Functional tests link distortions of reality with activation in the temporal lobe and the frontal Cingular cortex<sup>68</sup>. However, other researchers document that wider areas of the brain can be active during auditory hallucinations, and that there is activation of the corresponding sensory areas in the onset of hallucinations in this modality<sup>71</sup>.

In terms of neuroanatomical correlates of observed social cognitive deficits, Bas and associates consider social cognition to be associated with the shift of brain activity into the medial orbitofrontal cortex, amygdala, and right insula<sup>72</sup>. Studies by Eilman and associates have shown a decrease in amygdala activity compared with the control group, which is in line with the results of previous research<sup>73</sup>. It is believed that the amygdala and the orbitofrontal cortex have a significant contribution in the facial expression process, while the medial and orbitofrontal cortex are related to the Theory of mind. The front cingular gyrus is a key component that controls affective and cognitive functions. As part of the rostral limbic system, ACG modulates internal emotional responses, and is considered an integral component of various exquisite functions as well as an essential component of social cognition or mentalization<sup>74</sup>. A disorder in this region can be associated with social cognition and psychopathology in schizophrenia<sup>75</sup>.

## CONCLUSION

Cognitive impairment is the basic domain of schizophrenia dysfunction. Relatively stable and permanent cognitive deficiencies limit the functional recovery capacity. The premorbid existence of cognitive deficits in people who will later develop schizophrenia, as well as the presence of defects in close non-infected relatives suggests that cognitive deficits may be a marker of vulnerability to schizophrenia. Considering the complex relationship between symptomatology and cognitive deficits, and their impact on the degree of functionality, neurocognitive rehabilitation represents a significant focus of treatment.

## REFERENCES

1. Green MF, Harvey P. Cognition in schizophrenia: past, present, and future. *Schizophr Res Cogn* 2014; 1: e1–e9.
2. Viney, Wayne. A history of psychology: ideas and context. 3rd ed. Boston: Pearson, 2003.
3. Blomberg O. Concepts of cognition for cognitive engineering. *Int J Aviat Psychol* 2011; 21: 85–104.
4. Pavlović D. Neuropsychology, behavioral neurology, and neuropsychiatry. Belgrade: Orion Art, 2013. (in Serbian).
5. Adolphs R. Investigating the cognitive neuroscience of social behavior. *Neuropsychologia* 2003; 41: 119–26.
6. Mazza M, De R A, Surian L, Roncone R, Casacchia M. Selective impairments of theory of mind in people with schizophrenia. *Schizophr Res* 2001; 47: 299–308.
7. Ochsner KN. The social-emotional processing stream: five core constructs and their translational potential for schizophrenia and beyond. *Biol Psychiatry* 2008; 64: 48–61.
8. Green MF, Penn DL, Bentall R, et al. Social cognition in schizophrenia: an NIMH workshop on definitions, assessment, and research opportunities. *Schizophr Bull* 2008; 34: 1211–20.
9. Fletcher PC, Frith CD. Perceiving is believing: a Bayesian approach to explaining the positive symptoms of schizophrenia. *Nat Rev Neurosci* 2009; 10: 48–58.
10. Sergi MJ, Rassovsky Y, Widmark C, et al. Social cognition in schizophrenia: relationships with neurocognition and negative symptoms. *Schizophr Res* 2007; 90: 316–24.
11. Fett AKJ, Viechtbauer W, Dominguez MD, Penn DL, van Os J, Krabbendam L. The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neurosci Biobehav Rev* 2011; 35: 573–88.
12. Zhang T, Cui H, Tang Y, et al. Correlation of social cognition and neurocognition on psychotic outcome: a naturalistic follow-up study of subjects with attenuated psychosis syndrome. *Sci Rep* 2016; 10: 35017.
13. Keshavan M, Montrose DM, Rajarethinam R, Diwadkar V, Prasad K, Sweeney JA. Psychopathology among offspring of parents with schizophrenia: Relationship to premorbid impairments. *Schizophr Res* 2008; 103: 114–20.
14. Lewandowski KE, Cohen BM, Ongur D. Evolution of neuropsychological dysfunction during the course of schizophrenia and bipolar disorder. *Psychol Med* 2011; 41: 225–41.
15. Bowie CR, Harvey PD. Cognitive deficits and functional outcome in schizophrenia. *Neuropsychiatr Dis Treat* 2006; 2: 531–6.
16. Cornblatt BA, Keilp JG. Impaired attention, genetics, and the pathophysiology of schizophrenia. *Schizophr Bull* 1994; 20: 31–46.
17. Kuperberg G, Heckers S. Schizophrenia and cognitive function. *Curr Opin Neurobiol* 2000; 10: 205–10.
18. Andreasen NC, Paradiso S, O’Leary DS. “Cognitive dysmetria” as an integrative theory of schizophrenia: a dysfunction in cortical-subcortical-cerebellar circuitry?” *Schizophr Bull* 1998; 24: 203–18.
19. Caspi A, Reichenberg A, Weiser M, et al. Cognitive performance in schizophrenia patients assessed before and following the first psychotic episode. *Schizophr Re* 2003; 65: 87–94.
20. Gourion D, Gourevitch R, Leprovost JB, Olié H, Lôo JP, Krebs MO. Neurodevelopmental hypothesis in schizophrenia. *Encephale* 2004; 30: 109–18. (in French).
21. Nieto R, Kukuljan M, Silva H. BDNF and schizophrenia: from neurodevelopment to neuronal plasticity, learning, and memory. *Front Psychiatry* 2013; 4: 45.
22. Kremen WS, Seidman LJ, Faraone SV, Toomey R, Tsuang MT. The paradox of normal neuropsychological function in schizophrenia. *J Abnorm Psychol* 2000; 109: 743–52.
23. Reichenberg A, Harvey PD, Bowie CR, et al. Neuropsychological function and dysfunction in schizophrenia and psychotic affective disorders. *Schizophr Bull* 2009; 35: 1022–9.
24. Heinrichs RW, Zakzanis KK. Neurocognitive deficit in schizophrenia: a quantitative review of the evidence. *Neuropsychology* 1998; 12: 426–45.
25. Totić-Poznanović S, Pavlović DM, Đorđević JR, Pavlović AM, Marinković DJ. Neuropsychological assessment and treatment possibilities of cognitive deficit in schizophrenic patients. *Vojnosanit Pregl* 2012; 69: 510–6.
26. Hill SK, Ragland JD, Gur RC, Gur RE. Neuropsychological profiles delineate distinct profiles of schizophrenia, an interaction between memory and executive function, and uneven distribution of clinical subtypes. *J Clin Exp Neuropsychol* 2002; 24: 765–80.
27. Corigliano V, De Carolis A, Trovini G, et al. Neurocognition in schizophrenia: from prodrome to multi-episode illness. *Psychiatry Res* 2014; 220: 129–34.
28. Buchsbaum MS, Nuechterlein KH, Haier RJ, et al. Glucose metabolic rate in normals and schizophrenics during the continuous performance test assessed by positron emission tomography. *Br J Psychiatry* 1990; 156: 216–27.
29. Hazlett EA, Buchsbaum MS, Jeu LA, et al. Hypofrontality in unmedicated schizophrenia patients studied with PET during performance of a serial verbal learning task. *Schizophr Res* 2000; 43: 33–46.
30. Zhu Y, Stornetta RL, Zhu JJ. Chandelier cells control excessive cortical excitation: characteristics of whisker-evoked synaptic responses of layer 2/3 nonpyramidal and pyramidal neurons. *J Neurosci* 2004; 24: 5101–8.
31. Lewis DA. GABAergic local circuit neurons and prefrontal cortical dysfunction in schizophrenia. *Brain Res Brain Res Rev* 2000; 31: 270–6.
32. Cho RY, Konecky RO, Carter CS. Impairments in frontal cortical gamma synchrony and cognitive control in schizophrenia. *Proc Natl Acad Sci USA* 2006; 103: 19878–83.

33. Dominguez Mde G, Viechtbauer W, Simons CJ, van Os J, Krabbendam L. Are psychotic psychopathology and neurocognition orthogonal? A systematic review of their associations. *Psychol Bull* 2009; 135: 157–71.
34. Ventura J, Helleman GS, Thames AD, Koellner V, Nuechterlein KH. Symptoms as mediators of the relationship between neurocognition and functional outcome in schizophrenia: a meta-analysis. *Schizophr Res* 2009; 113: 189–99.
35. Addington J, Addington D, Maticka-Tyndale E. Cognitive functioning and positive and negative symptoms in schizophrenia. *Schizophr Res* 1991; 5: 123–34.
36. Brazo P, Marie RM, Halbecq I, et al. Cognitive patterns in subtypes of schizophrenia. *Eur Psychiatry* 2002; 17: 155–62.
37. Bozikas VP, Kosmidis MH, Kioperlidou K, Karavatos A. Relationship between psychopathology and cognitive functioning in schizophrenia. *Compr Psychiatry* 2004; 45: 392–400.
38. Heydebrand G, Weiser M, Rabinowitz J, Hoff AL, DeLisi LE, Csernansky JG. Correlates of cognitive deficits in first episode schizophrenia. *Schizophr Res* 2004; 68: 1–9.
39. Minzenberg MJ, Poole JH, Vinogradov S, Shenaut GK, Ober B.A. Slowed lexical access is uniquely associated with positive and disorganised symptoms in schizophrenia. *Cogn Neuropsychiatry* 2003; 8: 107–27.
40. Lucas S, Fitzgerald D, Redoblado-Hodge MA, et al. Neuropsychological correlates of symptom profiles in first episode schizophrenia. *Schizophr Res* 2004; 71: 323–30.
41. Rund BR, Melle I, Friis S, et al. Neurocognitive dysfunction in first-episode psychosis: correlates with symptoms, premorbid adjustment, and duration of untreated psychosis. *Am J Psychiatry* 2004; 161: 466–72.
42. Harvey PD, Koren D, Reichenberg A, Bowie CR. Negative symptoms and cognitive deficits: what is the nature of their relationship? *Schizophr Bull* 2006; 32: 250–8.
43. Penn DL, Addington J, Pinkham A. Social cognitive impairments. In: Lieberman JA, Stroup TS, Perkins DO, eds. *American psychiatric association textbook of schizophrenia*. Arlington: American Psychiatric Publishing Press Inc., 2006: 261–74.
44. Couture SM, Penn DL, Roberts DL. The functional significance of social cognition in schizophrenia: a review. *Schizophr Bull* 2006; 32(Suppl 1): S44–S63.
45. Pinkham AE, Penn D, Perkins DO, Graham K, Siegel M. Emotion perception and social skill over the course of psychosis: a comparison of individuals at risk, and early and chronic schizophrenia spectrum illness. *Cogn Neuropsychiatry* 2007; 12: 198.
46. Koelkebeck K, Pedersen A, Suslow T, Kueppers KA, Arolt V, Ohrmann P. Theory of Mind in first-episode schizophrenia patients: correlations with cognition and personality traits. *Schizophr Res* 2010; 119: 115–23.
47. Irani F, Platek SM, Panyavin IS, et al. Self-face recognition and theory of mind in patients with schizophrenia and first-degree relatives. *Schizophr Res* 2006; 88: 151–60.
48. Bertrand MC, Sutton H, Achim AM, Malla AK, Lepage M. Social cognitive impairments in first episode psychosis. *Schizophr Res* 2007; 95: 124–33.
49. Kettle JW, O'Brien-Simpson L, Allen NB. Impaired theory of mind in first-episode schizophrenia: comparison with community, university and depressed controls. *Schizophr Res* 2008; 99: 96–102.
50. Green MF, Bearden CE, Cannon TD, et al. Social cognition in schizophrenia. Part 1: performance across phase of illness. *Schizophr Bull* 2012; 38: 854–64.
51. Hoekert M, Kahn RS, Pijnenborg M, Aleman A. Impaired recognition and expression of emotional prosody in schizophrenia: review and meta-analysis. *Schizophr Res* 2007; 96: 135–45.
52. Brune M, Abdel-Hamid M, Lehmkamper C, Sonntag C. Mental state attribution, neurocognitive functioning, and psychopathology: what predicts poor social competence in schizophrenia best? *Schizophr Res* 2007; 92: 151–9.
53. Rassovsky Y, Horan WP, Lee J, Sergi MJ, Green MF. Pathways between early visual processing and functional outcome in schizophrenia. *Psychol Med* 2011; 41: 487–97.
54. Sergi MJ, Rassovsky Y, Widmark C, et al. Social cognition in schizophrenia: relationships with neurocognition and negative symptoms. *Schizophr Res* 2007; 90: 316–24.
55. Kohler CG, Hoffman LJ, Eastman LB, Healey K, Moberg PJ. Facial emotion perception in depression and bipolar disorder: a quantitative review. *Psychiatry Res* 2011; 188: 303–9.
56. Woodward TS, Mizrahi R, Menon M, Christensen BK. Correspondences between theory of mind, jumping to conclusions, neuropsychological measures and the symptoms of schizophrenia. *Psychiatry Res* 2009; 170: 119–23.
57. Combs DR, Penn DL, Michael CO, et al. Perceptions of hostility by persons with and without persecutory delusions. *Cogn Neuropsychiatry* 2009; 14: 30–52.
58. Fornells-Ambrojo M, Garety PA. Attributional biases in paranoia the development and validation of the Achievement and Relationships Attributions Task (ARAT). *Cogn Neuropsychiatry* 2009; 14: 87–109.
59. Corcoran R, Rowse G, Moore R, et al. A transdiagnostic investigation of 'theory of mind' and 'jumping to conclusions' in patients with persecutory delusions. *Psychol Med* 2008; 38: 1577–83.
60. Corcoran R, Mercer G, Frith CD. Schizophrenia, symptomatology and social inference: investigating "theory of mind" in people with schizophrenia. *Schizophr Res* 1995; 17: 5–13.
61. Mazza M, De RA, Surian L, Roncone R, Casacchia M. Selective impairments of theory of mind in people with schizophrenia. *Schizophr Res* 2001; 47: 299–308.

62. San L, Ciudad A, Alvarez E, Bobes J, Gilaberte I. Symptomatic remission and social/vocational functioning in outpatients with schizophrenia: prevalence and associations in a cross-sectional study. *Eur Psychiatry* 2007; 22: 490-8.
63. Tuulio-Henriksson A. Cognitive dysfunction in schizophrenia: a familial and genetic approach. Publications of the National Public Health Institute A7/ 2005. Helsinki: National Public Health Institute, 2005.
64. Lam BYH, Raine A, MC Lee T. The relationship between neurocognition and symptomatology in people with schizophrenia: social cognition as the mediator. *BMC Psychiatry* 2014; 14: 138.
65. Yurgelun-Todd DA, Gruber SA, Kanayama G, Killgore WD, Baird AA, Young AD. fMRI during affect discrimination in bipolar affective disorder. *Bipolar Disord* 2000; 2: 237-48.
66. Vita A, Dieci M, Giobbio GM, et al. Language and thought disorder in Schizophrenia: Brain morphological correlates. *Schizophr Res* 1995; 15: 243-51.
67. Tregellas J. Connecting brain structure and function in schizophrenia. *Am J Psychiatry* 2009; 166: 134-6.
68. Allen P, Larøi F, McGuire PK, Aleman A. The hallucinating brain: a review of structural and functional neuroimaging studies of hallucinations. *Neurosci Biobehav Rev* 2008; 32: 175-91.
69. Barta PE, Pearlson GD, Powers RE, Richards SS, Tune LE. Auditory hallucinations and smaller superior temporal gyral volume in schizophrenia. *Am J Psychiatry* 1990; 147: 1457-62.
70. Sumich A, Chitnis XA, Fannon DG, O'Ceallaigh S, Doku VC, Faldrowicz A, Sharma T. Unreality symptoms and volumetric measures of Heschl's gyrus and planum temporal in first-episode psychosis. *Biol Psychiatry* 2005; 57: 947-50.
71. Shergill SS, Brammer MJ, Williams SC, Murray RM, McGuire PK. Mapping auditory hallucinations in schizophrenia using functional magnetic resonance imaging. *Arch Gen Psychiatry* 2000; 57: 1033-8.
72. Baas D, Aleman A, Vink M, Ramsey NE, de Haan EH, Kahn RS. Evidence of altered cortical and amygdala activation during social decision-making in schizophrenia. *Neuroimage* 2008; 40: 719-27.
73. Aleman A, Kahn R. Strange feelings: do amygdala abnormalities dysregulate the emotional brain in schizophrenia? *Prog Neurobiol* 2005; 77: 283-98.
74. Kopelman A, Andreasen NC, Nopoulos P. Morphology of the anterior cingulate gyrus in patients with schizophrenia: relationship to typical neuroleptic exposure. *Am J Psychiatry* 2005; 162: 1872-8.
75. Fujiwara H, Hirao K, Namiki C, et al. Anterior cingulate pathology and social cognition in schizophrenia: a study of gray matter, white matter and sulcal morphometry. *Neuroimage* 2007; 36: 1236-45.