**MATERIAL ON THE ANNOTATED CHECKLIST
OF VASCULAR FLORA OF SERBIA.
NOMENCLATURAL, TAXONOMIC AND FLORISTIC NOTES II**

MARJAN NIKETIĆ^{1*}, GORDANA TOMOVIĆ², GORAN ANAČKOV³, MIRJANA BARTULA⁴, SLADJANA DJORDJEVIĆ⁵, VLADAN DJORDJEVIĆ², SUZANA DJORDJEVIĆ-MILOŠEVIĆ⁶, ŠEMIJA DURAKI⁷, MILAN GAVRILOVIĆ², PEDJA JANAČKOVIĆ², EVA KABAS², NEVENA KUZMANOVIĆ², DMITAR LAKUŠIĆ², PREDRAG LAZAREVIĆ², RANKO PERIĆ⁸, VLADIMIR RANDJELOVIĆ⁹, DRAGIŠA SAVIĆ¹⁰, MIHAJLO STANKOVIĆ¹¹, IVANA STEVANOSKI², VERICA STOJANOVIĆ¹², OLGA VASIĆ¹, SNEŽANA VUKOJIČIĆ², BOJAN ZLATKOVIĆ⁹, VLADIMIR STEVANOVIĆ¹³

¹ Natural History Museum, Njegoševa 51, 11000 Belgrade, Serbia,
e-mail: mniketic@nhmbeo.rs, oljav27@gmail.com

² Institute of Botany and Botanical Garden, Faculty of Biology, University of
Belgrade, Serbia, e-mail: gtomovic@bio.bg.ac.rs

³ Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad,
Serbia, e-mail: goran.anackov@dbe.uns.ac.rs

⁴ Faculty of Applied Ecology Futura, Metropolitan University, Serbia

⁵ Faculty of Agriculture, University of Belgrade, Serbia

⁶ Singidunum University, Serbia

⁷ Eight Belgrade Grammar School, Grčica Milenka 71, 11000 Belgrade, Serbia

⁸ Institute for Nature Conservation of Vojvodina province, Novi Sad, Serbia,
e-mail: ranko.peric@pzzp.rs

⁹ Department of Biology and Ecology, Faculty of Natural Sciences and
Mathematics, University of Niš, Serbia,
e-mail: vladar@pmf.ni.ac.rs, bojanzlat@pmf.ni.ac.rs

¹⁰ National Park "Fruška Gora", Sremska Kamenica, Serbia,
e-mail: dragita@gmail.com

¹¹ Nature Conservation Movement of Sremska Mitrovica, Serbia,
e-mail: trogloxen@gmail.com

¹² Institute for Nature Conservation of Serbia, Belgrade, Serbia,
e-mail: verica.stojanovic@zzps.rs

¹³ Serbian Academy of Sciences and Arts, Belgrade, Serbia,
e-mail: vstev@bio.bg.ac.rs

The paper propose nine new nomenclature combinations and indicate eight new synonyms. There are 12 taxa (species and subspecies) of vascular flora which are new for the flora of Serbia (of which nine are autochthonous and three

allochthonous plants). An old or unreliable literature records for three autochthonous and one allochthonous plant in the flora of Serbia has been confirmed. Eight plant taxa are a novelty for proper Serbia, Vojvodina, or Kosovo and Metohija, and for three plants, presence in certain administrative units in Serbia has been proved. The last part of the paper contains information on eight taxa whose presence in Serbia or in its territorial units has been refuted.

Key words: vascular flora, checklist, nomenclature, taxonomy, Serbia

INTRODUCTION

This paper represents a continuation of the inventory of the flora of Serbia through intensive field research, identification and revision of specimens from herbarium collections, as well as a comprehensive review of extensive taxonomic, floristic and phytocoenological literature. The first such article was published two years ago (Niketić *et al.* 2018a) as a preparation for the publication of *An annotated checklist of vascular flora of Serbia 1* (Niketić & Tomović 2018). In that paper, new data on taxa were integrated, which included the following groups: Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida.

In the meantime, a number of floristic and chorological contributions have been published (Bogosavljević & Zlatković 2018, Duraki & Niketić 2018a,b,c,d, Jovanović *et al.* 2018, Lazarević & Zlatković 2018, Radak *et al.* 2019, Süveges *et al.* 2019, Zlatković & Bogosavljević 2020, Sabovljević *et al.* 2020; Tomović *et al.* 2020), as well as taxonomic (Novaković *et al.* 2018) and phylogeographic ones (Janković *et al.* 2019) related to plant taxa from Serbia, which also contributed to better knowledge of the presence and distribution of native and adventive flora in our country.

The second paper presents nomenclatural, taxonomic and floristic notes related to taxa from the Magnoliopsida group, as well as some novelties that have not been published in the *An annotated checklist of vascular flora of Serbia 1* (Niketić & Tomović 2018) because in the meantime some significant findings for the flora of Serbia or certain administrative units have emerged.

MATERIAL AND METHODS

In order to accomplish all the floristic and distribution data we performed intensive field investigation, identification and revision of herbarium material as well as checked numerous literature sources. Collected plant specimens are stored in the Herbarium of the Natural

History Museum in Belgrade (BEO), Herbarium of the Institute of Botany and Botanical Garden “Jevremovac”, University of Belgrade (BEOU), the Herbarium of Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad (BUNS), the Herbarium of the Institute for Nature Conservation of the Vojvodina province (PZZP) (Thiers 2021) and Herbarium of the Faculty of Sciences and Mathematics, Department of Biology and Ecology, University of Niš (HMN).

The names and classification of families and affinity of genera to certain families were presented in accordance with APG IV (2016) for angiosperm families (flowering plants), with certain deviations in the taxa of the highest rank. The nomenclature is harmonized with modern taxonomic studies for particular groups and new taxonomic and/or nomenclature solutions were also presented. Families, genera and species were listed in the alphabetical order of their Latin names. Short taxon description for newly registered taxa for Serbia is accomplished by using numerous botanical literature, with minor modifications and changes which are the result of herbarium specimens inspections and field observations.

In the nomenclature lists heterotypic and homotypic names are sorted chronologically. Nomenclatural provisions are applied from ICN (Turland *et al.* 2018).

Invasiveness status was determined using the terminology according to Niketić & Tomović (2008).

Distribution of the studied taxa within the territory of the Republic of Serbia was presented and mapped according to the grid map with squares of 10×10 km, based on the Military Grid Reference System (MGRS) projection (Lampinen 2001).

RESULTS AND DISCUSSION

Nomenclatural notes

LILIOPSIDA

IRIDACEAE

Crocus tommasinianus Herb., J. Hort. Soc. London 2: 273 (1847).

= *C. panicii* Janka, Oesterr. Bot. Z. 18: 57 (1868), p.p.?, **syn. nov.**

= *Crocus heuffelianus* var. *csapodyae* Horvát & Jáv. in Jáv., Bot. Közlem. 51(4): 180 (1964).

– *C. serbicus* A. Kern. ex Maw, Gard. Chron., n.s. 16: 368 (1881), nom. inval. (pro syn. of *C. tommasinianus* Herb.).

– “*C. minimus*” sensu Pančić, Verh. K.K. Zool.-Bot. Ges. Wien 6: 578 (1856) [non DC. Redouté, Liliac. 2: 81 (1804)].

In this article, the type of *C. pancicii* Janka was included in *C. tommasinianus* Herb. The first description and diagnosis of Janka's plant were presented by Pančić (1856) under the name "*C. minimus* DC." (from *C. biflorus* group) and it was recorded for the surroundings of Jagodina (Sugudbina village) in Šumadija region of Serbia. The geographical position of the locality was interpreted as 20 km west of Jagodina in Niketić *et al.* (2018b), referring to the village of Velika Sugudbina. However, Pančić (1874, 1884) specified that the locality is between the village Oparić and [Mala] Sugudbina, which is 25 km south of Velika Sugudbina, closer to Trstenik than Jagodina. Pančić claimed that this plant could be a new species to science, from the *C. vernus* group, and he did not bring it into connection with representatives of the *C. biflorus* group. Even earlier, based on Pančić's original description of "*C. minimus* DC." Janka (1868) described *C. pancicii* and mentions it also for Karlovo in C Bulgaria. Although so far the original Pančić's material has not been found, nor has any *Crocus* species been found at Pančić's locality, numerous populations of *C. tommasinianus* have been observed only 8 km southeast, between villages Lazarevac and Konjuh. Having in mind the dominant presence of this species in a wider area, it can be concluded that the name *C. pancicii* actually corresponds to *C. tommasinianus*.

Collected specimen:

C Serbia, Kruševac, Lazarevac village, 211 m, 21.163889 E, 43.656389 N, MGRS 34T EP13, coll. V. Randjelović, J. Stojanović 08-Mar-2019, det. V. Randjelović (HMN 13911).

V. Randjelović, M. Niketić

MAGNOLIOPSIDA

ASTERACEAE (COMPOSITAE)

Amphoricarpos neumayerianus subsp. *autariatus* (Blečić & Mayer) Niketić, Gavrilović & Janačković, **comb. nov.**

≡ *Amphoricarpos autariatus* Blečić & E. Mayer [subsp. *autariatus*], Phytion (Horn) 12: 155 (1967) [basionym].

= *Amphoricarpos autariatus* subsp. *bertisceus* Blečić & E. Mayer, Phytion (Horn) 12: 156 (1967), **syn. nov.**

– *A. neumayeri* var. *intermedia* Fukarek, Glasn. Zemaljsk. Muz. Bosne Hercegovine Sarajevu. Prir. Nauke 3–4: 161 (1965), nom. inval. (nom. nud.).

Amphoricarpus neumayerianus subsp. *murbeckii* (Bošnjak) Niketić, Gavrilović & Janačković, **comb. nov.**

≡ *Amphoricarpus neumayeri* subsp. *murbeckii* Bošnjak, Glasn. Hrvatsk. Prir. Društva 41-48: 62 (1936) [basionym] ≡ *A. neumayeri* var. *murbeckii* (Bošnjak) Fukarek, Glasn. Zemaljsk. Muz. Bosne Hercegovine Sarajevu. Prir. Nauke 3-4: 161 (1965), comb. inval.

= *Amphoricarpus neumayeri* var. *velezensis* Murbeck, Acta Univ. Lund. 27: 100 (1892).

= *A. neumayeri* f. *latifolius* Beck, Jahreskat. Wiener bot. Tauschverein: 20 (1894).

In the traditional botanical literature sources, one or two species of the relict orophytic genus *Amphoricarpus* Vis. are most often mentioned for the Balkan Peninsula. In two species concept (Blečić & Mayer 1967, Greuter 2006+) the type species of the genus, *A. neumayerianus* (Vis.) Greuter, was mainly known from SW oromediterranean Montenegro (Mt Orjen and Mt Lovćen) including the bordering part of Herzegovina. Previously long-used invalid name, *A. neumayeri* Vis., has been replaced by Greuter (2003). The second 'species', *A. autariatus* Blečić & E. Mayer, was mentioned for a wider area of the western Balkans, from Bosnia and Herzegovina to NW Greece, including two subspecies: *autariatus* (incl. *A. neumayeri* subsp. *murbeckii* Bošnjak) and *bertisceus* Blečić & E. Mayer. In a single species concept (Webb 1976, Tan 1991) there were two subspecies: "*neumayeri*" and *murbeckii* (described from Herzegovina, but by its distribution equivalent to the then known *A. autariatus*). In the recent comprehensive molecular and morphometric study of the representative populations of the genus, Caković *et al.* (2015) has shown that all of these taxa can be considered conspecific and without intraspecific taxa.

However, we believe that some obvious differences should not be overlooked and that at least three subspecies can be identified within a common species, *A. neumayerianus*. This primarily refers to a clear genetic split (ITS, AFLP) between the northwestern (from mountains around the Neretva river in Herzegovina and Mt Vlašić in Bosnia) and other populations. Herzegovinian plants were previously included within subsp. *murbeckii* (Bošnjak 1936, Webb 1976, Tan 1991), but with an illegitimate specific epithet (Art. 52.1) which was corrected on this occasion. The subspecies *murbeckii* is characterized by denser indumentum and fewer and smaller cauline leaves. The rest of the species range includes the type subspecies present in Mt Orjen, Mt Lovćen and Mt Njegoš, and subsp. *autariatus* (Blečić & Mayer) Niketić, Gavrilović & Janačković that occupies the largest part of the species distribution area (including SW Serbia). The type subspecies (*neumayerianus*) was broadly accepted taxon (at different ranks) in all traditional classifications. It has relatively narrow

leaves, more acute phyllaries and winged cypselae. It is also separated genetically (AFLP and partially in plastid data) and morphologically (PCA, DCA) in Caković *et al.* (2015). Transitional populations (Mt Maganik, Mt Sinjavina, Mt Žijevo, Mt Komovi) were observed in the overlapping zone by Fukarek (1965) (var. *intermedia*, nom. nud.) and also confirmed genetically (Caković *et al.* 2015) (AFLP). In this study not enough diagnostic characters were observed for “subsp. *bertisceus*” and therefore it was included in subsp. *autariatus*. Southeastern population from S Albania and NW Greece are also genetically isolated (Caković *et al.* 2015), but due to insufficient evidence, their status remained unresolved. Consequently, proposed classification recognizes three subspecies and it is basically similar to that proposed by Webb (1976) and Tan (1991), except the separation of subsp. *murbeckii* and subsp. *autariatus*.

Recent morphometric and molecular analyzes (Gavrilović 2019) and metabolomic study (Cvetković *et al.* 2018) of the Balkan representatives of the genus *Amphoricarpos* also confirmed our taxonomic treatment. According to Cvetković *et al.* (2018) guaianolide sesquiterpene lactones (amphoricarpolides), as reliable chemotaxonomic markers, most contribute to differences between taxa *neumayerianus* [“*neumayeri*”] and *autariatus* while the compounds that contributed to the differences between “subsp. *autariatus*” and “subsp. *bertisceus*” “do not have chemotaxonomic significance”.

M. Niketić, M. Gavrilović, P. Janačković

***Hieracium brachytrichophyes* (O. Behr, E. Behr & Zahn) Niketić, comb. et stat. nov.**

≡ *Hieracium sparsum* subsp. *brachytrichophyes* O. Behr, E. Behr & Zahn, Glasnik Skopskog Naučnog Društva 20, Sect. Sci. Nat. 7: 34 (1939) [basionym].

IND. LOC.: “Korab: Hänge an der Waldgrenze an der Dlaboka [“Luboka”] Reka, ca. 1600–1700 m”. LECTOTYPE (**here designated**): “Macedonia, in montibus Korabensibus, ad declives praeter Dlaboka [“Luboka”] reka prope finem si[l]varum, ca. 1600 m.”, 23-Jul-1937, E. Behr (987 S 13-19905 [photo!]) image available at <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.s12-19905>; isoelectotype: M.

Until recently, this plant was considered very rare and only recorded for Mt Korab in W North Macedonia (Behr *et al.* 1939a). A few years ago Duraki & Niketić (2018a) discovered many populations on Šar Planina Mts on the border of North Macedonia and Serbia (Metohija and Kosovo). The taxon belongs to Alpine-Carpatho-Balkan-W Asian complex *H. djimilense* (*prenanthoides-sparsum*) which includes 17 taxa (Greuter 2006+).

distribution of other *Hieracium* species in the environment, the new proposed combination *H. pannosum* s.l. – *scardicum* (*pannosum* s.l. > *naegelianum*) and specific rank would be much more appropriate. The characteristic dark hairs at the base of the involucre are somewhat noticeable and are probably inherited from *H. naegelianum* Pančić. The specific rank of *H. cernyanum* and its exclusion from *H. gaudryi* were indirectly pointed out by Buttler (1991), who considered *H. gaudryi* to be endemic to Greece: “Plants referred to the species by Zahn (1921, 1936) under different subspecific and varietal names do not belong here”. *Hieracium cernyanum* also includes forms with denser indumentum, described as *Hieracium pannosum* subsp. *doerflerianum* Hayek & Zahn and var. *pantocsekianum* Hayek & Zahn, from the same locality (Zahn 1921). The connection of these plants with *H. cernyanum* was also emphasized in the protologue of subsp. *doerflerianum* (“inter *Cernyanum* and *pannosum*”), enabling inclusion of subsp. *doerflerianum* and var. *pantocsekianum* in *H. cernyanum* on this occasion. Plants with varietal name were also recorded for N Albania (Mnëla) (Zahn 1936). Closely related species to *H. cernyanum* is *H. trikalense* Buttler from Pindus Mts in Greece.

M. Niketić

***Hieracium delpinoi* Bald., Malpighia 5: 76 (1891) [“*H. delpinii*”].**

≡ *Hieracium waldsteinii* subsp. *delpinoi* (Bald.) Zahn in Rchb., Icon. Fl. Germ. Helv. 19(2): 112 (1907).

IND. LOC.: “In rupestribus travertineis ad coenobium Moracae”. LECTOTYPE (**here designated**): [Montenegro] “In rupestribus travertineis ad coenobium Moracae”, Aug-1890, *A. Baldacii* [“*H. delpinii*”] (E barcode 00413068 [photo!] image available at <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.m0161496>).

= *Hieracium waldsteinii* subsp. *nipholeucum* Zahn in Rchb., Icon. Fl. Germ. Helv. 19(2): 109 (1907), **syn. nov.**

IND. LOC.: “Albanien:...Klementi...Baldacii...VIII...155.”. LECTOTYPE (**here designated**): [Albania] “In saxosis fl. Cem ad Selce distr. Klementi”, 09-Aug-1901, *A. Baldacii* (sub *H. delpinoi*) (155 P barcode 02806108 [photo!] image available at <https://api.gbif.org/v1/image/unsafe/http%3A%2F%2Fmediaphoto.mnhn.fr%2Fmedia%2F1441344678661sdUV3G6moFz71va0>).

A comparison of the type material of *Hieracium delpinoi* Bald. (E 00413068) and *H. waldsteinii* subsp. *nipholeucum* Zahn (P 02806108) showed their great similarity. In addition, there are numerous records in which these two names are listed for the same localities. Zahn (1921, 1936)

included both taxa within *H. waldsteinii* Tausch as separate subspecies within separate groups (*plumulosum* and *suborieni*), but he nevertheless pointed out their similarity. These two morphotypes differ slightly in the density of the indumentum, but this character is very variable. *Hieracium delpinoi* was recorded for Montenegro, Albania, Serbia (Metohija) and NW Greece. Further studies will show whether it needs to be combined with some other taxa from the *H. waldsteinii* complex.

M. Niketić

***Hieracium hayekianum* (Dörfl. & Zahn) Niketić, comb. et stat. nov.**

≡ *Hieracium gaudryi* subsp. *hayekianum* Dörfl. & Zahn in Engl., Pflanz. 76: 570 (1921) [basionym].

IND. LOC.: “Gipfelregion des Paštrik (Doerfler, It. Alb. n. 877)!”.
LECTOTYPE (**here designated**): [Serbian-Albanian border] “Distrikt Hasi, an felsen des Gipfelregion des Paštrik, ca. 1700 m”, 27-Jul-1918, *I. Dörfler* (B barcode 10 9004793 [photo!]) image available at https://api.gbif.org/v1/image/unsafe/https%3A%2F%2Fpictures.bgbm.org%2Fdigilib%2FScaler%2FIIIF%2FB%2F10%2F90%2F04%2F79%2FB_10_9004793.jpg%2Ffull%2F2048%2C%2F0%2Fdefault.jpg.

This taxon is only known from the border of Serbia (Metohija) and Albania (Mt Paštrik) (Zahn 1921, 1936). The reasons for exclusion from *H. gaudryi* are similar to those for *H. cernyanum* (see on page 93). Although it probably has a similar origin [*H. pannosum* s.l. – *scardicum* (*pannosum* s.l. > *naegelianum*)] and morphology as *H. cernyanum*, it has more elongated and acute leaves, which indicates a slightly increased resemblance to *H. naegelianum*.

M. Niketić

***Hieracium oroglaucum* O. Behr, E. Behr & Zahn, Glasnik Skopskog Naučnog Društva 18, Sect. Sci. Nat. 6: 66 (1937).**

= *Hieracium sparsum* subsp. *livadicanum* O. Behr, E. Behr & Zahn, Glasnik Skopskog Naučnog Društva 20, Sect. Sci. Nat. 7: 128 (1939), **syn. nov.**

IND. LOC.: “Felsen am Abhang der Livadica, ca. 1800–1900 m”.
LECTOTYPE (**here designated**): “Macedonia: in saxi clivium Livadicae, in montibus Šarplaninae sitae, ca. 1800–1900 m.”, 23-Jul-1938, *E. Behr* (1178 M barcode 0165172 [photo!]) image available at <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.m0165172>.

This species was previously known only from Mt Korab in W North Macedonia (Behr *et al.* 1937, 1939a) and recently found in several localities in E Montenegro, SW North Macedonia, SW Serbia (Kosovo and Metohija) and SE Serbia (Duraki & Niketić 2018b). The authors of this

polymorphic species interpreted its origin as “*H. naegelianum* > *pseudobifidum* (*transylvanicum-bifidum*)” (Behr *et al.* 1937). Later, the same authors also described *H. sparsum* subsp. *livadicanum* O. Behr, E. Behr & Zahn from the border of North Macedonia and Kosovo in Serbia (Behr *et al.* 1939b). According to them it is similar to *H. sparsum* subsp. *pilosifrons* Zahn from E Serbia. However, it is clearly noticeable on the type material (M 0165172) that the morphology of the involucre generally corresponds to the ‘*H. naegelianum* type’ and that these specimens actually correspond to *H. oroglaucum* O. Behr, E. Behr & Zahn.

M. Niketić

Hieracium platytrigonodon (O. Behr, E. Behr & Zahn) Niketić, **comb. et stat. nov.**

≡ *Hieracium bohatschianum* subsp. *platytrigonodon* O. Behr, E. Behr & Zahn, Glasnik Skopskog Naučnog Društva 18, Sect. Sci. Nat. 6: 65 (1937) [basionym].

IND. LOC.: “Korab: Hänge an der Waldgrenze an der Dlaboka [“Luboka”] Reka, ca. 1600–1700 m”. LECTOTYPE (**here designated**): “Macedonia: in montibus Korabensibus, in fageto ad stanicam Stalna patrola Gabrova dictam, ca. 1780 m.”, 24-Jul-1937, E. Behr (975 M barcode 0161409 [photo!]) image available at <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.m0161409>).

This taxon is only known from two localities: Mt Korab in W North Macedonia (Behr *et al.* 1937) and Crni Vrh peak on Šar Planina Mts in Serbia (Metohija) (Duraki & Niketić 2018c). It belongs to Carpatho-Balkan complex *H. bohatschianum* (*schmidtii* – *sparsum*) which includes several taxa. Due to its elongated basal and reduced cauline leaves it could be considered as a microspecies probably originated from sympatric *Hieracium schmidtii* subsp. *balkanum* (R. Uechtr. ex Pančić) Niketić.

M. Niketić

Hieracium squarrosobracchiatum (O. Behr, E. Behr & Zahn) Niketić, **comb. et stat. nov.**

≡ *Hieracium sparsum* subsp. *squarrosobracchiatum* O. Behr, E. Behr & Zahn, Glasnik Skopskog Naučnog Društva 20, Sect. Sci. Nat. 7: 32 (1939) [basionym].

Previously known only from Mt Korab in W North Macedonia (Behr *et al.* 1939a), this very rare plant was recently discovered on Šar Planina Mts in Serbia (Metohija) (Duraki & Niketić 2018d). Although its inclusion in *H. sect. Cernua* R. Uechtr. is not disputed, it is very different from *H. sparsum* Friv. and has to be treated as an independent microspecies. Morphologically it is much more similar to *H. naegelianiforme* (O. Behr, E. Behr &

Zahn) Buttler, and according to the author of this combination (Buttler 1991) species “only superficially resembling *H. sparsum*”. According to Behr *et al.* (1937) and Zahn (1938) possible origin of *H. naegelianiforme* was interpreted as “*sparsum* – *naegelianum*”). The similar relations can be concluded for *H. squarrosobracchiatum*. The assumption of its hybridogenous origin is further supported by recording the triploid number of chromosomes ($2n = 27$) (Ilnicki & Szeląg 2011).

M. Niketić

Hieracium trojanum* (Zahn) Niketić, *comb. et stat. nov.

≡ *Hieracium pannosum* subsp. *trojanum* O. Behr, E. Behr & Zahn in Engl., Pflanzenr. 76: 564 (1921) [basionym].

This plant is distributed in E Serbia and E and C Bulgaria (Stara Planina, Rila and Pirin Mts) (Zahn 1936). The type specimen (BRNM 7179/36) was designated from E Serbia (Mt Basara) by Szeląg (2015). According to Zahn (1936) it belongs to *H. bornmuelleri* group with numerous, more conspicuous and more dentate cauline leaves. The proposed origin of this aggregate is “*pannosum* >> *racemosum*” (Zahn 1936). Within aggregate *H. bornmuelleri* Freyn and *H. cappadocicum* Freyn represent the Anatolian species while *H. friwaldii* Rchb. f. is probably Cretan endemic. *Hieracium mokragorae* (Nägeli & Peter) Freyn from Serbia, Montenegro, Albania and Bosnia and Hercegovina, and recently described *H. crinitopannosum* Szeląg & Vladimir. from Bulgaria (Szeląg & Vladimirov 2013), could also be included in the same group.

M. Niketić

CARYOPHYLLACEAE

Dianthus lakusicii* (Wraber) Niketić, *stat. et comb. nov.

≡ *Dianthus nitidus* subsp. *lakusicii* Wraber, Biol. Vestn. 36: 97 (1988) [basionym].

This endemic SE Dinaric plant, described from Mt Bjelasica in Montenegro as *D. nitidus* subsp. *lakusicii* Wraber (1988), was initially treated as Scardo-Pindic *D. scardicus* Wettst. (Lakušić 1966). The name is currently only disputed in POWO (2020) [“Not accepted by Govaerts (2000): World Checklist of Seed Plants Database in ACCESS D: 1-30141”] and synonymized with *D. scardicus*. Our comparison of herbarium material also confirmed essential differences between this taxon and Tatra species, *D. nitidus* Waldst. & Kit., and similarity with geographically close *D. scardicus*. However, some morphological differences in relation to *D. scardicus* were also observed. Representatives of Dinaric populations have a higher stems, ± acute leaves, shorter (relatively to the stem) and wider basal leaves, more numerous cauline leaves, usually acuminate epicalyx

scales, 1–2(3) flowers in capitula and dentate petal-limb (Fig. 1). Molecular analyzes are necessary to establish definitive status and phylogenetic position of this taxon.

Specimen examined:

Montenegro, Mt Bjelasica, Lalovića dolovi above Biogradsko Lake, in pastures, 1740 m, MGRS 34T CN85, coll. M. Lovka, T. Wraber 21-Jul-1987, det. T. Wraber ISOTYPE (BEO A38766) (Fig. 1B).

Fig. 1. – A *Dianthus scardicus* Wettst. from three localities on Mts Šar Planina (BEO); B isotype of *D. nitidus* subsp. *lakusicii* Wraber [$\equiv$ *D. lakusicii* (Wraber) Niketić] (Mt Bjelasica, Lalovića Dolovi, BEO A38766).

CAPRIFOLIACEAE (DIPSACACEAE)

Knautia csikii Jáv. & Szabó, Magyar Bot. Lapok 21: 19 (1923).= *Knautia midzorensis* f. *glabrescens* Diklić in Josifović, Fl. Srbije 5: 615 (1973), **syn. nov.**

The species is distributed in Prokletije and Šar Planina Mts in the Central Balkan, as well as in Mt Ozren in SW Serbia. This name is still not accepted in the current lists (Domina 2017, POWO 2020) where it is treated

Fig. 2. – Holotype of *Knautia midzorensis* f. *glabrescens* Diklić (= *K. csikii* Jáv. & Szabó) (BEO A464).

as Alpine-Carpathian *K. longifolia* (Waldst. & Kit.) W. D. J. Koch. However, *K. csikii* clearly differs by its relatively wider and hairy leaves which makes it closer to *K. dinarica* (Murb.) Borbás. Their phylogenetic connection was also shown in Frajman *et al.* (2016), with a significant distance from *K. longifolia* group. Allopatric and vicarious *K. magnifica* Boiss. & Orph. (= *K. midzorensis* Formánek) occurs in the eastern and southern parts of the Central Balkans. However, Diklić (1973) described *Knautia midzorensis* f. *glabrescens* Diklić within the range of *K. csikii* (Mt Mokra Planina which is a part of Prokletije Mts). The analysis of the original material (BEO A464) showed that it in fact belongs to *K. csikii*.

Specimen examined:

Metohija, Prokletije Mts, Mt Mokra Planina, Oklačka Glava (Razvrš), MGRS 34T EN98, coll. M. Lovka, coll. I. Rudski 9-Jul-1932, det. N. Diklić HOLOTYPE (BEO A464) (Fig. 2).

M. Niketić

EUPHORBIACEAE

Euphorbia maglicensis Rohlena, Sitzungsber. Königl. Böhm. Ges. Wiss., Math.-Naturwiss. Cl. 1912(1): 111 (1913) ["Rasse der *E. esula* L. (= *E. maglicensis* m.)"].

≡ *Euphorbia esula* var. *maglicensis* (Rohlena) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(1): 132 (1924).

IND. LOC.: "Auf dem Maglicgebirge, unter Gerölle". LECTOTYPE (**here designated** by D. Geltman, Oct-2000): [Montenegro] "in lapidosis mtis Maglić", Aug-1906, *J. Rohlena* (sub "*E. esula* b) *magličensis* m.") (PRC barcode 452975 [photo!] <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.prc452975>); isolectotype: Aug-1906, *J. Rohlena* (58 BM barcode 000751501).

= *Euphorbia panicii* Beck, Glasn. Zemaljsk. Muz. Bosni Hercegovini 32: 95 (1920) ["*Pančićii*"], **syn. nov.** ≡ *Tithymalus panicii* (Beck) Soják, Čas. Nár. Mus., Odd. Přír. 140: 175 (1972).

IND. LOC.: "Na vapnenastim stijenama u dolini Rzave, kod Dobruna, u Razdolini kod 400 do 450 m, blizu Strmice na Limu (K. Maly kao *E. variabilis*)". LECTOTYPE (**here designated** by A. R. Smith, 08-Oct-1964): [Bosnia] "An felsen nächst Strmica am Lim, Kalk felsen und Shutt; ca. 450 m", 12-Apr-1910, *K. Malý* (sub "? *E. variabilis*") (K barcode 000911842 [photo!] <https://plants.jstor.org/stable/viewer/10.5555/al.ap.specimen.k000911842?loggedin=true>).

- “*Euphorbia variabilis*” sensu Pančić, Fl. Serbiae: 612 (1874) [non Ces., Bibl. Ital. (Milan) 91: 348 (1838)]; K. Malý, Magyar Bot. Lapok 8(1–4): 93.

The first indication that this is a new taxon (“*E. species proxima virgatae sed diversa*”) was given by Pančić (1867) for W Serbia (vicinity of Mokra Gora). The same data was represented in the *Flora of the Principality of Serbia* (Pančić 1874) as *E. variabilis* Ces. (currently known only from the Alps). The first description and diagnosis of the taxon were presented by Malý (1909) who found the same plant in E Bosnia (Dobrun) near the mentioned Pančić’s site in W Serbia. He expressed suspicion that the plant belongs to *E. variabilis*, but did not formally describe a new taxon. He was followed by Beck (1920), who described *E. pancicii* based on Pančić’s and Malý’s material. Under that name, the plant was recognized in *Prodromus Florae peninsulae Balcanicae* (Hayek 1924) and *Flora of SR Serbia* (Janković & Nikolić 1972), and it was also mentioned for the flora of Montenegro (Blečić 1953). However, in the *Flora Europaea* (Smith & Tutin 1968) species was wrongly treated as a synonym of *E. esula* L. subsp. *esula*. Although this calciphyllous East Illyrian plant really belongs to a very variable complex of the widespread Eurasian *E. esula*, it differs in many morphological and ecological characteristics, that were thoroughly presented by Wraber (1982) and accepted in Niketić (2014). In addition, it differs phenologically by flowering already in April-May in the lower altitudes. Nevertheless, in the current lists (e.g. Govaerts *et al.* 2020) the taxon is still treated as a variety of *E. esula*.

The same plant was independently described by Rohlena (1913) from NW Montenegro on the border with Bosnia and Herzegovina (Mt Maglić) under the name *E. maglicensis*. Based on the following citation in the protologue: “Rasse der *E. esula* L. (= *E. maglicensis* m.)”, it remains unclear whether the author described the species or the variety, especially if we keep in mind that on the original herbarium label (PRC, BM), as well as in Rohlena (1942), the taxon was treated as a variety. However, it is more probable that the author initially used the term “race” in the sense of microspecies, not as an indication of rank. Hayek (1924), in addition to *E. pancicii*, also mentioned this plant as *E. esula* var. *maglicensis* (Rohlena) Hayek. Inspection of the original material of *E. pancicii* (K) and *E. maglicensis* (PRC, BM) showed that they are conspecific, but clearly different from *E. esula*. Although the name *E. maglicensis* has long been neglected, it has priority over *E. pancicii* (Art. 11).

Floristic notes

New and confirmed taxa for the flora of Serbia

LILIOPSIDA

LILIACEAE (XANTHORRHOEACEAE)

Asphodeline taurica (Pall. ex M. Bieb) Endl., Cat. Horti Vindob. 1: 142 (1842).

Two species of the genus *Asphodeline* Rchb. are reported for the flora of Serbia: *A. liburnica* (Scop.) Rchb. and *A. lutea* (L.) Rchb. (Anačkov *et*

Fig. 3. – *Asphodeline taurica* (Pall. ex M. Bieb) Endl. (E Serbia, Mt Suva Planina, BEO 100027).

al. 2018). During the field trip in Mt Suva Planina in E Serbia two individuals of East-Mediterranean *A. taurica* (Pall. ex M. Bieb) Endl. was found on the edge of a mountain ridge. In the Balkan Peninsula it is known from Bulgaria, Greece and Albania (Govaerts 2020) as well as from North Macedonia (Matveeva 1968). Notable diagnostic characters for this plant are white petals and conspicuous and dense scarious whitish bracts. Unfortunately, the species was not found in the same micro-locality last year. In the wider area of Mt Suva Planina and the surroundings of Niš, within a radius of 12 km, all three species listed for the flora of Serbia were reported. *Asphodeline liburnica* are reported for the Jelašnica Gorge (Petrović 1882) and also was recently found in Koritnik hill in the foothill of the Mt Suva Planina (coll. M. Niketić BEO). *Asphodeline lutea* was mentioned by Pančić (1884) for Sićevačka Gorge (Vis peak), but after that it was no longer found at that locality, but only *A. liburnica* previously reported by Petrović (1882). The original Pančić's material of *A. lutea* was not found in BEOU. Niketić (2010) preliminary announced its alleged presence in Mt Suva Planina, but this record actually correspond to *A. taurica*.

First record:

E Serbia, Suva Planina, Sokolov Kamen peak, limestone, rocky ground along the ridge, 1350 m, MGRS 34T EN98, coll. M. Niketić 04-Jul-1989, det. M. Niketić (BEO 100027) (Figs 3, 16).

A new species for the flora of Serbia. Probably extinct.

M. Niketić

POACEAE (GRAMINEAE)

Poa stiriaca Dörf., Jahreskat. Wiener Bot. Tauschvereins 1904: 226 (1904).

This perennial grass is the only representative of *P.* sect. *Leptophyllae* J. R. Edmondson in *Flora Europaea* and its distribution encompasses E and C Europe (Austria, Czech Republic, Poland and Romania). In the Balkan Peninsula it is recorded for Montenegro (Rohlena 1942, Edmondson 1980), Bosnia and Herzegovina (Stefanović 1960), SE Albania (De Sanctis *et al.* 2018) and NW Greece (Dimitrov & Vutov 2015). Previous unreferenced and unverified data of its presence in Serbia (Svoboda 2012) was not supported either by herbarium specimens, nor by specific data for a certain locality, and it was disputed by Lakušić *et al.* (2018) and Niketić *et al.* (2018b). The newly-discovered locality of *P. stiriaca* in Mt Suvobor represents the most continental finding of the species in the Balkan Peninsula. In all three localities in Serbia (Fig. 16) this species grows in ultramaphitic geological substrate, in *Erico-Quercetum dalechampii* forest

or different types of *Pinus nigra* J. F. Arnold forests at the elevation between 700 and 1520 m a.s.l. Within its range, *P. stiriaca* was found on various types of bedrock, also including ultramaphitic substrate in Austria (Eggler 1955) and Greece (Dimitrov & Vutov 2015).

Fig. 4. – *Poa stiriaca* Dörf. (SW Serbia, Mt Ozren BEO, 100009).

First records:

SW Serbia, Mt Ozren near Sjenica, Čir, ultramaphite, roadside in the *Pinus nigra* forest, 1520 m, MGRS 34T DN08, 19.8050839 E, 43.2448787 N, coll. M. Niketić, G. Tomović 30-Jul-2020, det. M. Niketić (BEO 100009) (Fig. 4).

NW Serbia, Mt Maljen, Divčibare, Crni Vrh peak, Kičer, ultramaphite, *Brachypodio-Pinetum nigrae*, 1042 m, MGRS 34T DP18, 19.99488 E, 44.09031 N, coll. D. Lakušić 27-Sep-2020, det. D. Lakušić, (BEOU 50851).

Fig. 5. – *Poa stiriaca* Dörf. (NW Serbia, Mt Maljen) (photo D. Lakušić).

NW Serbia, Mt Maljen, Divčibare, Crni Vrh peak, Kičer, ultramaphite, *Brachypodio-Pinetum nigrae*, 1010 m, MGRS 34T DP18, 19.99347 E, 44.09083 N, D. Lakušić 27-Sep-2020 (field observ.).

NW Serbia, Mt Maljen, Tometino Polje, Rastovac, ultramaphite, *Erico-Pinetum nigrae*, 818 m, MGRS 34T DP18, 19.99781 E, 44.07844 N, D. Lakušić 27-Sep-2020 (field observ.).

NW Serbia, Mt Maljen, Tometino Polje, Rastovac, ultramaphite, *Pinetum nigrae*, 806 m, MGRS 34T DP18, 19.99755 E, 44.07894 N, D. Lakušić 27-Sep-2020 (field observ.).

NW Serbia, Mt Suvobor, Nateme, ultramaphite, *Erico-Quercetum dalechampii* forest, 688 m, MGRS 34T DP38, 20.168641 E, 44.134539 N, coll. N. Kuzmanović, I. Stevanoski, 08-Oct-2019, det. N. Kuzmanović (BEOU 63632) (Fig. 5).

Confirmed and for the first time located species for the flora of Serbia.

M. Niketić, G. Tomović, D. Lakušić, N. Kuzmanović, I. Stevanoski

MAGNOLIOPSIDA

APIACEAE (UMBELLIFERAE)

Sison amomum L., Sp. Pl.: 252 (1753).

In the flora of Europe and the Mediterranean there are three species of the genus *Sison* L. (Tutin 1968, Hand 2011): *S. amomum* L., *S. exaltatum* Boiss. and *S. segetum* L. of which *S. exaltatum* is present in E Mediterranean (Israel, Lebanon and Syria), *S. segetum* is distributed in the W Europe, while *S. amomum* has the widest distribution range that encompasses W and S Europe, Asia Minor and N Africa (Maslo *et al.* 2020 and literature therein). In the Balkan Peninsula this species is present in Albania (Barina *et al.* 2011), Bulgaria, Croatia, Greece, Romania and Turkey in Europe (Hand 2011) and only recently it was recorded as a new species for the flora of Bosnia and Herzegovina (Maslo *et al.* 2020).

Sison amomum was mentioned for the first time for the flora of Serbia by Stojanović & Stevanović (2008) who presented the list of vascular plant taxa for Mt Gučevo in NW Serbia. Although this was the first mention both of the genus *Sison* and the species *S. amomum* in the flora of Serbia, the authors did not point this out in their paper, and they did not provide any details about the exact locality in which this species was collected.

By checking the herbarium material of *S. amomum* in the Herbarium of the Institute of Botany and Botanical Garden “Jevremovac”, University of Belgrade (50835 BEOU) it was confirmed that this specimens was correctly identified. By providing additional detail about exact locality, habitat and the date of collecting as well as picture of the herbarium specimen (Fig. 6) we confirm and point out the existence both of the new genus as well as new species for the flora of Serbia. Detail description of

the genus as well as the species could be found in Tutin (1968) and Maslo *et al.* (2020).

Fig. 6. – *Sison amomum* L. (NW Serbia, Mt Gučevo, 50835 BEOU).

Finding the species *S. amomum* on Mt Gučevo (locality Velika Ravan) is not so surprising considering that the nearest newly recorded localities in central and eastern Bosnia (Gornji Srebrenik near Tuzla and Hrvati near

Banovići) are only c. 60 km to the west. According to Maslo *et al.* (2020) this submediterranean-subatlantic plant inhabits the edges of the local roads near Tuzla (the villages Gornji Srebrenik and Hrvati) et the elevations between 360 and 420 m. On Mt Gučevo (locality Velika Ravan) the species was found next to a macadam road, near the wooden fence of a rural abandoned house where water is retained, et elevation c. 530 m.

First record:

NW Serbia, Mt Gučevo, on the road to Gučevo, Velika Ravan, MGRS 34T CQ52, coll. V. Mitrović, 19-Aug-1999, det. V. Mitrović (50835 BEOU) (Figs 6, 16).

Confirmed and for the first time located genus and species for the flora of Serbia.

V. Stojanović, V. Stevanović

ASTERACEAE (COMPOSITAE)

Artemisia dracunculus L., Sp. Pl.: 849 (1753).

Artemisia dracunculus belongs to the subgenus *Dracunculus* Besser, series *Dracunculus* (Poljakov 1961). The species is characterized by a wide but disjunctive Holarctic distribution. Most of the area covers the temperate and boreal areas of Eurasia from the European part of Russia in the west to Manchuria and NE China in the east (Meusel & Jäger 1992). The northern part of the area in Eurasia includes the northeastern part of the boreal zone of Yakutia, and the southern mountainous areas of the Turanian region. The other part of the range includes temperate areas of western and central North America. European part of its natural range includes Belarus, Ukraine, Crimea and Russia to the Urals. Due to the spice tarragon, *A. dracunculus* is grown throughout the Holarctic, as well as in Europe where in many countries such as Poland, Romania, Austria, Switzerland and France the species is naturalized (subspontaneous) (Greuter 2006+).

Its status in the countries where it is still grown (Norway, Belgium, Germany, Latvia, Estonia) is naturalized in some places, while Serbia is listed as a country where it is only grown. Data on tarragon cultivation in Serbia are scarce and localized. It is stated as cultivated for Deliblato Sands, where after the Second World War there were trial areas where the possibilities of its cultivation were explored. According to Šajinović & Mihajlov (1979), it was introduced in the period between 1960 and 1965 in the localities of Devojački Bunar and Dubovac, and after only ten years it spreads outside the plantations. According to the same authors, experimental studies of the introduction of about 400 medicinal plants have shown that *A. dracunculus* is one of the four species that are best adapted to the climate.

Fig. 7. – *Artemisia dracunculus* L. (Banat, Deliblatska Sands, 11038 BEOU).

Almost 30 years later, this species was found along the Sušara – Korn road, c. 1 km from the place on the Korn Bunar (BEOU). The population of *A. dracunculus* in this locality is small (only a dozen specimens were found) and is located on a steamed surface of *Robinia pseudoacacia* L. forest surrounded by shrubs *Crataegus monogyna* Jasq., *Ligustrum vulgare* L. and *Euonymus europaeus* L. It can be assumed that this small population

originates from nearby plantations of medicinal herbs, most likely from Korn or Sušara. It is possible that this species could be also found in other parts of the Deliblato Sands.

First record:

Banat, Deliblatska Sands, Korn Bunar, steppe and the edge of a planted *Robinia pseudoacacia* forest, sand, MGRS 34T EQ07, coll. V. Stevanović, 16-Aug-1998, det. V. Stevanović (11038 BEOU) (Fig. 7).

Confirmed and for the first time located allochthonous species for the flora of Serbia.

V. Stevanović, M. Niketić, G. Tomović

Centaurea austrobalcanica subsp. ***prisadana*** (Skokanová) Raus & Strid, Phytol. Balcan. 26(3): 505 (2020).

≡ *Cyanus austrobalcanicus* Skokanová, Pl. Syst. Evol. 305: 590 (2019) [basionym].

- “*Centaurea nissana*” sensu Randelović & Stamenković, Leskovački zbornik 24: 377 (1984) [non Petrović, Addit. Fl. Agr. Nyss.: 110 (1885)].
- “*Centaurea napulifera* subsp. *tuberosa*” sensu Zlatković, V. Randjelović & N. Randjelović, III Simp. fl. jugoist. Srbije: 104 (1993) [non (Vis.) Stoj. & Acht., Stud. Centaur. Bulg.: 22 (1935)].

Randelović & Stamenković (1984) were the first botanists who found this Balkan endemic plant in Mt Rujan and near Preševo in S Serbia. However, they published this record under the name *C. nissana* Petrović. This data was also cited by Randelović & Stamenković (1985) (sub *C. nissana* Petrović) and Zlatković *et al.* (1993) (sub *C. napulifera* subsp. *tuberosa*) from the same locality. According to Greuter (2006+) *Cyanus tuberosus* (Vis.) Soják (= *Centaurea tuberosa* Vis.) is distributed in Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Greece, Montenegro, North Macedonia and Serbia.

However, according to the most recent genetic, morphological, ecological and phenological study of Skokanová *et al.* (2019) the *C. tuberosus* group is extensively radiated and comprises nine almost allopatric genetic lineages within eight Balkan endemic species. One of them is *Cyanus austrobalcanicus* Skokanová which comprises two subspecies: *C. a.* subsp. *austrobalcanicus* distributed in N Greece and S Bulgaria and *C. a.* subsp. *prisadanus* Skokanová present in North Macedonia and possibly in NC Greece. Although Skokanová *et al.* (2019) did not mention that the last

Fig. 8. – *Centaurea austrobalcanica* subsp. *prisadana* (Skokanová) Raus & Strid (S Serbia, Mt Rujan Planina, BEO 100019).

subspecies grows also in Serbia, the close proximity of the North Macedonian populations (e.g. Mt Vodno), as well as comparison of the herbarium specimens from S Serbia with diagnosis presented in the paper for subsp. *prisadanus*, confirms its presence in Mt Rujan.

Most recently Raus & Strid (2020) published new combination in some Balkan *Centaurea* taxa. One of them is *Centaurea austrobalcanica* subsp. *prisadana* (Skokanová) Raus & Strid and their attitude is accepted in this paper.

First records:

S Serbia, Mt Rujan Planina, Cer village, solo silicicolo, in pascuis, 500 m, MGRS 34T EM57, coll. M. Niketić 14-May-1991, det. S. Bancheva (sub *C. tuberosa*), rev. M. Niketić 01-Apr-2020 (BEO 100019) (Figs 8, 16).

S Serbia, Mt Rujan Planina, Orljak, 742 m, limestone, stony pastures, coll. & det. M. Niketić, G. Tomović 03-May-2009 (sub *C. tuberosa*) (29576 BEOU).

A new species and subspecies for the flora of Serbia.

M. Niketić, G. Tomović

Hieracium pseudofastigiatum Degen & Zahn, Magyar Bot. Lapok 5: 68 (1906).

≡ *Hieracium umbrosum* subsp. *pseudofastigiatum* (Degen & Zahn) Zahn in Engl., Pflanzenr. 76: 794 (1921).

Hieracium pseudofastigiatum belongs to *H. umbrosum* group (*H. murorum* – *prenanthoides*) together with *H. eugraptum* Omang, *H. umbrosum* Jordan, *H. vipetinum* Huter ex Freyn (Sell & West 1976) and 16 additional taxa (Greuter 2006+). It is a Balkan subendemic species previously known from Albania, Bulgaria, North Macedonia and Romania (Greuter 2006+). During floristic exploration of the Đerdap Gorge in NE Serbia it was also found on locality Ploča, near the road in the zone of *Quercus cerris* L. and *Q. frainetto* Ten. It was previously also recorded from Mt Domogled in Romania (Zahn 1937), 33 km northeast from Serbian population.

First record:

NE Serbia, Đerdap Gorge, Ploča, limestone, asphalt road, MGRS 34T FQ04, coll. S. Vukojičić, G. Tomović, “ENDEMIT” 15-Jul-2001, det. M. Niketić (13374 BEOU) (Fig. 9).

A new species for the flora of Serbia.

Fig. 9. – *Hieracium pseudofastigiatum* Degen & Zahn (NE Serbia, Đerdap Gorge, 13374 BEOU).

M. Niketić, G. Tomović, S. Vukojičić

Leucanthemum illyricum (Horvatić) Vogt & Greuter, Willdenowia 33: 41 (2003).

[Papeš, Acta Bot. Croat. 32: 244 (1973), comb. inval.]

- ≡ *Chrysanthemum croaticum* var. *illyricum* Horvatić, Acta Bot. Inst. Bot. Univ. Zagreb. 3: 89 (1928) [basionym]; *Leucanthemum atratum* subsp. *illyricum* (Horvatić) Soó, Acta Bot. Acad. Sci. Hung. 23: 386 (1978).
- “*Chrysanthemum chloroticum*” sensu K. Malý, Glasn. Zem. Muz. Saraj. 46: 53 (1934) [non (A. Kern. & Murb.) Horvatić, Acta Bot. Inst. Bot. Univ. Zagreb 3: 102 (1928)].
- “*Chrysanthemum croaticum*” sensu Vukićević, Glasnik Šumarskog fakulteta 29: tab. 10 (1965) [non Horvatić, Acta Bot. Inst. Bot. Univ. Zagreb 3: 89 (1928)].

This Balkan endemic species inhabits rocky grasslands and rock crevices in subalpine belt and gorges in C and E Dinarides (Horvatić 1928, Papeš 1973). It was very often confused with parapatric Dinaric species, *L. visianii* (Gjurašin) Vogt & Greuter [= *L. croaticum* (Horvatić) Bjelčić] and *L. chloroticum* Kern. & Murb. ex Murb., that also have ± fleshy glabrescent leaves. *Leucanthemum illyricum* is characterized by more robust growth, coarsely dentate leaves and larger capitula. In the past period, this species was also found in Mt Tara in W Serbia in many places in the rocky habitats of *Ostrya carpinifolia* Scop. and *Picea omorika* (Pančić) Purk. communities (BEO 100016). Previous record of *L. chloroticum* (Malý 1934) [“*Chrysanthemum chloroticum*”] from the same mountain undoubtedly corresponds to *L. illyricum*. Through the revision of herbarium material, its presence in the area of Prokletije Mts in Metohija was confirmed (BEO 25584) (Fig. 16). Previous report of *L. visianii* (Vukićević 1965) from the Serbian part of Prokletije Mts probably also refers to *L. illyricum*.

First record:

W Serbia, Mt Tara, Kozje Stene, limestone, 800-900 m, MGRS 34T CP76, coll. M. Niketić 3-Jul-1988, det. M. Niketić (BEO 100016) (Figs 10, 16).

Metohija, Prokletije Mts, monast. Dečani, in the bush near Dečanska Bistrica river, MGRS 34T DN31, coll. P. Černjavski, I. Rudski, V. Lindtner 14-Jul-1933, det. P. Černjavski (sub *L. vulgare*), rev. M. Niketić 16-Mar-2020 (BEO 25584).

A new species for the flora of Serbia.

Fig. 10. – *Leucanthemum illyricum* (Horvatić) Vogt & Greuter (W Serbia, Mt Tara, BEO 100016).

M. Niketić

***Rudbeckia hirta* L., Sp. Pl.: 907 (1753).**

This species is native to North America and widely cultivated for ornamental and naturalized in waste places, river banks and woods mainly in C Europe (Hansen 1976). Concerning neighboring countries it has

naturalized status in Hungary and unknown status in Romania [Greuter (2006+)]. In Italy it has status of naturalized alien (NAT) (Galasso *et al.* 2018) while in Croatia it is treated as allochthonous species out of cultivation (Nikolić 2020). According to criteria of Niketić & Tomović (2018) its invasive status in Serbia is considered to be: A(A) (initial phase of naturalization).

First record:

NE Serbia, Kučevo, Vrelo, Marica, gravelly substrate, roadside, 320 m, MGRS 34T EQ52, 21.6961 E, 44.51587 N, coll. M. Niketić, G. Tomović 13-Jun-2020, det. M. Niketić (BEO 100028) (Figs 11, 16).

A new allochthonous species for the flora of Serbia.

Fig. 11. – *Rudbeckia hirta* L. (NE Serbia, Kučevo, Vrelo) (photo M. Niketić).

BORAGINACEAE

Brunnera macrophylla (Adams) I. M. Johnston, Contr. Gray Herb. 73: 54 (1924).

≡ *Myosotis macrophylla* Adams, Beitr. Naturk. 1: 46 (1805) [basionym].

It is a single representative of the genus *Brunnera* Steven in the flora of Europe native to the Transcaucasia (Azerbaijan, Armenia and Georgia) and the Caucasus (Turkey). *Brunnera macrophylla* is introduced in Great Britain,

Fig. 12. – *Brunnera macrophylla* (Adams) I. M. Johnston (Šumadija, Beograd, Zvezdara) (photo M. Niketić).

Germany, Switzerland, Austria, Slovakia and Czech Republic (Chater 1972, Valdés 2011), while in Italy it has status of naturalized alien (NAT) (Galasso *et al.* 2018). To our knowledge, this species has not been found in neighboring countries so far. In the urban part of Belgrade it was found in the park-forest Zvezdara in the zone of *Quercus robur* L., but with domination of some invasive and cultivated trees. Although it was found to have escaped from cultivation, it spread to the forest ecosystem. Its invasive status in Serbia according to criteria of Niketić & Tomović (2018) is: A(A)c (initial phase of naturalization, also cultivated).

First record:

Šumadija, Beograd, Zvezdara, park-forest Zvezdara, MGRS 34T DQ66, 20.50238 E, 44.81161 N, coll. M. Niketić 07-Apr-2020, det. M. Niketić (BEO 100017) (Figs 12, 16).

A new allochthonous species for the flora of Serbia, escaped from cultivation.

M. Niketić, G. Tomović

BRASSICACEAE (CRUCIFERAE)

Alyssum wierzbickii Heuff., Flora 18: 242 (1835).

Species was described from SW Romania and it is also distributed in E Serbia and W Bulgaria (Španiel *et al.* 2017). For the flora of Serbia it was reported for Mt Ozren (Leskovik) near Sokobanja, Mt Stol near Bor and Mt Rtanj (Pančić 1874, Formanek 1896, Diklić 1972, Nikolić *et al.* 1986). So far, only material from Mt Ozren has been detected (coll. J. Pančić BP, W) (Španiel *et al.* 2017). Its presence was confirmed on the border with Bulgaria during field research as well as in herbarium material.

New records:

E Serbia, Stara Planina, Toplodolska river - right bank, beach forest, MGRS 34T FN39, FP30, coll. V. Nikolić, N. Diklić 25-Jun-1958, det. M. Niketić (BEO 42200).

E Serbia, Mt Ruj Planina, from Štrbi Kamik to Zdravči Kamik, rocky grounds and high-mountain pastures, limestone, 1273 m, MGRS 34T FN24, coll. M. Niketić, G. Tomović 30-Jun-2016, det. M. Niketić (BEO 100023) (Figs 13, 16).

E Serbia, Mt Ruj Planina, Štrbi Kamik peak, rocky grounds and high-mountain pastures and herb vegetation, limestone, 1480 m, rudine, MGRS 34T FN24, coll. M. Niketić, G. Tomović 30-Jun-2016, det. M. Niketić (BEO 100024).

Confirmed species for the flora of Serbia.

Fig. 13. – *Alyssum wierzbickii* Heuff. (Mt Ruj Planina, from Štrbi Kamik to Zdravči Kamik, BEO 100023).

M. Niketić, G. Tomović

CARYOPHYLLACEAE

Silene chalcedonica (L.) E. H. L. Krause in Sturm, *Deutschl. Fl.* (Sturm), ed. 2, 5: 96 (1901).

≡ *Lychnis chalcedonica* L., *Sp. Pl.*: 436 (1753).

The species was originally described and treated within the genus *Lychnis* L., which has recently been included in the genus *Silene* (Jafari *et al.* 2020). It is native for Central and Eastern Asia and widely cultivated in gardens elsewhere. According to Marhold (2011) it is introduced, as escaped from cultivation, in Finland, Italy, Moldova and Ukraine. There are two historic specimens (1922) in BEO (5261, 5262) from NE Serbia and Pomoravlje, but it is not entirely clear from the labels whether the plant was domesticated or cultivated. During the pre-congress excursion of the XVII International Botanical Congress (Vienna 2005) a locally abundant population of *S. chalcedonica* was discovered along the edge of the indigenous oak forest (*Quercetum confertae-cerridis* Rudski) in Mt Rtanj.

Fig. 14. – *Silene chalcedonica* (L.) E. H. L. Krause (E Serbia, Mt Rtanj, between villages Rtanj and Nikolinac, BEO 100031) (photo O. Vasić).

First records:

Pomoravlje, Svilajnac and the whole of Resava, MGRS 34T EP19, coll. M. S. Dimitrijević, det. Ž. Jurišić (sub *Lychnis chalcedonica*) (BEO 5261).

NE Serbia, Bor, Trnjane, MGRS 34T FP09, coll. P. S. Pavlović Jun-1922, det. Ž. Jurišić (sub *Lychnis chalcedonica*) (BEO 5262).

E Serbia, Mt Rtanj, between villages Rtanj and Nikolinac, limestone, the edge of an oak forest, MGRS 34T EP74, coll. O. Vasić 15-Jul-2005, det. M. Niketić (BEO 100031) (Figs 14, 34).

A new allochthonous species for the flora of Serbia, escaped from cultivation.

O. Vasić, M. Niketić

***Dianthus lakusicii* (Wraber) Niketić**

≡ *Dianthus nitidus* subsp. *lakusicii* Wraber, Biol. Vestn. 36: 97 (1988) [basionym].

This orophytic plant, closely related to *D. scardicus* Wettst., is endemic for SE Dinarides (see on page 97). It was preliminary enounced by Niketić (2000) for the part of Prokletije Mts in Metohija, near the border with

Fig. 15. – *Dianthus lakusicii* (Wraber) Niketić (Metohija, Prokletije Mts, Mt Hajla, BEO 100039).

Fig. 16. – Distribution of some new taxa for the vascular flora of Serbia as well as *Erechites hieraciifolius*, new for Vojvodina (small circles within symbol: white – new records; black – literature record).

Montenegro (Mt Hajla). After that it was proposed for protection under *The Rulebook on the proclamation and protection of strictly protected and protected wild species of plants, animals and fungi* (Sl. glasnik RS 2010–

2011) and the presence of this taxon was reported for Serbia and Kosovo and Metohija in several publications (Ferakova *et al.* 2011, Amidžić *et al.* 2013, Tomović *et al.* 2014, Vuksanović 2016, Gavrilović *et al.* 2017), but without a specific locality, citation of data sources or any other information. Therefore, all of the above reports can be considered as preliminary. Area of occupancy was estimated at about 1 km² and subpopulation size at more than 5000 mature individuals. It was found in subalpine belt in carbonate rocky pastures and in the zone of *Pinus mugo* Turra.

First record:

Metohija, Mts Prokletije, Mt Hajla, 1800-2400 m, limestone, rocky pastures, MGRS 34T DN23, coll. M. Niketić 03-Sep-1997, det. M. Niketić (BEO 100039) (Figs 15–16).

A new species for the flora of Serbia.

M. Niketić

EUPHORBIACEAE

Euphorbia glareosa Pall. ex M. Bieb., Fl. Taur.-Caucas. 1: 373 (1808).

≡ *Euphorbia nicaeensis* subsp. *glareosa* (Pall. ex M. Bieb.) Radcl.-Sm., Rept. Spec. Nov. Regni Veg. 79: 55 (1968).

During floristic investigation in Mt Rujan Planina in S Serbia an interesting plant from *E. nicaeensis* group was found on rocky and eroded areas around the quarry (BEO 100005) (Fig. 17A–B). After revision of herbarium material from Serbia and neighbouring countries, it was identified as *E. glareosa* Pall. ex M. Bieb. Although the same taxon has already been reported in the *Flora of SR Serbia* (Janković & Nikolić 1972) and many other sources, everything previously mentioned for this taxon actually refers to *E. pannonica* Host (= *E. glareosa* var. *lasiocarpa* Boiss.) (Fig. 17C), which is much larger plant, with stiff and thicker, relatively wider leaves, with 3–5(7) prominent basal nerves and larger number of rays. According to Geltman (2015, 2020) and Pahlevani (2017) *E. pannonica* is a Pannonian-East Submediterranean plant (described from Pannonian Austria) which grows most often in steppe or steppe-like grasslands on different substrates, while *E. glareosa* is an Oriental-Pontian-East Submediterranean plant (described from Georgia and absent from the Pannonian Basin) and more associated with rocky substrate. The occurrence of both species in sympatry was observed in E Mediterranean (Transcaucasia, North Anatolia and E part of the Balkan Peninsula). In the current *World Checklist* (Govaerts *et al.* 2020) *E. pannonica* is incorrectly not listed for the Pannonian countries, including Serbia. According to the same source *E. glareosa* is also absent from that region and reaches out to the west to Bulgaria, while according to Geltman (2015) it also reaches to Dobruja in E Romania. The new locality of the species on Mt Rujan in S

Serbia is currently the westernmost point in the species range. Its presence can also be expected in Kosovo and Metohija, as well as in North Macedonia. Since all previous data on the presence in Serbia were erroneous and correspond to *E. pannonica*, the new record from Mt Rujan Planina at the same time represents a new addition to the flora of Serbia.

Fig. 17. – A-B *Euphorbia glareaosa* Pall. ex M. Bieb. (S Serbia, Mt Rujan Planina BEO 100005); C *E. pannonica* (Vojvodina, Srem, Banstol, BEO 37263).

First record:

S Serbia, Mt Rujan Planina, Cer, quarry, limestone, eroded areas around the quarry, MGRS 34T EM57, coll. B. Zlatković 04-Jun-2020, det. B. Zlatković, M. Niketić (BEO 100005) (Figs 17A, 34).

A new species for the flora of Serbia.

B. Zlatković, M. Niketić

LAMIACEAE (LABIATAE)

Stachys recta* subsp. *nitens (Janka) Ciocârlan, Fl. Il. României: 180 (1990).

[Beldie, Fl. Rom. Determ. Ilust. Pl. Vasc.2: 75 (1979), comb. inval.]

= *Stachys nitens* Janka, Linnaea 30: 597 (1860) [basionym] ≡ *Stachys recta* var. *nitens* (Janka) Chrtek f., Folia Geobot. Phytotax. 27: 182 (1992).

= *Stachys recta* var. *ramosissima* Rochel ex Griseb. & Schenk, Arch. Naturgesch. 18(1): 330 (1852); Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(2): 294 (1929) [subsp. *subcrenata*], non sensu Diklić, Fl. Srbije 6: 427 (1974).

Fig. 18. – *Stachys recta* subsp. *nitens* (Janka) Ciocârlan (NE Serbia, Mt Veliki Greben, Gradašnica cave – Avramica, BEO 100037).

Independently described as a variety (Grisebach & Schenk 1852) and a species (Janka 1860) from SW Romania, near the Serbian border, the plant was also mentioned for the eastern parts of Romania, near the Black Sea coast (Răvăruț 1960, Beldie 1979). In our opinion the populations of this plant are morphologically and ecologically well defined and deserve the rank of a subspecies. It differs from the type subspecies by its glabrescent habitus, thin and fragile branches, narrow leaves, as well as by longer calyx teeth. For the flora of the Balkan Peninsula it was also recorded in Serbia and Bosnia and Hercegovina (Hayek 1929) (as *S. recta* var. *ramosissima*) without specific localities, but since this author emphasizes that the stems and calyx are hairy, these data probably do not refer to *S. recta* subsp. *nitens*. It was later mentioned under the same name for S Serbia (Diklić 1974), but this record has to be treated as erroneous. In the same publication Diklić also mentioned possibly correct location of this taxon (Mt Miroč in NW Serbia), but under the misapplied name “*S. recta* var. *fragilis*”. Niketić (2005) preliminarily recorded this subspecies (as *S. nitens*) for NW (Đerdap Gorge and Podvrška: Ms Velika Čuka) and E (Mt Suva Planina and Jerma Gorge) Serbia. However, the specimens from E Serbia do not belong to this subspecies. Since all previous data for Serbia were erroneous or preliminary, the taxon should be considered as new for the flora. Its correct distribution in Serbia is shown in Fig. 34.

First records:

NE Serbia, Đerdap Gorge, between Tekija and Kovilovski stream, MGRS 34T FQ14, coll. V. Nikolić 07-Aug-1966, det. N. Diklić (sub *S. recta* subsp. *labiosa* var. *glabrata*), rev. M. Niketić 14-Feb-2004 (s.n. BEO).

NE Serbia, Đerdap Gorge, Tekija, peat, MGRS 34T FQ14, coll. V. Nikolić 03-Oct-1968, det. N. Diklić (sub *S. recta* subsp. *recta*), rev. M. Niketić 14-Feb-2004 (s.n. BEO).

NE Serbia, Donji Milanovac, Mt Veliki Greben, *Carpinus orientalis* underbrush, MGRS 34T EQ92, coll. N. Diklić 31-Jul-1964 (sub *S. recta* subsp. *subcrenata* var. *hyssopifolia* subvar. *fragilis*), rev. M. Niketić 01-Feb-2007 (s.n. BEO).

NE Serbia, Đerdap Gorge, Donji Milanovac, Porečka river - Velika Gradišnica river, MGRS 34T EQ92, coll. V. Nikolić, N. Diklić 04-Aug-1966, det. M. Niketić (s.n. BEO).

NE Serbia, Đerdap Gorge, Mali Štrbac peak, rocky grounds along the ridge, MGRS 34T FQ04, coll. V. Nikolić, N. Diklić 18-Jun-1977, det. M. Niketić (s.n. BEO).

Fig. 19. – *Stachys recta* subsp. *nitens* (Janka) Ciocârlan (NE Serbia, Mt Veliki Greben, Gradašnica cave) (photo M. Niketić).

NE Srbija: Mt Miroč, Podvrška, Ms Velika Čuka; 370 m; limestone; rock crevices. MGRS 34T 34T EQ92FQ13, coll. M. Niketić, V. Stevanović, G. Tomović 15-Jun-1998, det. M. Niketić (100040 BEO).

NE Serbia: Mt Veliki Greben, Gradašnica cave - Avramica; 350 m; limestone; rock crevices. MGRS 34T 34T EQ92, coll. M. Niketić, G. Tomović 14-Jun-2009, det. M. Niketić (BEO 100037) (Figs 18–19).

NE Serbia: Mt Veliki Greben, Gradašnica cave; 350 m; limestone; rock crevices. MGRS 34T 34T EQ92, coll. M. Niketić 26-Jul-2010, det. M. Niketić (BEO 100038).

NE Serbia: Đerdap Gorge, Tekija, Misija; 150 m; serpentinite; steppe vegetation. MGRS 34T 34T FQ14, coll. M. Niketić 18-Sep-2018, det. M. Niketić (BEO 100034).

NE Serbia: Đerdap Gorge, Tekija, Džervin Kamen; 300 m; limestone; rock crevices. MGRS 34T 34T FQ15, coll. M. Niketić 17-Sep-2018, det. M. Niketić (BEO 100035).

NE Serbia: Negotinska Krajina, Vratna Gorge; 250 m; limestone; rock crevices. MGRS 34T 34T FQ01, coll. M. Niketić 14-Nov-2018, det. M. Niketić (BEO 100036).

A new subspecies for the flora of Serbia.

M. Niketić

Stachys recta subsp. *rhodopaea* (Velen.) Chrtek f., Folia Geobot. Phytotax. 27: 182 (1992).

≡ *Stachys fragilis* var. *rhodopaea* Velen., Fl. Bulg.: 456 (1891) [basionym]

≡ *Stachys recta* [subsp. *subcrenata* var. *ramosissima*] subvar. *rhodopaea* (Velen.) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(2): 294 (1929).

Due to the thin and fragile branches, narrow leaves, and longer calyx teeth, this subspecies was very often confused with previous subspecies, *S.*

Fig. 20. – *Stachys recta* subsp. *rhodopaea* (Velen.) Chrtek f. (SE Serbia, Pčinja Gorge, Šajince– Trgovište, coll. V. Nikolić, N. Diklić 14-Jul-1972, BEO s.n.).

recta subsp. *nitens* (Janka) Ciocârlan (Hayek 1929, Diklić 1974). It differs from the mentioned subspecies primarily by its conspicuously hairy habitus. It was described from Bulgaria, and also known from North Macedonia (Chrtek, 1992). Preliminary recorded from SE Serbia (Pčinja Gorge) in Niketić (2005).

First records:

SE Serbia, Pčinja Gorge, between Trgovište and Radovnica, rocks, MGRS 34T EM99, coll. V. Nikolić, N. Diklić 13-Jul-1972, det. N. Diklić (sub *S. recta* subsp. *subcrenata* subvar. *fragilis*), rev. M. Niketić 01-Feb-2007 (s.n. BEO).

SE Serbia, Pčinja Gorge, between Šajince and Trgovište, rocks by the road, MGRS 34T EM89, coll. V. Nikolić, N. Diklić 14-Jul-1972, det. N. Diklić (sub *S. recta* subsp. *subcrenata* subvar. *fragilis*), rev. M. Niketić 01-Feb-2007 (s.n. BEO) (Fig. 20).

SE Serbia, Pčinja Gorge, Trgovište, Vražji Kamen, MGRS 34T EM89, coll. M. Niketić, V. Vladimirov 07-Jul-2014, det. M. Niketić (BEO 100033).

A new subspecies for the flora of Serbia.

M. Niketić

SCROPHULARIACEAE

Verbascum dieckianum Borbás & Degen in Degen, Magyar Bot. Lapok 4: 82 (1905).

This C Balkan endemic species of mullein was described from North Macedonia near Skopje (Degen 1905), and was also reported from N Greece and Bulgaria (Marhold 2011+). In the region of Mt Rujan Planina in S Serbia it was also found in dry grasslands on different type of substrate (Fig. 34). The plant can be easily recognized by its brown and membranous bracts in maturity. Population size was estimated to more than 1000 reproductive individuals that are widespread within the area of Mt Rujan Planina. It was previously preliminary enounced for the same area by Niketić (2000).

First records:

S Serbia, Mt Rujan Planina, Orljak peak, limestone, MGRS 34T EM67, coll. M. Niketić 15-Apr-1991, det. M. Niketić (BEO 100020).

S Serbia, Mt Rujan Planina, Mali Orljak peak, from the pass to the top, silicate, forest and steppe-like pastures, 680-710 m, MGRS 34T EM57, coll. M. Niketić, G. Tomović 15-Aug-2010, det. M. Niketić (BEO 100021) (Fig. 21).

S Serbia, Mt Rujan Planina, Mamince village, MGRS 34T EM58, coll. M. Niketić 30-May-1991, det. M. Niketić (BEO 100022).

A new species for the flora of Serbia.

Fig. 21. – *Verbascum dieckianum* Borbás & Degen (S Serbia, Mt Rujan Planina, Mali Orljak) (photo M. Niketić).

M. Niketić, G. Tomović

New and confirmed taxa for the flora of administrative units**LILIOPSIDA**

CYPERACEAE

***Cyperus odoratus* L., Sp. Pl. 46 (1753).**

Since its first record in SE Europe in the Danube valley in Bulgaria (as "*C. strigosus*" Tzonev *et al.* 2003), this North and South American species was expected to be found upstream along the river Danube and its tributaries. This proved to be correct with the first records in Serbia originating from habitats along the river Danube in the vicinity of Dubovac, Kovin (as "*C. strigosus*" Stevanović *et al.* 2004, 2005), Labudovo Okno [<http://iasv.dbe.pmf.uns.ac.rs/index.php>] and Pančevo (Perić & Rilak 2017). According to Verloove (2014) earlier records of *C. strigosus* from Bulgaria and Serbia are erroneous and refer to *C. odoratus*. Our field investigations confirm identity of this species in Serbia as *C. odoratus* and its further spread to the west. We found this species at four localities along the mouth of the river Drina, which are the first data for this species for the flora of Serbia proper (it was previously published in abstract form, Stanković *et al.* 2018). Also, additional data for two localities in Vojvodina (vicinity of Kovilj and Gardinovci) confirm its upstream spread along the Danube (Fig. 34).

First records for Vojvodina province:

NW Serbia, Bogatić, Crna Bara, Parašnica, along the river Drina, 75 m, 19.368975 E, 44.882930 N, MGRS 34T UTM CQ77, coll. M. Stanković det. R. Perić 08-Aug-2017 (s.n. collection of Zasavica Special Nature Reserve).

NW Serbia, Zvornik, Radalj, Luke, along the Radalj river, 193 m, 19.19253 E, 44.40794 N, MGRS 34T UTM CQ51, coll. M. Stanković det. R. Perić 30-Sep-2017 (s.n. collection of Zasavica Special Nature Reserve).

NW Serbia, Velika Reka, along the Velika Reka river, 164 m, 19.23906 E, 44.27736 N, MGRS 34T UTM CQ50, coll. et det. M. Stanković 21-Sep-2016 (s.n. Private herbarium collection of M. Stanković).

NW Serbia, Ljubovija, Gračanica, along the river Gračanička, about 200 m from the rivermouth, 176 m, MGRS 34T UTM CQ69, coll. et det. M. Stanković 23-Sep-2016 (s.n. Private herbarium collection of M. Stanković) (Fig. 22).

NW Serbia, Ljubovija, Donja Ljubovidja, along the river Ljubovidja, 225 m, 19.55889 E, 44.23583 N, MGRS 34T UTM CQ79, coll. et det. M. Stanković 22-Sep-2016 (s.n. Private herbarium collection of M. Stanković).

Fig. 22. – *Cyperus odoratus* L. (NW Serbia, Ljubovija, Gračanica, s.n. collection of Zasavica Special Nature Reserve).

Additional records for Vojvodina province:

Bačka, Gardinovci, Gardinovački liman, 73 m, ≈ 20.1301 E, 45.19549 N, MGRS 34T UTM DR30, coll. et det. R. Perić 13-Sep-2011 (s.n. PZZP).

Bačka, Kovilj, Koviljski rit, Golić, along Stari Dunav oxbow, 75 m, 20.02553 E, 45.20725 N, MGRS 34T UTM DR20, coll. et det. R. Perić 23-Sep-2009 (s.n. PZZP).

A new allochthonous species for the flora of Serbia proper.

R. Perić, M. Stanković, S. Djordjević, M. Bartula, S. Djordjević-Milošević

IRIDACEAE

Crocus tommasinianus Herb., J. Hort. Soc. London 2: 273 (1847)

An indirect indications of the presence of this species in the flora of Serbia were first presented by Pančić (1856) and Janka (1868) (see on page 89) and also through the invalid name of one herbarium specimen – *C. serbicus* Kern. ex Maw (1881). The first reliable data on the distribution of the species *C. tommasinianus* Herb. in Serbia comes from Pančić (1884) for Sokolovik near Zaječar. This species was also mentioned for Mt Kopaonik (Maw 1886, sub *C. serbicus*). Later, the species was found in Serbia in many other localities south of the Sava and Danube rivers (Randjelović *et al.* 1990).

Fig. 23. – *Crocus tommasinianus* Herb. (Srem, Fruška Gora, near village Lug) (photo V. Randjelović).

The area of *C. tommasinianus* includes Croatia, Montenegro, Bosnia and Herzegovina, Serbia, Bulgaria, and North Macedonia (Pulević 1977, Randjelović *et al.* 1990). According to Barina *et al.* (2018) the only record for Albania (Meyer 2011) is doubtful. In the Pannonian plain, *C. tommasinianus* are known as allochthonous species from three localities in Hungary (Bartha & Király 2015): village Gyulaj in Tolna County (Jávorka 1964, Priszter 1964a,b, Suranyi *et al.* 2010), Villanyi hills in Baranya County (Kevey 1990, Dénes 2000) and Soproni hills in Győr-Moson-Sopron County (Király *et al.* 2009). It is also introduced as horticultural plant in UK, Belgium, Netherland, Germany and USA (POWO 2020).

A substantial population of *C. tommasinianus* was found in a beech forest not far from the village of Lug on Mt Fruška Gora. Area of occupancy was estimated at about 200 m² and subpopulation size at more than 1000 mature individuals. It is the first finding of this species for Vojvodina and the possibility that species presence in Mt Fruška Gora is due to anthropogenic factors can not be excluded.

First record for Vojvodina province:

Srem, Fruška Gora, near village Lug, beech forest, 228 m, MGRS 34T CR80, 19.5309 E, 45.1647 N, coll. et det. D. Savić, V. Randjelović, 26-Feb-2020 (HMN 14265) (Figs 23, 34).

A new species for the flora of Vojvodina province.

V. Randjelović, D. Savić

ORCHIDACEAE

Anacamptis laxiflora (Lam.) R. M. Bateman, Pridgeon & M. W. Chase, Lindleyana 12(3): 120 (1997).

Knowledge of the distribution of this Atlantic-Submediterranean species in Serbia is not sufficient (Djordjević *et al.* 2018), bearing in mind that the taxon was cited in the literature only at the specific rank for the flora of Serbia, including two subspecies of *Anacamptis* (*Orchis*) *palustris*: *elegans* and *palustris* (Diklić 1976). Consequently, published data on the distribution of *A. laxiflora* cannot be considered with certainty and therefore there is a need for further confirmation of the presence of this taxon in Serbia. It is assumed that most of the published records actually refer to the species *A. palustris*, and primarily subspecies *elegans*. Recent research has confirmed that *A. laxiflora* grows in the area of S Serbia, in the vicinity of Preševo (Niketić *et al.* 2018a). According to the recent *An annotated checklist of vascular flora of Serbia I*, its status in Kosovo and Metohija is marked as “n”, which means that there is no reliable data, but that the presence is possible (Djordjević *et al.* 2018). Several authors have published data on the distribution of this taxon in Kosovo and Metohija,

Fig. 24. – *Anacamptis laxiflora* (Lam.) R. M. Bateman (Kosovo, Priština, Kišnica, 57648 BEOU).

such as: localities in the vicinity of Dečani and Uroševac (Nikolić *et al.* 1986); Prokletije Mts (Amidžić & Panjković 2013); Kosovska Vitina (Kojić 2002); Šar Planina Mts (ZZPS 2011); Šar Planina Mts – Brezovica (Leute 1978); and Dečane – Rastavica and Uroševac – Grlica (Hundozi 1981). However, no herbarium material was found in the Herbarium of the Institute of Botany and Botanical Garden “Jevremovac”, University of Belgrade (BEOU) and in the Herbarium of Natural History Museum in Belgrade (BEO), which confirms that these findings are still unreliable.

New record for Kosovo and Metohija province:

Kosovo, Priština, Kišnica, meadows, MGRS 34T EN11, coll. P. Lazarević, det. P. Lazarević, V. Djordjević, 15-May-1996 (57648 BEOU) (Figs 24, 34).

Confirmed orchid species for the flora of Kosovo and Metohija, and the second confirmed record of this species in Serbia.

V. Djordjević, P. Lazarević

POACEAE (GRAMINEAE)

***Bromus moesiacus* Velen., Fl. Bulg.: 616 (1891).**

For the flora of Serbia it was first published by Kožuharov *et al.* (1981) for Mt Kopaonik, locality Srebrnac. Afterwards, the same record was retrieved for the Kopaonik area by Lakušić (1996), but it was never confirmed in the field. We have found herbarium specimens from Kosovo and Metohija province along the Ibar river valley (Kamenica – Donje Jarinje, Srbovac), Kosovska Mitrovica and Šar Planina Mts. The supplemented distribution in Serbia is shown in Fig. 34.

First records for Kosovo and Metohija province:

Kosovo, Ibar river gorge, Kamenica village, sporadically to Donje Jarinje, mountain pastures, sporadically on dry embankments around the road, MGRS 34T DN78, coll. D. Prodanović, Z. Krivošej May-2004, Jun-2005, det. Z. Krivošej (sub *B. erectus* subsp. *fibrosus*), rev. E. Kabaš 12-Mar-2020 (BEOU 17711).

Kosovo, Kosovska Mitrovica, Srbovac, MGRS 34T DN85, coll. B. Petković 10-Jul-1991 (sub *B. fibrosus*), rev. E. Kabaš 12-Mar-2020 (BEOU 60350).

Kosovo, Šar Planina Mts, Ostrovica, MGRS 34T DM97, coll. et det. V. Stevanović, S. Jovanović 20-Jul-1989 (sub *B. erectus*), rev. E. Kabaš 05-Feb-2020 (184/89 BEOU) (Fig. 25).

A new species for the flora of Kosovo and Metohija province.

Fig. 25. – *Bromus moesiacus* Velen. (Kosovo, Šar Planina Mts, Ostrovica, 184/89 BEOU).

E. Kabaš, S. Vukojičić, P. Lazarević

Stipa tirsa Steven, Bull. Soc. Nat. Moscou 30(2): 115 (1857).

In the *Flora of SR Serbia* it was published for Mts Rtanj and Suva Planina, and Brestovačka Banja spa in Eastern and Northeastern Serbia (Vukićević 1976). It is also known from Vršачki Breg in Vojvodina, where it was found in meadow steppe communities (Vučković 1991). Recently it was discovered in Central Serbia, on the Mt Kopaonik (Tomović *et al.* 2020) in ultramafic dry grassland steppe-like communities. Lakušić *et al.* (2018) refuted its presence in Vojvodina, however a herbarium sample from Mt Vršачke Planine (Široko Bilo) have been found (BUNS 00004340). Also new field records for Vojvodina were noted in the vicinity of Mesić (Mt Vršачke Planine).

Fig. 26. – *Stipa tirsa* Steven (Vojvodina, Mt Vršачke Planine, in the vicinity of Mesić monastery) (photo E. Kabaš).

Unpublished record for Vojvodina province:

Vojvodina, Mt Vršacke Planine, Široko Bilo, MGRS 34T EQ29, coll. et det. V. Matanović 04-Jun-1978 (sub *S. joanis*), rev. E. Kabaš 26-Mar-2019 (00004340 BUNS).

New records for Vojvodina province:

Vojvodina, Mt Vršacke Planine, Manastir Mesić, 21.387646 E, 45.103455 N, MGRS 34T EQ39, coll. et det. S. Vukojičić, E. Kabaš, P. Lazarević 30-May-2019 (BEOU 68137) (Fig. 26).

Vojvodina, Mt Vršacke Planine, Kuštilj, 21.363228 E, 45.091527 N, MGRS 34T EQ29, coll. et det. S. Vukojičić, E. Kabaš, P. Lazarević 30-May-2019. (BEOU 68220).

Confirmed species for the flora of Vojvodina province (Fig. 34).

E. Kabaš, S. Vukojičić, P. Lazarević

MAGNOLIOPSIDA

ASTERACEAE (COMPOSITAE)

Carpesium abrotanoides L., Sp. Pl. 860 (1753).

Since its first official record in the vicinity of Boljevcu near Belgrade (Radulović 2002a, 2002b, also later published in Mataruga *et al.* 2016, Jarić *et al.* 2019) and preliminary report from Apatin, published as abstract (Perić *et al.* 2005), this East-Asian adventive species was almost unknown in Serbia. Our herbarium data and field observations suggest that it is well-established species in lowland riparian forests at few previously unknown or, at least, officially unpublished localities along the rivers Danube, Bosut and Sava. Its oldest herbarium record dates back to 1996. when this species was collected at Obedska Bara. Majority of our records originates from localities along the Danube river in the vicinity of Bezdan and Apatin, where this species creates densely populated stands in poplar-willow forests understory, especially at places not directly affected by floods with dense wild boar and deer populations. Species distribution in Serbia is presented in Fig. 34.

Its additional known records are in nearby transboundary areas of Croatia (Panjković 1990) and Hungary (Kovács & Deme 2008) from where it was probably firstly introduced in Serbia (its sticky and tiny fruits are easily transported by floodwater and wild animals moving across the river). According to criteria of Niketić & Tomović (2018) its invasive status in Serbia is considered to be: A(A) (initial phase of naturalization).

First records for Vojvodina province:

Bačka, Kolut, Sirota, along the channel Sirota, MGRS 34T UTM CR38, R. Perić 16-Nov-2010 (*pers. comm.*).

Bačka, Apatin, city center, beside the wall, MGRS 34T UTM CR45, coll. et det. R. Perić 28-Aug-2004 (*s.n.* BUNS).

Fig. 27. – *Carpesium abrotanoides* L. (Srem, Bosut forests, *s.n.* PZZP).

Bačka, Apatin, Bestrement, forest edge, 79 m, 18.9471 E, 45.56927 N, MGRS 34T UTM CR34, coll. et det. R. Perić 05-Oct-2012 (*s.n.* PZZP).

Bačka, Apatin, Duboki Jendek, Veliki revir, MGRS 34T UTM CR34, R. Perić 26-Feb-2011 (*pers. comm.*).

Srem, Bosut forests, Gutiševo Bara, near a forest road, 86 m, ≈ 19.16518 E, 44.9897 N, MGRS 34T UTM CQ58, coll. et det. R. Perić 17-Sep-2013 (*s.n.* PZZP) (Fig. 27).

Srem, Obedska Bara, MGRS 34T DQ25, coll. V. Stojšić, det. R. Perić 11-Aug-2003 (*s.n.* PZZP).

Srem, Obedska Bara, Čenjin, at the mouth of Revenica river into the river Sava, MGRS 34T DQ14, coll. V. Stojšić, det. R. Perić 29-Aug-1996 (*s.n.* PZZP).

A new allochthonous species for the flora of Vojvodina province.

R. Perić

Centaurea markiana J. Wagner, Magyar Bot. Lapok 2: 284 (1903), pro hybr.

= *Centaurea magocsyana* J. Wagner, Magyar Bot. Lapok 2: 281 (1903), pro hybr.

This centaury of hybrid origin belongs to the group of transitional taxa between *C. jacea* and *C. phrygia* groups which also includes *C. macroptilon* Borbás, *C. oxylepis* Hayek, etc. It was discovered and described from Mt Vršacke Planine in Vojvodina (Banat) (Wagner 1903) as a hybrid, *C. ×markiana* J. Wagner (*C. [jacea* subsp.] *banatica* × *stenolepis*). In the same paper author also described another hybrid, *C. ×magocsyana* J. Wagner (*C. [jacea* subsp.] *banatica* × *indurata*). The same author later synonymized *C. ×magocsyana* with *C. ×markiana* within the same combination, *C. [jacea* subsp.] *banatica* × *stenolepis* (Wagner 1939, 1943). According to him there is a whole palette of transitional forms in terms of phyllaries shape, where the 'markiana type' is more reminiscent of *C. jacea* subsp. *banatica*, while 'magocsyana type' has elongated phyllaries with pectinate-fimbriate appendages, more similar to *C. stenolepis* A. Kern. (from *C. phrygia* group). In the *Flora Europaea* Dostál (1976) neglected the name *C. markiana* and treated *C. magocsyana* as a separate micro-species, similar to *C. macroptilon*. In the case of inclusion of *C. markiana* in *C. magocsyana*, the plant have to be named *C. markiana*, because Wagner (1939) established the priority of the chosen name as first to unite them (Art. 11.5). Further research will show whether this fusion is appropriate or not. Plant with 'C. magocsyana type' have been mostly observed in Vojvodina, while some forms of 'markiana type' have been found in NE Serbia for the first time (Fig. 16). These specimens are similar to *C. jacea* but differ in tomentose leaves and phyllary appendages with ± short pectinate-fimbriate acumens.

First records for Serbia proper:

NE Serbia, Miroč, from Veliki Štrbac to Miroč village, MGRS 34T EQ92, FQ03, coll. N. Diklić, M. Pajović 26-Jul-1955, det. N. Diklić (sub *C. jacea*), rev. M. Niketić 01-Apr-2020 (BEO 100010) (Fig. 28).

NE Serbia, Đerdap Gorge, Donji Milanovac, another stream towards Greben, *Carpinus orientalis* underbrush, 80-100 m, MGRS 34T EQ92,

coll. V. Nikolić, N. Diklić, M. Rakin 13-Sep-1965, det. N. Diklić (sub *C. jacea* subsp. *banatica*), rev. M. Niketić 01-Apr-2020 (BEO 100011).

Fig. 28. – *Centaurea markiana* J. Wagner (NE Serbia, Đerdap Gorge, Donji Milanovac, BEO 100010).

NE Serbia, Đerdap Gorge, Donji Milanovac, between Kazanski and Varoški stream, beach forest, 80 m, MGRS 34T EQ82, coll. V. Nikolić, N. Diklić, M. Rakin 11-Sep-1965, det. N. Diklić (sub *C. jacea* subsp. *banatica*), rev. M. Niketić 01-Apr-2020 (BEO 100012).

NE Serbia, Đerdap Gorge, Golubac, Golubac fortress, slopes towards the Danube, MGRS 34T EQ54, coll. V. Nikolić, N. Diklić 19-Sep-1967, det. V. Nikolić (sub *C. jacea* subsp. *banatica*), rev. M. Niketić 01-Apr-2020 (BEO 100013).

A new species for the flora of Serbia proper.

M. Niketić

Erechtites hieraciifolius (L.) Raf. ex DC., Prodr. 6: 294 (1838).

This species originating from temperate North & South America (Tutin 1976) has been precisely recorded in Serbia for the first and last time in the

Fig. 29. – *Erechtites hieraciifolius* (L.) Raf. ex DC. (Srem, Bosut forests, Raškovića, s.n. PZZP).

Đerdap Gorge (Iron Gate area near Tekija: Kovilovski stream) more than 50 year ago (Nikolić & Diklić 1968). Its later citation for Vojvodina province (Obradović & Panjković-Matanović 1986), despite remark that it is “domesticated” species here, has been published without any precise description of its exact habitats nor substantiated with herbarium specimens. Our field and herbarium data indicate that this species should still be considered rare casual with only two confirmed records in Serbia (Vojvodina province – Bosut forests and in the vicinity of Bezdan) (Fig. 16). Its typical habitats in Serbia are ruderal communities in wetland forests: along a forest paths and at open habitats disturbed by wild boars. Its closest next record is in the broad vicinity of Zagreb in Croatia (Nikolić 2020). Its invasive status in Serbia according to criteria of Niketić & Tomović (2018) is: A(A) (initial phase of naturalization).

First records for Vojvodina province:

Srem, Bosut forests, Raškovica, from Manješevo Bara to Široka Bara, between forest compartments 28e/28c, along a forest path, MGRS 34T UTM CQ58, coll. et det. R. Perić 17-Sep-2013 (*s.n.* PZZP) (Fig. 29).

Bačka, Bezdan-Bački Monoštor, Štrbac, Crna Bara pond, western edge, MGRS 34T UTM CR47, R. Perić 22-Oct-2013 (*pers. comm.*).

Banat, Radojevo, field road to the Hetin village, the old airport for agricultural aviation, MGRS 34T UTM DR86, G. Anačkov 29-Sep-2020 (*pers. comm.*).

Confirmed and for the first time located species for the flora of Vojvodina province.

R. Perić, G. Anačkov

***Erigeron sumatrensis* Retz.,** Observ. Bot. 5: 28 (1788).

Erigeron sumatrensis is native to South America, from where it has expanded to warm and temperate regions of the world, except to Antarctica (Vladimirov 2009 and literature therein). Concerning neighboring countries it has naturalized status in Slovenia, Croatia, Greece and Bulgaria and unknown status in Albania, Montenegro and European Turkey, while in Romania it is treated as an archeophyte (Greuter 2006+).

This adventive species was first reported for the territory of Serbia in the form of an abstract by Niketić & Jovanović (2002) for the urban area of Belgrade. Later on, it was also confirmed for several localities in Serbia proper: E Serbia – surroundings of Niš, Leskovac and Gadžin Han by Zlatković & Bogosavljević (2014) and NE Serbia – surroundings of Kladovo and Knjaževac and Šumadija – Ripanj near Belgrade by Bogosavljević & Zlatković (2018).

Fig. 30. – *Erigeron sumatrensis* Retz (Metohija, Prizren, BEO 100029).

Erigeron sumatrensis grows in sunny warm habitats, usually ruderal one such as: along railroad tracks, abandoned arable land, road embankments and field margins, but it was also recorded in secondary grasslands, in near proximity to railways and roads (Vladimirov 2009). The results of our intensive field research in the area of Serbia Proper have confirmed its great invasive potential, so *E. sumatrensis* recently been observed on clearings within the forests of the lower belt in many locations.

This plant was included in the lists of invasive species of Serbia and Vojvodina province respectively (Vrbničanin *et al.* 2004, Stojanović & Jovanović 2018, Rat *et al.* 2016). According to criteria of Niketić & Tomović (2018) its invasive status in Serbia is: A(N) Naturalised (to a greater extent domesticated). A very large and dense population was also recorded for the first time for Kosovo and Metohija along the river Prizrenka Bistrica.

First record for Kosovo and Metohija province:

Metohija, Prizren, Prizrenka Bistrica river, alluvium, ruderal sites near the river bank, 400 m, MGRS 34T DM77, coll. M. Niketić, G. Tomović, Š. Duraki 19-Jul-2007, det. M. Niketić (BEO 100029) (Fig. 30).

A new species for the flora of Kosovo and Metohija province.

M. Niketić, G. Tomović, Š. Duraki

Carlina acaulis subsp. *caulescens* (Lam.) Schübl. & G. Martens, Fl. Württemberg: 523 (1834).

≡ *Carlina caulescens* Lam., Fl. Franç. 2: 7 (1779) [basionym].

The species is listed for all neighboring countries of Serbia (Greuter 2006+) as well as for Kosovo and Metohija province (Košanin 1939, Meusel & Kästner 1994). For the flora of Serbia proper it was preliminary announced in Niketić (2000) for Mt Suva Planina in E Serbia.

First record for Serbia proper:

E Serbia, Mt Suva Planina, between Golemi Vrh and Rakoš peaks, limestone, rocky grounds and high-mountain pastures, 1500 m, MGRS 34T FN07, coll. M. Niketić 14-Aug-1988, det. M. Niketić (BEO 100014) (Figs 31, 34).

A new subspecies for the flora of Serbia proper.

Fig. 31. – *Carlina acaulis* subsp. *caulescens* (Lam.) Schübl. & G. Martens (E Serbia, Mt Suva Planina, BEO 100014).

M. Niketić

CACTACEAE

Opuntia humifusa (Raf.) Raf., Med. Fl. 2: 247 (1830).

≡ *Cactus humifusus* Raf., Ann. Nat. 1: 15 (1820) [basionym].

The natural range of this species of cactus includes arid areas of the temperate and subtropical zones of North America (POWO). It was introduced in a number of European countries where it is naturalized, especially in the Mediterranean. However, it is almost the only species of cactus that grows in the conditions of temperate-continental climate, which

is characterized by cold winters. In the surrounding countries in Croatia, Romania and Bulgaria, it has status of naturalized species and grows in similar habitats as in Serbia (Korotkova & Raab-Straube 2017).

Fig. 32. – *Opuntia humifusa* (Raf.) Raf. (Banat, Deliblatska Sands, Mramorak) (photo M. Niketić).

The first (unpublished) record of this species in Serbia comes from NE Serbia (Korbovska Sands near Kladovo), where it was found in 1998 (coll. V. Stevanović, M. Niketić, G. Tomović, BEOU 12417). Ten years later, it was observed that this small population had expanded and was showing

vitality. The origin of this population is unknown but it can be assumed that the species was introduced at Korbovska Sands from neighboring Romania. *Opuntia humifusa* was also recorded from SE Serbia in the valley of the river Pčinja near the village of Jablanica in dry sandy and rocky places on 475 m a.s.l. (coll. B. Zlatković, S. Bogosavljević, 09-Jul-2016, 13837 HMN!) (Bogosavljević & Zlatković 2018, the first published record for Serbia) (Fig. 34).

The first locality where this species of cactus was found in Vojvodina is Deliblatska Sands: Mramorak vineyards where it was observed a few years ago by V. Stevanović. During the field investigation in October 2020, *O. humifusa* was found in several micro-localities in the wider area of Mramorak vineyards c. about 6 km away from the center of the village Mramorak. The origin of this population can be related to the German inhabitants, who were the majority in the village of Mramorak before World War II and who had its vineyards on the edge of the Deliblatska Sands. It is assumed that *O. humifusa* was transferred from Germany where it was grown on the sands of the Baltic seashore. As the vineyards were abandoned *O. humifusa* spread in this part of the Deliblatska Sands in the sandy vegetation as well as in the more open *Robinia pseudoacacia* forests.

First record for Vojvodina province:

Banat, Deliblatska Sands, Mramorak, Mramorak vineyards, forest and steppe vegetation, sand, 160 m, MGRS 34T EQ07, 44.92519 N, 21.0069 E, coll. M. Niketić, V. Stevanović, G. Tomović 27-Oct-2020, det. M. Niketić, V. Stevanović (BEO 100026) (Figs 32, 34).

Additional record for the flora of Serbia proper:

NE Serbia, Korbovo, Vajuga, sand, MGRS 34T FQ33, V. Stevanović, M. Niketić, G. Tomović 15-Jun-1998, det. M. Niketić, V. Stevanović (12417 BEOU).

A new allochthonous species for the flora of Vojvodina province.

V. Stevanović, M. Niketić, G. Tomović

Erroneously reported taxa for the flora of Serbia

MAGNOLIOPSIDA

ASTERACEAE (COMPOSITAE)

Amphoricarpos neumayerianus (Vis.) Greuter, Willdenowia 33: 51 (2003) subsp. *neumayerianus*

≡ *Jurinea neumayeriana* Vis., Fl. Dalmat.: t. 10, f. 2 (1842) [basionym].

In the *Flora of SR Serbia* (Gajić 1975), as well as in older literature sources, the presence of the species *Amphoricarpos neumayerianus*

[“*Amphoricarpus neumayeri*”] in Prokletije Mts was reported without specifying its infraspecific classification. Since the type subspecies occurs outside of Serbia (see on page 90), it should be considered that this data actually refers to *A. neumayerianus* subsp. *autariatus* (Blečić & Mayer) Niketić.

M. Niketić

Carlina graeca Heldr. & Sartori ex Boiss., Exsicc. (Herb. Graec. Norm.) 1855: n. 423 (1855).

Carlina graeca is an E Mediterranean plant recorded in Albania, Greece and Anatolia (Greuter 2006+). Although Zlatković & Bogosavljević (2014) did not mention *C. graeca* as a new species for the flora of Serbia, they used this name accidentally, considering that it also includes Submediterranean *C. corymbosa* L. as a synonym. Their citation of older records (Ilić 1902 [“Ničić”], Zlatković 2006) undoubtedly refers to *C. corymbosa*. The herbarium voucher (BEOU 16835) of their new finding (Mt Rujan Planina) also corresponds to this species.

Specimen examined:

Carlina corymbosa – S Serbia, Mt Rujan Planina, Slavujevac village, silicate, dry pastures, 800 m, *Festuco-Brometea*, MGRS 34T EM68, coll. B. Zlatković, N. Smiljković 31-Jul-2008 (sub *C. graeca*), rev. B. Zlatković 15-Sep-2020 (BEOU 16835).

B. Zlatković, M. Niketić

Centaurea phrygia subsp. ***moesiaca*** (Urum. & J. Wagner) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(2): 789 (1931).

≡ *Centaurea moesiaca* Urum. & J. Wagner, Magyar Bot. Lapok 6: 165 (1907) [basionym].

The first literature data on taxon *C. phrygia* subsp. *moesiaca* (Urum. & J. Wagner) Hayek in Serbia come from Kušan (1936): “Suva planina (Petrović), Vlasina (Pančić), Zlot kod Provalija u istočnoj Srbiji (Pelivanović)”. These records have not been confirmed by field research so far. The same author also mentioned its presence in the flora of E Bosnia, but his descriptions of plants from certain localities (tomentose leaves and phyllary appendages with arcuate recurved, pectinate-fimbriate acumens) did not fit the description of taxon in the protologue and they are more in line with the characteristics of *C. stenolepis* A. Kern. Inspection of the original Pančić’s material (identified by F. Kušan) from Vlasina plateau in SE Serbia (BEOU 10518) confirmed this suspicion and specimen has been revised as *C. stenolepis*. The same specimen was previously cited in Randelović & Zlatković (2010), under the name “*C. moesiaca*”. Consider-

ring the great complexity of the *C. phrygia* group, the presence of this taxon in E Serbia could still be expected. For now, the presence of the species (*C. phrygia*) and subspecies (*moesiaca*) should be denied.

Specimen examined:

Centaurea stenolepis – SE Serbia, Vlasina plateau, coll. J. Pančić, det. F. Kušan 7-Mar-1934 (sub *C. phrygia* subsp. *moesiaca*), rev. M. Niketić 15-Feb-2020 (BEOU 10518).

M. Niketić

CARYOPHYLLACEAE

***Arenaria biflora* L., Mant. Pl. 1: 71 (1767).**

Despite different treatments on the taxonomic status and distribution of *A. biflora* L. and *A. rotundifolia* M. Bieb., the original opinion of McNeill (1963) that they are sister allopatric species has recently been resolved (Marhold 2011, POWO 2020). *Arenaria biflora* is an Alpine-Pyrenean species which presence in Bulgaria (Rila Mts) and Romania (Carpathians) should be checked. *Arenaria rotundifolia* is a Central Asiatic-Oriental-East Mediterranean orophyte which has inflorescences with several flowers and relatively narrower and ± acute petals (vs. 1–2 flowers and obtuse petals in *A. biflora*). According to McNeill (1963), in the central part of the Balkan Peninsula grows subsp. *pancicii* (Degen & Bald.) McNeill, with obovate (not orbicular) leaves, which “closely resembles the Central and East European *A. biflora*”. In the *Flora of Serbia* (Diklić & Stevanović 2012) and checklist of Albania (Barina *et al.* 2018) *A. rotundifolia* is synonymized with *A. biflora*. In Greek checklist (Dimopoulos *et al.* 2016) *A. rotundifolia* replace *A. biflora*. The study of the collected material showed that all specimens from Serbia, Montenegro and Albania belong to *A. rotundifolia*. This consequently excludes the presence of *A. biflora* in the flora of Serbia, although there are numerous literature data for it.

M. Niketić

EUPHORBIACEAE

***Euphorbia agraria* M. Bieb., Fl. Taur.-Caucas. 1: 375 (1808).**

This Pontic plant is reported for the western part of its range to the flora of Romania, Bulgaria, Greece, Macedonia and Serbia (Hayek 1924). In the *Flora of SR Serbia* (Janković & Nikolić 1972) only the presence of the variety *E. agraria* var. *subhastata* (Vis. & Pančić) Hayek was mentioned, while for the type variety it was noted that it does not grow in Serbia. Lazarević *et al.* (2013) and Niketić (2014) restored the status of the species for the Balkan endemic *E. subhastata* Vis. & Pančić which is primarily E Dinaric plant, but also was reported from Bulgaria, Dobruja and North

Fig. 33. – *Euphorbia lucida* f. *latifolia* (W. D. J. Koch) Prodan (E Serbia, Niš, Duvanište, Nišava river bank, BEO 100030).

Fig. 34. – Distribution of some new taxa for the vascular flora of Serbia or administrative units. Only the new or confirmed localities in Vojvodina are mapped for *Crocus tommasinianus* and *Stipa tirsia* as well as a new locality in Kosovo and Metohija for *Anacamptis laxiflora*. Small squares, triangles and circles within symbols: white – new record; black – literature record.

Macedonia (Hayek 1924). However, it remained questionable whether Pontic *E. agraria* M. Bieb. really grows in Serbia, and its presence has not been confirmed yet. Some sources about its presence probably refer to *E. subhastata* (treated within *E. agraria*) (Krasnići 1972) or maybe to some other species (Sigunov 1965), e.g. *E. esula* L. On the other hand, field verification (2005, BEO 100030) of the previous record for *E. agraria* var. *subhastata* from vicinity of Niš (Niketić 1985) showed that the population of this plant actually contains extremely broadleaved morphotype of *Euphorbia lucida* [f. *latifolia* (W. D. J. Koch) Prodan]. Since this morphotype is dominant in many populations of lowland areas of Serbia, its relations with the common Pontic-C European *Euphorbia lucida* Waldst. & Kit. should be clarified in the future.

Specimen examined:

Euphorbia lucida f. *latifolia* – E Serbia, Niš, Duvanište, Nišava river bank, MGRS 34T EN79, coll. M. Niketić, G. Tomović 05-Jun-2005, det. M. Niketić, G. Tomović (BEO 100030, 23831 BEOU) (Fig. 33).

M. Niketić, G. Tomović

Erroneously reported taxa for the flora of administrative units

MAGNOLIOPSIDA

ASTERACEAE (COMPOSITAE)

Scorzonera parviflora Jacq., Fl. Austriac. 4: 3 (1776). [**Serbia proper**]

Randelović *et al.* (2008) listed this taxon for the flora of Lalinac saltmarsh area, located in the vicinity of Niš. In fact, the basis for this record refers to poorly developed specimens of *T. pratensis* subsp. *orientalis* (L.) Čelak. A revision of herbarium material, as well as a detailed observation of salt marsh habitats in a given area in the recent period, confirmed the absence of *S. parviflora* from Serbia proper.

B. Zlatković

BRASSICACEAE (CRUCIFERAE)

Erysimum crepidifolium Rchb., Icon. Fl. Germ. Helv. 1: 8 (1834). [**Serbia proper**]

The distribution of this C European species in Serbia has been usually confused with the distribution of *E. diffusum* group. It has been partially corrected in the *Atlas Florae Europae* (Jalas & Suominen 1994) in which authors expressed doubt about the presence of the species in Serbia. According to Niketić (1999) “only two specimens, collected in Serbia and kept in old herbarium material, are true *E. crepidifolium*”. The first one was collected by J. Pančić in the region of Srem in Vojvodina (BEOU). The other one was collected from the modern urban area of Belgrade (Dedinje,

Košutnjak), where natural habitats under trees are still preserved (BEO). Since the collected material was in the juvenile stage, its identification was not certain. Recent field exploration has shown that *E. odoratum* Ehrh. actually grows in this locality, which has similar dentation and hairiness of the lower leaves in the juvenile stage as *E. crepidifolium*. Therefore, the presence of *E. crepidifolium* in the area of Serbia proper should be denied. However, aforementioned Pančić's specimen from Vojvodina (BEOU) still corresponds to this species, which is considered extinct from the flora of Serbia.

Specimen examined:

Erysimum odoratum – Šumadija, Beograd, Dedinje, Careva česma, MGRS 34T DQ55, coll. O. Grebenščikov 03-Apr-1945, det. O. Grebenščikov (sub *E. crepidifolium*), rev. M. Niketić 28-Feb-2020 (*s.n.* BEO).

M. Niketić

CAPRIFOLIACEAE (DIPSACACEAE)

Knautia magnifica Boiss. & Orph., Bull. Congr. Bot. Petersb. 1869: 138 (1870). [**Kosovo and Metohija**]

= *Knautia midzorensis* Formánek, Deutsche Bot. Monatsschr. 16: 19 (1898).

In Diklić (1973) and other sources there are several data on the presence of “*K. midzorensis*” in the territory of Kosovo and Metohija. According to Niketić (2014) and POWO (2020) this name has to be included in very variable *K. magnifica* Boiss. & Orph. As mentioned on page 99 its presence in administrative unit Kosovo and Metohija is unlikely and all records of *K. magnifica* (incl. *K. midzorensis*) should be treated as erroneous and attributed to *K. csikii* Jáv. & Szabo.

M. Niketić

Acknowledgements

The authors would like to thank Dr. Nicholas Turland from the Botanischer Garten und Botanisches Museum Berlin, Germany who helped us to overcome certain nomenclature concerns for *Stachys nitens*. Dr. Jindrich Chrtek jun. from the National Museum in Prague, Czech Republic and Dr Bogdan Iuliu Hurdu from the Institute of Biological Research, National Institute for Research and Development in Biological Sciences, Cluj-Napoca, Romania offered us the literature sources for *Stachys nitens*. We are thankful to Dr Božo Frajman from the University of Innsbruck, Austria for providing chorological observation for the representatives of *Euphorbia nicaeensis* complex. Special thanks to Dr Svetlana Banchева for sending and allowing the inspection of original specimens of *Centaurea phrygia* subsp. *moesiaca* from the Bulgarian Academy of Sciences, Sofia, Bulgaria (SOM).

REFERENCES

- Amidžić, L., Panjković, B., Perić, R. (2013): Chorological analysis of the endemic flora in the Metohian Prokletije mountains. – **Archives of Biological Sciences** **5(2)**: 645–650.
- Anačkov, G., Niketić, M., Tomović, G. (2018): Liliopsida: Liliaceae s.l. (Amaryllidaceae–Xanthorrhoeaceae). In: Niketić, M., Tomović, G. (2018): An annotated Checklist of Vascular Flora of Serbia 1. Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida [Catalogue of taxa]: 91–102. – Serbian Academy of Sciences and Arts, Belgrade.
- APG IV (2016): An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. – **Botanical Journal of the Linnean Society** **181**: 1–20.
- Ball, P. W. (1993): *Hesperis* L. In: Tutin, T. G., Burges, N. A., Chater, A. O., Edmondson, J. R., Heywood, V. H., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): *Flora Europaea*, ed. 2, 1: 336–337. – University Press, Cambridge.
- Barina, Z., Pifko, D., Mesterhazy, A. (2011): Contributions to the flora of Albania, 3. – **Willdenowia** **41(2)**: 329–339.
- Barina, Z., Somogyi, G., Pifkó, D., Rakaj, M. (2018): Checklist of vascular plants of Albania. – **Phytotaxa** **378(1)**: 1–339.
- Bartha, D., Király, G. (2015): Distribution atlas of vascular plants of Hungary (Atlas Florae Hungariae). – University of West Hungary, Faculty of Forestry, Institute of Botany and Nature Conservation, Sopron.
- Beck, G. (1920): Flora Bosne, Hercegovine i bivšeg Sandžaka Novog Pazara 2(9). – **Glasnik Zemaljskog Muzeja Bosne i Hercegovine** **32**: 83–127. [in Serbo-Croatian]
- Behr, O., Behr, E., Zahn, K. H. (1937): Beitrag zur Kenntnis der Hieracien der Balkanhalbinsel. – **Glasnik Skopskog Naučnog Društva** **18**, Sect. Sci. Nat. **6**: 51–67.
- Behr, O., Behr, E., Zahn, K. H. (1939a): Beiträge zur Kenntnis der Hieracien von Südserbien. – **Glasnik Skopskog Naučnog Društva** **20**, Sect. Sci. Nat. **7**: 23–34.
- Behr, O., Behr, E., Zahn, K. H. (1939b): Beiträge zur Kenntnis der Hieracien von Südserbien, Montenegro und Griecheland. – **Glasnik Skopskog Naučnog Društva** **20**, Sect. Sci. Nat. **7**: 121–129.
- Beldie, A. (1979): Flora României: determinator ilustrat al plantelor vasculare 2. – Academiei R.S.R., București.
- Blečić, V. (1953): Prilog poznavanju flore severne Crne Gore. – **Glasnik Prirodnačkog muzeja Srpske zemlje** **5–6**: 21–28. [in Serbian]
- Blečić, V., Mayer, E. (1967): Die europäischen Sippen der Gattung *Amphoricarpos* Visiani. – **Phyton** **12**: 150–158.
- Bogosavljević, S., Zlatković, B. (2018): Report on the new floristic data from Serbia II. – **Biologica Nyssana** **9(2)**: 63–75.
- Bošnjak, K. (1936): Iz hercegovačke flore. – **Glasnik Hrvatskog Prirodoslovnoga Društva** **41–48**: 62–63.

- Buttler, K. P. (1991): *Hieracium* L. In: Strid, A., Tan, K. (eds): Mountain flora of Greece 2: 595–642. – University Press, Edinburgh.
- Caković, D., Stešević, D., Schönswetter, P., Frajman, B. (2015): How many taxa? Spatiotemporal evolution and taxonomy of *Amphoricarpus* (Asteraceae, Carduoideae) on the Balkan Peninsula. – **Organisms Diversity and Evolution** 15: 429–445.
- Chater, O. A. (1972): *Brunnera* Steven. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds.): Flora Europaea 3: 106. – University Press, Cambridge.
- Chrtek, J. jun. (1992): Taxonomical Remarks on the *Stachys recta* Group. – **Folia Geobotanica and Phytotaxonomica** 27(2): 177–183.
- De Sanctis, M., Fanelli, G., Gjeta, E., Mullaj, A., Attorre, F. (2018): The forest communities of Shebenik-Jabllanicë National Park (Central Albania). – **Phytocoenologia** 48(1): 51–76.
- Degen, A. (1905): Bemerkungen über einige orientalische Pflanzenarten. – **Magyar Botanikai Lapok** 4(4–5): 82–83.
- Dénes, A. (2000): A Villányi-hegység flóra- és vegetációkutatásának története, eredményeinek összefoglalása, különös tekintettel a védett és ritka fajok előfordulására. – **Dunántúli Dolgozatok Természettudományi Sorozat** 10: 47–77. [in Hungarian with English summary]
- Diklić, N. (1972): *Alyssum* L. In: Josifović, M. (ed.): Flora SR Srbije 3: 286–311. – Srpska akademija nauka i umetnosti, Beograd [in Serbian].
- Diklić, N. (1973): Dipsacaceae B. Juss. In: Josifović, M. (ed.): Flora SR Srbije 5: 536–607 [615]. – Srpska akademija nauka i umetnosti, Beograd [in Serbian].
- Diklić, N. (1974): *Stachys* L. [In: Diklić, N., Janković, M. M.: Lamiaceae Lindley]. In: Josifović, M. (ed.): Flora SR Srbije 6: 408–432. – Srpska akademija nauka i umetnosti, Beograd [in Serbian].
- Diklić, N. (1976): Orchidaceae Lindl. In: Josifović, M. (ed.): Flora SR Srbije 8: 36–116. – Srpska akademija nauka i umetnosti, Beograd. [in Serbian]
- Diklić, N., Stevanović, V. (2012): *Arenaria* L. In: Stevanović, V. (ed.): Flora Srbije, ed. 2, 2: 251–256. – Srpska akademija nauka i umetnosti, Beograd. [in Serbian]
- Dimitrov, D. S., Vutov, V. M. (2015): *Poa stiriaca* Dörf. In: Greuter, W., Raab-Straube, E. (eds): Euro+Med-Checklist Notulae, 5: 461. – **Willdenowia** 45(3): 449–464.
- Dimopoulos, P., Raus, T., Bergmeier, E., Constantinidis, T., Iatrou, G., Kokkini, S., Strid, A., Tzanoudakis, D. (2016): Vascular plants of Greece: An annotated checklist. Supplement. – **Willdenowia** 46(2): 301–347.
- Djordjević, V., Niketić, M., Tomović, G. (2018): Liliopsida: Orchidaceae. In: Niketić, M., Tomović, G. (2018): An Annotated Checklist of Vascular Flora of Serbia 1. Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida [Catalogue of taxa]: 102–110 – Serbian Academy of Sciences and Arts, Monographs 690, Department of Chemical and Biological Sciences 10, Belgrade.

- Domina, G. (2017): Dipsacaceae. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved May 1st 2020).
- Dostál, J. (1976): *Centaurea* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 4: 254–301. – University Press, Cambridge.
- Duraki, Š., Niketić, M. (2018a): *Hieracium djimilense* subsp. *brachytrichophyes* O. Behr & al. In: Raab-Straube, E., Raus, T. (eds): Euro+Med-Check Notulae, 9: 201. – **Willdenowia** 48(2): 195–220.
- Duraki, Š., Niketić, M. (2018b): *Hieracium oroglaucum* O. Behr & al. subsp. *oroglaucum*. In: Raab-Straube, E., Raus, T. (eds): Euro+Med-Check Notulae, 9: 201–202. – **Willdenowia** 48(2): 195–220.
- Duraki, Š., Niketić, M. (2018c): *Hieracium bohatschianum* subsp. *platytrigonodon* O. Behr & al. In: Raab-Straube, E., Raus, T. (eds): Euro+Med-Check Notulae, 9: 200. – **Willdenowia** 48(2): 195–220.
- Duraki, Š., Niketić, M. (2018d): *Hieracium sparsum* subsp. *squarrosobracchiatum* O. Behr & al. In: Raab-Straube, E., Raus, T. (eds): Euro+Med-Check Notulae, 9: 202–203. – **Willdenowia** 48(2): 195–220.
- Edmondson, J. R. (1980): *Poa* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 5: 159–167. – Cambridge University Press, Cambridge, New York, Melbourne.
- Eggler, J. (1955): Ein Beitrag zur Serpentinvegetation in der Gulsen bei Kraubath in Obersteiermark. – **Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark** 85: 27–72.
- Ferakova, V., Mered'a, P., Hodálová, I., Eliáš, P., Petrović, D., Stevanović, V. (2011): *Dianthus nitidus*. In: The IUCN Red List of Threatened Species 2011: e.T161959A5519631.
- Frajman, B., Rešetnik, I., Niketić, M., Ehrendorfer, F., Schönschetter, P. (2016): Patterns of rapid diversification in heteroploid *Knaulia* sect. *Trichera* (Caprifoliaceae, Dipsacoideae), one of the most intricate taxa of the European flora. – **BMC Evolutionary Biology** 16(204): 1–20
- Formánek, E. (1896): Zweiter Beitrag zur Flora von Serbien, Macedonien und Thessalien. – **Verhandlungen des Naturforschenden Vereins in Brünn** 34: 255–365.
- Fukarek, P. (1965): Rasprostranjenost i ekološke karakteristike krčagovine *Amphoricarpus neumayeri* Vis. – **Glasnik Zemaljskog muzeja u Sarajevu, Prirodne nauke** 3–4: 159–180.
- Gajić, M. (1975): Asteraceae Dumortier. In: Josifović, M. (ed.): Flora SR Srbije 7: 1–465. – Srpska akademija nauka i umetnosti, Beograd. [in Serbian]
- Galasso, G., Conti, F., Peruzzi, F., Ardenghi, N. M. G., Banfi, E., Celesti-Grappow, L., Albano, A., Alessandrini, A., Bacchetta, G., Ballelli, S., Bandini Mazzanti, M., Barberis, G., Bernardo, L., Blasi, C., Bouvet, D., Bovio, M., Cecchi, L., Del Guacchio, E., Domina, G., Fascetti, S., Gallo, L., Gubellini, L., Guiggi, A., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E., Masin, R. R., Medagli, P., Passalacqua, N. G., Peccenini, S.,

- Pennesi, R., Pierini, B., Podda, L., Poldini, L., Prosser, F., Raimondo, F. M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Scortegagna, S., Selvaggi, A., Selvi, A., Soldano, A., Stinca, A., Wagensommer, R. P., Wilhalm, T., Bartolucci, F. (2018): An updated checklist of the vascular flora alien to Italy. – **Plant Biosystems** 152(2): 179–303.
- Gavrilović, B., Tomović, G., Niketić, M., Milenković, M., Simić, S., Radovanović, M. (2017): Geocological characteristics of plant endemism in the Balkan part of Serbia. – **Botanica Serbica** 41(2): 177–197.
- Gavrilović, M. M. (2019): Taxonomy and phylogeny of the *Xeranthemum* group (Asteraceae) – morphological, anatomical, phytochemical and molecular aspects. – University of Belgrade, Faculty of Biology, Belgrade. (PhD Thesis, manuscr.) [in Serbian with English Abstract]
- Geltman, D. V. (2015): Podrod *Esula* roda *Euphorbia* (Euphorbiaceae): sistema, filogenija, geografičeskij analiz. – Federal'noe gosudarstvennoe bjužetnoe učreždenie nauki Botaničeskij institut im. V.L. Komarova Rossijskoj akademii nauk. (PhD Thesis, manuscr.) [in Russian]
- Geltman, D. V. (2020): A synopsis of *Euphorbia* (Euphorbiaceae) for the Caucasus. – **Novitates Systematicae Plantarum Vascularium** 51: 43–78.
- Govaerts, R. (2020): World Checklist of Liliaceae. – Royal Botanic Gardens, Kew. [<http://wmsp.science.kew.org>] (retrieved September 1st 2020)
- Govaerts, R., Fernández Casas, F. J., Barker, C., Carter, S., Davies, S., Esser, H-J., Gilbert, M., Hoffmann, P., Radcliffe-Smith, A., Steinmann, V., Welzen, P., Whitmore, T. (2020): World Checklist of Euphorbiaceae. – Royal Botanic Gardens, Kew. [<http://wmsp.science.kew.org>] (retrieved October 1st 2020)
- Greuter, W. (2003): The Euro Med treatment of Cardueae (Compositae) – generic concepts and required new names. – **Willdenowia** 33(1): 49–61.
- Greuter, W. (2006+): Compositae (pro parte majore). In: Greuter, W. & Raab-Strube, E. von (eds): Compositae. Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. [<http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameId=14103&PTRefFk=7000000>] (retrieved September 1st 2020)
- Grisebach, A., Schenk, A. (1852): Iter hungaricum a. 1852 susceptum. Beiträge zur Systematik der ungarischen Flora. – **Archiv für Naturgeschichte** 18(1): 291–362.
- Hand, R. (2011): Apiaceae. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved June 1st 2020).
- Hansen, A. (1976): *Rudbeckia* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 4: 141. – Cambridge University Press, Cambridge, New York, Melbourne.
- Hayek, A. (1924): Prodromus florae peninsulae balcanicae 1(1). – **Repertorium specierum novarum regni vegetabilis, Beihefte** 30(1): 1–160.
- Hayek, A. (1929): Prodromus florae peninsulae balcanicae 2(3). – **Repertorium specierum novarum regni vegetabilis, Beihefte** 30(2): 241–336.
- Horvatić, S. (1928): Oblici sekcije *Leucanthemum* iz roda *Chrysanthemum* u flori Jugoslavije. – **Acta Botanica Instituti Botanici Universitatis Zagrebiensis**.

- Izvešća Botaničkog Zavoda kr. Sveučilišta u Zagrebu** 3: 61–140. [in Croatian with German Summary]
- Hundozi, B. (1981): Përsëri mbi disa lloje bimore të florës së Kosovës (Again about some plant species of flora of Kosovo). – **Zbornik radova PMF Priština Ser. A, Biologija-Hemija** 8: 127–131 [in Albanian with Serbian summary].
- Ilić, Đ. I. (1902): Još nekoliko biljka iz Vranjske okoline koje nisu navedene u građi za floru u okolini Vranja. Izveštaj - gimnazija Nemanjina u Vranju. – Štamparija D. Dimitrijevića. Beograd. [in Serbian]
- Ilnicki, T., Szeląg, Z. (2011): Chromosome Numbers in *Hieracium* and *Pilosella* (Asteraceae) From Central and Southeastern Europe. – **Acta Biologica Cracoviensia, Series Botanica** 53(1): 102–110.
- Jafari, F., Zarre, S., Gholipour, A., Eggens, F., Rabeler, R. K., Oxelman, B. (2020): A new taxonomic backbone for the infrageneric classification of the species-rich genus *Silene* (Caryophyllaceae). – **Taxon** 69(2): 337–368.
- Jalas, J., Suominen, J. (eds) (1994): Atlas Florae Europaeae 10. Cruciferae (*Sisymbrium* to *Aubrietia*). – Committee for Mapping the Flora of Europe, Helsinki, Societas Biologica Fennica, Vanamo.
- Janka, V. (1860): Adnotationes in plantas dacicas nonnullasque alias europaeas. – **Linnaea** 30: 549–622.
- Janka, V. (1868): Correspondenz. – **Österreichische botanische Zeitschrift** 18(2): 57.
- Janković, I., Šatović, Z., Liber, Z., Kuzmanović, N., Di Pietro, R., Radosavljević, I., Nikolov, Z., Lakušić, D. (2019): Genetic and morphological data reveal new insights into the taxonomy of *Campanula versicolor* s.l. (Campanulaceae). – **Taxon** 68(2): 340–369.
- Janković, M. M., Nikolić, V. (1972): Euphorbiaceae J. St.-Hill. In: Josifović, M. (ed.): Flora SR Srbije 3: 537–570. – Srpska akademija nauka i umetnosti, Beograd [in Serbian].
- Jarić, S., Karadžić, B., Mataruga, Z., Kostić, O., Mitrović, M., Pavlović, P. (2019): Alohtone biljne vrste u flori i vegetaciji Crnog luga (jugozapadni Srem). – **Acta herbologica** 28(1): 31–58. [in Serbian]
- Jávorka, S. (1964): Hazai *Crocus*-aink. – **Botanikai Közlemenyek** 51(2): 177–182. [in Hungarian with German summary]
- Jovanović, S., Hlavati-Širka, V., Lakušić, D., Jogan, N., Nikolić, T., Anastasiu, P., Vladimirov, V., Šinžar-Sekulić, J. (2018): *Reynoutria* taxa niche modelling and protected areas prioritization for restoration and protection from invasion: Southeastern Europe case study. – **Journal for Nature Conservation** 41: 1–15.
- Kevey, B. (1990 [1989]): Adatok Magyarország flórájának és vegetációjának ismeretéhez V. – **Botanikai Közlemenyek** 76(1–2): 83–96. [in Hungarian with German summary]
- Király, G., Baraniai-Nagy, A., Kerekes, S., Király, A., Korda, M. (2009): Kiegészítések a magyar advenív-flóra ismeretéhez IV. – **Flora Pannonica** 7: 3–31. [in Hungarian with English abstract]
- Kojić, C. S. (2002): Biogeografija Gornje Morave. – NIJP Kosovsko Pomoravlje, Gnjilane [in Serbian].

- Korotkova, N. & Raab-Straube, E. (2017): Cactaceae. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved September 1st 2020)
- Košanin, N. (1939): Über die Vegetation von Nordalbanien. – **Srpska kraljevska akademija, Spomenik 89, Prvi razred 20**: 73–107.
- Kovács, Gy., Deme, T. (2008): Idős tölgyesek az Alsó-Duna árterén. – **Somogyi Múzeumok Közleményei 18**: 43–50. [in Hungarian]
- Kožuharov, S., Petrova, A., Ehrendorfer, F. (1981): Evolutionary patterns in some brome grass species (*Bromus*, Graminae) of the Balkan Peninsula. – **Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 102**: 381–391.
- Krasnići, F. (1972): Šumska vegetacija brdskog regiona Kosova. – Zajednica naučnih ustanova Kosova. Studije 27, Priština. [in Serbian Latin].
- Kušan, F. (1936): Oblici sekcije *Eujacea* i *Lepteranthus* u flori Jugoslavije. – **Prirodoslovna istraživanja Kraljevine Jugoslavije 20**: 1–79. [in Croatian]
- Lakušić, D. (1996): Survey of the flora of Mt. Kopaonik (S. W. Serbia, Yugoslavia). – **Ekologija 31(2)**: 1–35.
- Lakušić, D., Niketić, M., Tomović, G. (2018): Liliopsida: Poaceae (Gramineae). In: Niketić, M., Tomović, G. (2018): An Annotated Checklist of Vascular Flora of Serbia 1. Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida [Catalogue of taxa]: 110–138 – Serbian Academy of Sciences and Arts, Monographs 690, Department of Chemical and Biological Sciences 10, Belgrade.
- Lakušić, R. (1966): Vegetacija livada i pašnjaka na planini Bjelasici. – **Godišnjak Biološkog instituta Univerziteta u Sarajevu 19**: 1–186.
- Lampinen R. (2001): Universal Transverse Mercator (UTM) and Military Grid Reference System (MGRS). [<http://www.luomus.fi/english/botany/afe/map/utm.htm>]
- Lazarević, P., Krivošej, Z., Tomović, G., Stevanović, V. (2013): *Euphorbia subhastata* Vis. & Pančić. In: Vladimirov, V., Dane, F., Stevanović, V., Tan. K. (eds): New floristic records in the Balkans, 22: 282. – **Phytologia Balcanica 19(2)**: 279–283.
- Lazarević, P., Zlatković, B. (2018): *Ranunculus aconitifolius* L. (Ranunculaceae), a taxon new for the flora of Serbia. – **Botanica Serbica 42(1)**: 147–151.
- Leute, G. H. (1978): Ergebnisse einer botanischen Sammelreise durch das südliche Jugoslawien im Jahre 1969. – **Annalen des Naturhistorischen Museums in Wien 81**: 75–105.
- Lozina-Lozinskaya, A. S. (1970): *Rumex* L. In: Komarov, V. L. (ed.): Flora of the U.S.S.R. 5: 445–501. – Akademii Nauk, Moskva-Leningrad.
- Maly, K. (1909): ? *Euphorbia variabilis* Ces. – **Magyar Botanikai Lapok 8(1–4)**: 93.
- Maly, K. (1934): Beiträge zur Kenntnis der *Picea omorika*. – **Glasnik Zemaljskog muzeja u Bosni i Hercegovini 46**: 37–64.
- Marhold, K. (2011+): *Verbascum*. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved May 1st 2020).

- Marhold, K. (2011): Caryophyllaceae. In: Euro+Med PlantBase - the information resource for Euro-Mediterranean plant diversity [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved May 1st 2020).
- Maslo, S., Šarić, Š., Sarajlić, N. (2020): *Sison amomum* L. (Apiaceae), a new species in the flora of Bosnia and Herzegovina. – **Glasnik Hrvatskog botaničkog društva** 8(1): 15–18.
- Mataruga, Z., Jarić, S., Karadžić, B., Mitrović, M., Kostić, O., Marković, M., Pavlović, P. (2016): Prilog poznavanju alohtone flore u donjem toku reke Save. – **Acta herbologica** 25(1): 57–70. [in Serbian]
- Matveeva, J. (1968): Floristička karakteristika na Osoj. – **Fragmenta Balcanica** 6: 141–148 [in Macedonian].
- Maw, G. (1881): A synopsis of the genus *Crocus*. – The Gardeners' Chronicle 16: 367–368.
- Maw, G. (1886): A monograph of the genus *Crocus*. – Dulau et co., London.
- McNeill, J. (1963): Taxonomic studies in the Alsinoideae: II. A revision of the species in the Orient. – **Notes From The Royal Botanic Garden Edinburgh** 24: 241–404.
- Meusel, H., Jäger, E. (1992): Vergleichende Chorologie der zentraleuropäischen Flora 3. Text. – Gustav Fischer, Jena, Stuttgart, New York.
- Meusel, H., Kästner, A. (1994): Lebensgeschichte der Gold- und Silberdisteln Monographie der mediterran-mitteuropäischen Compositen-Gattung *Carlina* 2. – Springer, Wien.
- Meyer, F. K. (2011): Beiträge zur Flora von Albanien. – **Hausknechtia**, suppl. 15: 1–220.
- Niketić, M. (1985): Prilog flori okoline Niša. In: Randelović, N. (ed.): Simpozijum Stogodišnjica flore okoline Niša, Zbornik radova: 45–49. – Niš [in Serbian Latin].
- Niketić, M. (1999): *Erysimum crepidifolium* Reichenb. In: Stevanović, V. (ed.): The Red Data Book of Flora of Serbia 1. Extinct and critically endangered taxa: 115–116, 434–435. – Ministry of Environment of the Republic of Serbia, Faculty of Biology, University of Belgrade, Institution for protection of nature of the Republic of Serbia, Belgrade. [in Serbian with English Summary]
- Niketić, M. (2000): Novi taksoni za floru cvetnica Srbije i susednih područja. In: 6. Simpozijum o flori jugoistočne Srbije i susednih područja sa međunarodnim učešćem: 32–33. – Sokobanja (Abstract) [in Serbian Latin]
- Niketić, M. (2005): Novi podaci o rasprostranjenju nekih ređih biljaka u Srbiji. In: Randjelović, N. (ed.): 8. Simpozijum o flori jugoistočne Srbije i susednih područja sa međunarodnim učešćem: 47. – Prirodno-matematički fakultet, Niš. (Abstract) [in Serbian Latin]
- Niketić, M. (2010): Novi taksoni za floru cvetnica Srbije i susednih područja 2. In: 10. Simpozijum o flori jugoistočne Srbije i susednih regiona: 32–33. – Niš (Abstract) [in Serbian Latin]
- Niketić, M. (2014): Nomenclature review of the plants published by Josif Pančić (*Nomenclator Pancicianus novus*). – **Botanica Serbica** 38(2): 209–236.

- Niketić, M., Jovanović, S. (2002): *Conyza sumatrensis* nova adventivna vrsta u flori Srbije. In: 7. Simpozijum o flori jugoistočne Srbije i susednih područja sa međunarodnim učešćem: 23. – Dimitrovgrad (Abstract) [in Serbian Latin].
- Niketić, M., Tomović, G. (2018): An Annotated Checklist of Vascular Flora of Serbia 1. Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida – Serbian Academy of Sciences and Arts, Monographs 690, Department of Chemical and Biological Sciences 10, Belgrade.
- Niketić, M., Tomović, G., Perić, R., Zlatković, B., Anačkov, G., Djordjević, V., Jogan, N., Radak, B., Duraki, Š., Stanković, M., Kuzmanović, N., Lakušić, D., Stevanović, V. (2018a): Material on the Annotated Checklist of Vascular Flora of Serbia. Nomenclatural, taxonomic and floristic notes I. – **Bulletin of the Natural History Museum in Belgrade** 11: 101–180.
- Niketić, M., Tomović, G., Anačkov, G., Djordjević, V., Lakušić, D., Randjelović, V., Jogan, N. (2018b): Komentari / Comments. In: Niketić, M., Tomović, G.: An annotated Checklist of Vascular Flora of Serbia 1. Lycopodiopsida, Polypodiopsida, Gnetopsida, Pinopsida and Liliopsida: 175–208. – Serbian Academy of Sciences and Arts, Monographs 690, Department of Chemical and Biological Sciences 10, Belgrade.
- Nikolić, T. (2020): Flora Croatica Database. – Faculty of Science, University of Zagreb, Zagreb. [<http://hirc.botanic.hr/fcd/>] (accessed: October 1st 2020).
- Nikolić, V., Diklić, N. (1968): O nekim retkim i novim vrstama biljaka za floru Jugoslavije i Srbije sa područja Đerdapske klisure. – **Glasnik Prirodnjačkog muzeja u Beogradu, serija B**, 23: 39–58. [in Serbian]
- Nikolić, V., Sigunov, A., Diklić, N. (1986): Dopuna Flori SR Srbije novim podacima o rasprostranjenju biljnih vrsta. In: Sarić, M., Diklić, N. (eds): Flora SR Srbije 10: 259–351. – Srpska akademija nauka i umetnosti, Beograd [in Serbian].
- Novaković, J., Zlatković, B., Lazarević, M., Garcia-Jacas, N., Susanna, A., Marin, P. D., Lakušić, D., Janačković, P. (2018): *Centaurea zlatiborensis* (Asteraceae, Cardueae–Centaureinae), a new endemic species from Zlatibor mountain range, Serbia. – **Nordic Journal of Botany** 36(6): e01893, 1–8.
- Obradović, M., Panjković-Matanović, V. (1986): Adventivna flora Vojvodine. – **Matica Srpska, zbornik za prirodne nauke** 70: 99–114. [in Serbian]
- Pahlevani, A. H. (2017): Diversity of the genus *Euphorbia* (Euphorbiaceae) in SW Asia. – Fakultät Biologie Chemie Geowissenschaften der Universität Bayreuth. (PhD Thesis, manuscr.).
- Pančić, J. (1856): Verzeichniss der in Serbien wildwachsenden Phanerogamen, nebst den Diagnosen einiger neuer Arten. – **Verhandlungen des Zoologisch-botanischen Vereins in Wien** 6: 475–598.
- Pančić, J. (1867): Botanische Ergebnisse einer i. J. 1866 unternommenen Reise in Serbien. – **Österreichische botanische Zeitschrift** 17(6): 166 173.
- Pančić, J. (1874): Flora Kneževine Srbije. – Državna štamparija, Beograd. [in Serbian]
- Pančić, J. (1884): Dodatak Flori Kneževine Srbije. – Kraljevsko-srpska državna štamparija, Beograd. [in Serbian]

- Panjковиć, B. (1990): Analiza životnih oblika i flornih elemenata u flori Baranje (Hrvatske). – **Acta Botanica Croatica** 49: 107–123. [in Croatian]
- Papeš, D. (1973): *Leucanthemum illyricum* (Horvatić) Papeš, comb. nov. et stat. nov. – **Acta Botanica Croatica** 32: 243–251. [in Croatian with English Summary]
- Perić, R., Knežević, J., Ručando, M., Feješ, E., Nikolić, K., Beljin, V., Dačić, J. (2005): Prilog poznavanju flore Zapadne Bačke. In: Randjelović, N. (ed.): 8. Simpozijum o flori jugoistočne Srbije i susednih područja sa međunarodnim učešćem: 58. – Prirodno-matematički fakultet, Niš. (Abstract) [in Serbian Latin]
- Perić, R., Rilak, S. (2017): *Eclipta prostrata* (L.) L. (Compositae), an adventive species new to the flora of Serbia. – **Botanica Serbica** 41(1): 89–93.
- Petrović, S. (1882): Flora okoline Niša. – Kraljevsko-srpska državna štamparija, Beograd. [in Serbian]
- Poljakov, P. P. (1961): *Artemisia* L. In: Schischkin, B. K., Bobrov, E. G. (eds): Flora USSR 26: 425–631. – Institutum botanicum nomine V. L. Komarovii Academiae scientiarum URSS, Moskva [“Mocqua”], Leningrad. [in Russian]
- POWO (2020): Plants of the world online. – The Royal Botanic Gardens, Kew. [http://www.plantsoftheworldonline.org/]
- Priszter, S. (1964a): Occurrence of *Crocus tommasinianus* Herb in Hungary. – **Acta Biologica Academiae Scientiarum Hungaricae** 15: 49. [in Hungarian with German summary] (retrieved August 25th 2020).
- Priszter, S. (1964b): Új safrányfaj (*Crocus tommasinianus* Herb.) Magyarországon. – **Botanikai Közlemények** 51(2): 183–185.
- Pulević, V. (1977): Prilog poznavanju taksonomije i horologije *Crocus tommasinianus* Herbert (Iridaceae). – **Poljoprivreda i šumarstvo (Titograd)** 223(1): 53–60.
- Radak, B. Đ., Vuku, A. Z. Peškanov, J. M., Matevski, V. S., Anačkov, G. T. (2019): Morphological characterization of three natural hybrid orchid taxa, new for Serbia, Montenegro and North Macedonia. – **Archives of Biological Sciences** 71(4): 597–607.
- Radulović, S. (2002a): *Carpesium abrotanoides* L., nova vrsta za floru Srbije u lovištu „Crni Lug“. – **Glasnik Šumarskog fakulteta** 86: 173–180 [in Serbian with English summary].
- Radulović, S. (2002b): The basic site characteristics of the species *Carpesium abrotanoides* L. – **Acta herbologica** 11(1–2): 15–22.
- Randelović, N., Stamenković, V. (1984): Flora i vegetacija Rujan planine u jugoistočnoj Jugoslaviji. – **Leskovački zbornik** 24: 375–392.
- Randelović, N., Stamenković, V. (1985): Iščezle, ugrožene i retke biljke jugoistočne Srbije predložene za zaštitu. In: Randjelović, N. (ed.): Simpozijum „Stogodišnjica flore okoline Niša“. – Zbornik radova. Univerzitet u Nišu, Tehnološki fakultet u Leskovcu. Podružnica Srpskog biološkog društva Niš, 57–65. [in Serbian Latin]
- Randelović, V., Zlatković, B. (2010): Flora i vegetacija Vlasinske visoravni. – Prirodno-matematički fakultet, Niš. [in Serbian Latin]

- Randelović, V., Zlatković, B., Dimitrijević, D. (2008): Fitogeografska analiza flore Lalinačke slatine. In: 9th Symposium on the flora of Southeastern Serbia and neighboring regions, Proceeding: 73–82. – University of Niš, Faculty of Sciences and Mathematics, Niš. [in Serbian with English abstract]
- Randjelović, N., Hill, D. A., Randjelović, V. (1990): The genus *Crocus* L. in Serbia. – Serbian Academy of Sciences and Arts, Monographs 601, Department of Natural and Mathematical Sciences 66, Belgrade.
- Rat, M., Simonović, P., Glavendekić, M., Paunović, M., Stojanović, V., Karaman, M., Radišić, D., Anačkov, G. (2016): Overview of the invasive alien species in Serbia. In: Rat, M., Trichkova, T., Scalera, R., Tomov, R., Uludag, A. (eds): ESENIAS Scientific Reports 1. State of the Art of Alien Species in South-Eastern Europe. [Country reports]. – University of Novi Sad, Novi Sad, Serbia, IBER-BAS, Bulgaria, ESENIAS, Novi Sad, 91–114.
- Raus, Th., Strid, A. (2020): New combinations in some Balkan *Centaurea* taxa. – **Phytologia Balcanica** 26(3): 505–507.
- Răvărut, M. (1960): *Stachys* L. In: Săvulescu, T., Nyárády, E. I. (eds): Flora Republicii Populare Romîne 8: 208–233. – Academia Republicii Populare Romîne, București. [in Romanian]
- Rohlena, J. (1913): Fünfter Beitrag zur Flora von Montenegro. – **Sitzungsberichte der Königlich Böhmisches Gesellschaft der Wissenschaften. Mathematisch-naturwissenschaftliche Classe** 1912: 1–143.
- Rohlena, J. (1942): Conspectus Florae Montenegrinae. – **Preslia** 20–21: 1–506.
- Sabovljević, M. S., Tomović, G., Niketić, M., Lazarević, P., Lazarević, M., Latinović, J., Latinović, N., Kabaš, E., Đurović, S. Z., Kutnar, L., Skudnik, M., Pantović, J., Stevanoski, I., Vukojičić, S., Veljić, M. (2020): New records and noteworthy data of plants, algae and fungi in SE Europe and adjacent regions, 1. – **Botanica Serbica** 44(1): 81–87.
- Sell, P. D., West, C. (1976): *Hieracium* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 4: 358–410. – Cambridge University Press, Cambridge, New York, Melbourne.
- Sigunov, A. (1965): Prilog poznavanju flore i šumske vegetacije okoline Majdanpeka – **Glasnik Muzeja šumarstva i lova** 5: 55–102. [in Serbian Latin].
- Skokanová, K., Hodálová, I., Mered'a Jr., P., Slovák, M., Kučera, J. (2019): The *Cyanus tuberosus* group (Asteraceae) in the Balkans: biological entities require correct names. – **Plant Systematics and Evolution** 305(8): 569–596.
- Sl. glasnik RS (2010-2011): Pravilnik o proglašenju i zaštiti strogo zaštićenih i zaštićenih divljih vrsta biljaka, životinja i gljiva. – „Sl. glasnik RS“, br. 5/2010 i 47/2011. [in Serbian]
- Smith A. R., Tutin, T. G. (1968): *Euphorbia* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 2: 213–226. – University Press, Cambridge.
- Stanković, M., Đorđević, S., Bartula, M., Đorđević-Milošević, S., Milošević, S., Pejić, D. (2018): New sites of allochthonous species *Cyperus odoratus* L. (fam. Cyperaceae) in the middle flow of the Drina river watershed. In: Anačkov, G., Zorić, L. (eds): 7th Balkan Botanical Congress, Book of

Abstracts. – University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology, Novi Sad, Botanical Society “Andreas Wolny”, Novi Sad. – **Botanica Serbica** 42 (Supplement 1): 121.

Stefanović, V. (1960): Tipovi šuma bijelog bora na području krečnjaka istočne Bosne. – Naučno društvo Bosne i Hercegovine, Radovi, XVI, Odjeljenje privredno-tehničkih nauka, knjiga 4, Sarajevo.

Stevanović, V., Šinžar-Sekulić, J., Stevanović, B. (2004): Expansion of the adventive species *Paspalum paspaloides* (Michx.) Scribn. and *Echinochloa oryzoides* (Ard.) Fritsch and *Cyperus strigosus* L. in the Yugoslav part of the Danube reservoir (km 1090-1075). – **The Proceedings of the International Association for Danube Research** 35: 399–405.

Stevanović, V., Tan, K., Tomašević, M., Uotila, P. (2005): The occurrence of *Cyperus strigosus* (Cyperaceae) in Serbia and Montenegro. – **Phytologia Balcanica** 11(2): 137–138.

Stojanović, V., Jovanović, I. (2018): Pregled invazivnih i potencijalno invazivnih vrsta biljaka u Republici Srbiji i okruženju u cilju utvrđivanja njihovog statusa na nacionalnom nivou. – **Zaštita prirode** 68(1-2): 41–59. [in Serbian with English summary]

Stojanović, V., Stevanović, V. (2008): Prikaz flore planine Gučevo u Severozapadnoj Srbiji. – **Zaštita prirode** 59(1–2): 93–108.

Suranyi, G., Máthé, C., Mosolygó, A. Borbély, G., Vasas, G. (2010): Analysis of genetic diversity in crocuses with Carpathian Basin origin using AFLP-markers. – **Acta Biologica Hungarica** 61 (Suppl.): 149–155.

Süveges, K., Löki, V., Lovas-Kiss, Á., Ljubka, T., Fekete, R., Takács, A., Vincze, O., Molnár, V. A. (2019): From European priority species to characteristic apophyte: *Epipactis tallosii* (Orchidaceae). – **Willdenowia** 49(3): 401–409.

Svoboda, C. (2012): Grasses (Poaceae) from the Balkan Peninsula. – **Studii și cercetări, Biology** 17: 19–45.

Szeląg, Z. (2015): *Hieracia balcanica* X. Typification of the *Hieracium* (Asteraceae) names described by K. H. Zahn from the Balkan Peninsula. – **Polish Botanical Journal** 60(1): 41–65.

Szeląg, Z., Vladimirov, V. (2013): A new species of *Hieracium* sect. *Pannosa* (Asteraceae) from Bulgaria. – **Phytotaxa** 108(1): 57–60.

Šajinović, B., Mihajlov, M. (1979): Jovan Tucakov and the Deliblato Sand. In: Čanak, M. (ed.): II Međunarodni simpozijum o zaštiti i unapređivanju Deliblatskog peska. – Društvo ekologa Vojvodine, Novi Sad. [in Serbian with English summary]

Španiel, S., Zozomová-Lihová, J., Marhold, K. (2017): Revised taxonomic treatment of the *Alyssum montanum*-*A. repens*. – **Plant Systematics and Evolution** 303(10): 1413–1442.

complex in the Balkans: a multivariate morphometric analysis

Tan, K. (1991): *Amphoricarpos* Vis. In: Strid, A., Tan, K. (eds): Mountain flora of Greece 2: 472–473. – University Press, Edinburgh.

- Thiers, B. (2021): Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih> (retrieved October 10th 2020)
- Tomović, G., Niketić, M., Lakušić, D., Randjelović, V., Stevanović, V. (2014): Balkan endemic plants in Central Serbia and Kosovo regions: distribution patterns, ecological characteristics and centres of diversity. – **Botanical Journal of the Linnean Society** 176(2): 173–202
- Tomović, G., Sabovljević, M. S., Djokić, I., Petrović, P., Djordjević, V., Lazarević, P., Mašić, E., Barudanović, S., Ștefănuț, S., Niketić, M., Butorac, B., Pantović, J., Hajrudinović Bogunić, A., Bogunić, F., Kabaš, E., Vukojičić, S., Kuzmanović, N., Djurović, S. Z., Buzurović, U. (2020): New records and noteworthy data of plants, algae and fungi in SE Europe and adjacent regions, 2. – **Botanica Serbica** 44(2): 251–259.
- Turland, N. J., Wiersema, J. H., Barrie, F. R., Greuter, W., Hawksworth, D. L., Herendeen, P. S., Knapp, S., Kusber, W.-H., Li, D.-Z., Marhold, K., May, T. W., McNeill, J., Monro, A. M., Prado, J., Price, M. J., Smith, G. F. (2018): International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. – *Regnum Vegetabile* 159.
- Tutin, T. G. (1968): *Sison* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds.): *Flora Europaea* 2: 352. – University Press, Cambridge.
- Tutin, T. G. (1976): *Erechtites* Rafin. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds.): *Flora Europaea* 4: 191. – Cambridge University Press, Cambridge, New York, Melbourne.
- Tzonev, R., Zieliński, J., Tan, K. (2003): *Cyperus strigosus* (Cyperaceae), a naturalized species new to Bulgaria. – **Polish Botanical Journal** 48(1): 47–49.
- Valdés, B. (2011): Boraginaceae. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. [<http://ww2.bgbm.org/EuroPlusMed/>] (retrieved May 1st 2020).
- Verloove, F. (2014): A conspectus of *Cyperus* s.l. (Cyperaceae) in Europe (incl. Azores, Madeira and Canary Islands), with emphasis on non-native naturalized species. – **Webbia** 69(2): 179–223.
- Vladimirov, V. (2009): *Erigeron sumatrensis* (Asteraceae): a recently recognized alien species in the Bulgarian flora. – **Phytologia Balcanica** 15(3): 361–365.
- Vrbničanin, S., Karadžić, B., Dajić-Stevanović, Z. (2004): Adventivne i invazivne korovske vrste na području Srbije. – **Acta herbologica** 13(1): 1–12. [in Serbian with English summary]
- Vučković, M. (1991): Livadska i livadsko-stepska vegetacija Vršackih Planina. – Matica Srpska, Odeljenje za prirodne nauke, Novi Sad. [in Serbian with English summary]
- Vukićević, E. (1965): Sukcesija vegetacije i prirodno obnavljanje šuma na šumskim požarištima u Srbiji. Doktorska disertacija. – **Glasnik Šumarskog fakulteta** 29: 1–87.

- Vukićević, E. (1976): *Stipa* L. In: Josifović, M. (ed.): Flora Srbije 8: 316–322. – Srpska akademija nauka i umetnosti, Beograd. [in Serbian]
- Vuksanović, S. (2016): Rasprostranjenje, horološka struktura i centri diverziteta balkanske endemične flore u Crnoj Gori. – University of Belgrade, Faculty of Biology, Belgrade. (PhD Thesis, manuscr.) [in Serbian with English Summary].
- Wagner, J. (1903): Neue *Centaurea*-Bastarde. – **Magyar Botanikai Lapok** 2(9–10): 281–287.
- Wagner, J. (1939): Bemerkungen über *Centaureen*. – **Acta Universitatis Szegediensis, Section Science Naturalis, Pars Bot.** 5(3–4): 73–113.
- Wagner, J. (1943): Észrevételek Nyárády E. Gy.: Adatok a hazai *Centaureák* “*Jacea*” és “*Phrygia*” csoportjainak bővebb ismeretéhez, különös tekintettel Kolozsvár környékének flórájára. – **Borbásia Nova** 15: 1–23. [in Hungarian]
- Webb, D. A. (1976). *Amphoricarpus* Vis. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. (eds): Flora Europaea 4: 243. – University Press, Cambridge.
- Wraber, T. (1982): De *Euphorbia pancicii* G. Beck notula. – **Glasnik Republičkog zavoda za zaštitu prirode – Prirodnjačkog muzeja u Titogradu** 15: 77–84.
- Wraber, T. (1988): The discovery of *Dianthus nitidus* Waldst. & Kit. in Montenegro (Yugoslavia). – **Biološki vestnik** 36(4): 95–102.
- Zahn, K. H. (1921): Compositae (*Hieracium*). In: Engler, A. (ed.): Das Pflanzenreich: 289–576. – Regni Vegeabilis Conspectus 76 (IV.280), Wilhelm Engelmann, Leipzig.
- Zahn, K. H. (1936): *Hieracium* L. In: Ascherson, P. F. A. & Graebner, K. O. P. P. (eds): Synopsis der mitteleuropäischen Flora, 12(3): 1–320. – Gebrüder Borntraeger, Berlin.
- Zahn, K. H. (1937): *Hieracium* L. In: Ascherson, P. F. A. & Graebner, K. O. P. P. (eds): Synopsis der mitteleuropäischen Flora, 12(3): 321–480. – Gebrüder Borntraeger, Berlin.
- Zahn, K. H. (1938): *Hieracium* L. In: Ascherson, P. F. A. & Graebner, K. O. P. P. (eds): Synopsis der mitteleuropäischen Flora, 12(3): 481–708. – Gebrüder Borntraeger, Berlin.
- Zlatković, B. (2006): *Carlina corymbosa* L. In: Greuter, W., Raab-Straube, E. (eds): Euro+Med Notulae, 2: 709. – **Willdenowia** 36 (2): 707–717.
- Zlatković, B. K., Bogosavljević, S. S. (2020): Risk analysis of alien plants recorded in thermal waters of Serbia. – **Weed Research** 60(1): 85–95.
- Zlatković, B., Bogosavljević, S. (2014): Report on the new floristic data from Serbia. – **Biologica Nyssana** 5(2): 123–129.
- Zlatković, B., Randjelović, V., Randjelović, N. (1993): Građa za floru jugoistočne Srbije. In: Randjelović, N. (ed.): III Simpozijum o flori jugoistočne Srbije. Flora i vegetacija. Zbornik radova. – Univerzitet u Nišu, Tehnološki fakultet u Leskovcu. Leskovac-Pirot, 95–110. [in Serbian Latin]
- ZZPS (2011): Stručna osnova za izradu zakona o Nacionalnom parku „Šarplanina“. Studija zaštite. – Zavod za zaštitu prirode Srbije, Beograd. [in Serbian]

**МАТЕРИЈАЛ ЗА КРИТИЧКУ ЛИСТУ ВРСТА
ВАСКУЛАРНЕ ФЛОРЕ СРБИЈЕ.
НОМЕНКЛАТУРНИ, ТАКСОНОМСКИ
И ФЛОРИСТИЧКИ ПРИЛОЗИ II**

Р Е З И М Е

МАРЈАН НИКЕТИЋ, ГОРДАНА ТОМОВИЋ, ГОРАН АНАЧКОВ, МИРЈАНА
БАРУЛА, СЛАЂАНА ЂОРЂЕВИЋ, ВЛАДАН ЂОРЂЕВИЋ, СУЗАНА
ЂОРЂЕВИЋ-МИЛОШЕВИЋ, ШЕМИЈА ДУРАКИ, МИЛАН ГАВРИЛОВИЋ,
ПЕЋА ЈАНАЋКОВИЋ, ЕВА КАБАШ, НЕВЕНА КУЗМАНОВИЋ, ДМИТАР
ЛАКУШИЋ, ПРЕДРАГ ЛАЗАРЕВИЋ, РАНКО ПЕРИЋ, ВЛАДИМИР
РАНЂЕЛОВИЋ, ДРАГИША САВИЋ, МИХАЛЈО СТАНКОВИЋ, ИВАНА
СТЕВАНОВИЋ, ВЕРИЦА СТОЈАНОВИЋ, ОЛГА ВАСИЋ, СНЕЖАНА
ВУКОЛИЧИЋ, БОЈАН ЗЛАТКОВИЋ, ВЛАДИМИР СТЕВАНОВИЋ

У раду је предложено девет нових номенклатурних комбинација и назначено је осам нових синонима. Поред тога, представљено је 12 таксона (врста и подврста) васкуларне флоре који су нови за флору Србије (од тога девет аутохтоних и три алохтоне биљке). Потврђени су стари или непоуздани наводи за три аутохтоне и једну алохтону биљку у флори Србије. Осам биљних таксона представљају новину за ужу Србију, Војводину или Косово и Метохију, а за три биљке је доказано присуство у одређеним административним јединицама у Србији. Последњи део рада садржи информације о осам таксона чије је присуство у Србији или у њеним територијалним јединицама оповргнуто.