

DOI: 10.5937/SustFor2490119V

UDK: 630*443.3(497.11)=111

Original scientific paper

THE MOST COMMON FUNGI ASSOCIATED WITH A DECLINE OF TURKEY OAK (*QUERCUS CERRIS* L.) IN URBAN CONDITIONS IN SERBIA

Aleksandar VEMIĆ¹*, Zlatan RADULOVIĆ¹, Katarina MLADENović¹,
Ljubinko RAKONJAC¹

Abstract: Turkey oak (*Quercus cerris*) is one of the most important tree species in park forests and parks in Serbia. Despite this, there has been no available information in domestic literature about the parasitic fungi that affect Turkey oak in urban areas. Research carried out between 2016 and 2023 identified 19 taxa of parasitic and saprotrophic fungi that colonize Turkey oak in urban conditions: one on the roots, two on the leaves, two on the bark, seven on the trunk, two on the branches, two on the fruits, two on the stumps, and one on both leaves and fruits. The most significant fungi found were *Fomes fomentarius*, *Inonotus nidus-pici*, and *Fuscoporia torulosa*, which cause heart rot and are typically found on individual trees. Most of the fungi identified occurred in succession. Following primary damage, the most frequently occurring fungus was *Stereum hirsutum*, while after mechanical injuries, *Schizophyllum commune* was most frequently recorded. *Alternaria* spp. was found on old leaves and heavily damaged leaves. To protect the urban Turkey oak trees, measures should be focused on reducing tree density and preventing mechanical injuries. The findings from this research also contribute to understanding the ecological characteristics of these fungal taxa based on their frequent occurrence in urban conditions.

Keywords: oak decline, causes, urban areas, first record.

NAJČEŠĆE GLJIVE POVEZANE SA PROPADANJEM STABALA CERA (*QUERCUS CERRIS* L.) U URBANIM PODRUČJIMA SRBIJE

Sažetak: Cer (*Quercus cerris*) predstavlja jednu od najvažnijih vrsta drveća u park šumama i parkovima Srbije. Do sada u domaćoj literaturi nije bilo podataka vezano za parazitski kompleks gljiva koje kolonizuju cer u urbanim uslovima. Istraživanje izvršeno u periodu 2016-2023 je pokazalo 19 taksona parazitskih i saprofitskih gljiva; jedan na korenu, dva na listovima, dva na kori, sedam na deblu, dva na granama, dva na plodovima, dva na panjevima i jedan na listovima i plodovima. Najvažnije gljive su bile *Fomes fomentarius*, *Inonotus nidus-pici* i *Fuscoporia torulosa* prouzrokovajući centralne truleži, koje su konstatovane na pojedinačnim stablima. Većina konstatovanih gljiva se javljala u sukcesiji. Posle primarnih uzročnika oštećenja, najčešća je *Stereum hirsutum*, dok je posle mehaničkih oštećenja najčešće konstatovana *Schizophyllum commune*. *Alternaria* spp. je zabeležena na

¹ Institute of Forestry, Kneza Višeslava 3, 11030 Belgrade, Serbia

*Corresponding author. E-mail: aleksandar.vemic2@gmail.com

starim listovima, kao i posle teških oštećenja listova. Mere zaštite cera treba da budu usmerene u smanjenju gustine i izbegavanju mehaničkog oštećenja stabala. Dobijeni rezultati takođe doprinose poznavanju ekoloških karakteristika taksona gljiva na osnovu njihove česte konstatacije u urbanim sredinama.

Ključne reči: propadanje hrastova, uzročnici, urbana područja, prvi nalaz.

1. INTRODUCTION

The *Quercus* genus comprises approximately 500 tree species in the northern hemisphere (Cvjetićanin et al., 2016). One particularly significant species in Europe and Asia Minor is the Turkey oak (*Quercus cerris* L.), which can be found in mixed forests in the Mediterranean basin and is commonly planted in parks due to its ability to withstand urban conditions (Cvjetićanin et al., 2016; de Rigo et al., 2016). Climate change is influencing the increasing importance of Turkey oak because of its enhanced adaptability to drier conditions and ability to recover after periods of drought. Its adaptability is somewhat lower in Southern Europe (Kasper et al., 2022; Mészáros et al., 2022; Šimková et al., 2023). Consequently, it is crucial to initiate research on the biotic factors contributing to the decline of these trees to enhance the likelihood of preserving as many trees as possible. Currently, there is limited detailed information available in domestic literature on the diseases affecting Turkey oak trees, particularly in urban areas where the trees are more exposed to air pollution and extremely high temperatures.

Fungi that harm Turkey oak can be found at all stages of its growth. To properly assess the impact of these fungi, it is important to know the stage of development of the tree's vegetative organs (Bercea, 2010). Additionally, in the case of Turkey oak, drought, and insect defoliation can contribute to the appearance of specific pathogens, such as *Biscogniauxia mediterranea* (De Not.) Kuntze (Capretti & Battisti, 2007; Vannini et al., 2009).

The health status of trees in urban areas reflects the development and quality of life in those areas (Solomou et al., 2019). Moreover, specific genera of fungi, such as *Alternaria* and *Cladosporium*, which appear on trees, can cause allergic reactions in humans (Kasprzyk et al., 2021). Experiences from various parts of Europe have demonstrated that fungal diseases are the most common cause of decay in Turkey oak trees (Georgieva et al., 2018).

In light of the above, a multi-year study was conducted in Serbia to investigate the diversity of fungi associated with symptoms of Turkey oak deterioration. The initial hypotheses were: i) there are not the same symptoms caused by different fungi on Turkey oak trees; ii) the causes of Turkey oak decay in urban areas are not different from those in rural areas.

2. MATERIAL AND METHODS

2.1 Field methods

Field research was conducted in all major cities of Serbia, with a focus on Belgrade. The research involved inspecting symptomatic trees and collecting

samples from these trees for laboratory analysis. The trees were inspected three times a year: in spring, summer, and autumn. Every part of the trees was visually inspected, and samples were taken for laboratory analysis in case of visible damage. Samples of tree parts or fruiting bodies of fungi were placed in paper bags and stored in a refrigerator until the analyses began.

2.2 Laboratory methods

The fungi were isolated and identified based on their morphological characteristics from the samples collected. When identifying fungi directly from plant material, we used temporary histological preparations following the procedure described by Muntanola Cvetković (1990). We followed the description by Ellis & Ellis (1985) to identify microfungi from histological preparations. The fungi were isolated using MEA (30 g/l Biolab, Hungary; 20 g/l Torlak, Serbia) and PDA (LAB M, UK) nutrient media. Pure culture identification was carried out using the keys provided by Nobles (1948, 1964) and Stalpers (1978).

3. RESULTS AND DISCUSSION

Based on the samples taken, 19 parasitic and saprophytic fungi taxa were identified on Turkey oak in urban areas of Serbia (see Table 1).

Table 1. Identified fungi on Turkey oak in urban areas of Serbia

Species	Tree part	Type of damage	Frequency	Significance
<i>Alternaria</i> spp.	Leaves, fruits	- Fruit rot	++	+
<i>Armillaria mellea</i> (Vahl) Kumm.	Root	Butt rot	+	+++
<i>Botrytis cinerea</i> Pers.	Fruits	Fruit rot	++	+++
<i>Cladosporium</i> spp.	Fruits	Fruit rot	++	+
<i>Cytospora ambiens</i> (Pers.) Sacc.	Bark	Bark necrosis	+	++
<i>Diplodia</i> spp.	Leaves	Spotting	+	++
<i>Fomes fomentarius</i> (L.) Fr.	Trunk	Heart rot	+	+++

Species	Tree part	Type of damage	Frequency	Significance
<i>Gnomonia quercina</i> Kleb.	Leaves	Leaf anthracnose	+	+++
<i>Inonotus hispidus</i> (Bull.) Karst.	Trunk	Heart rot	++	+++
<i>Irpex lacteus</i> Fr.	Stump	-	+	+
<i>Inonotus nidus-pici</i> Pilát	Trunk	Heart rot	+++	+++
<i>Peniophora</i> spp.	Branches	Sap rot	+	+
<i>Peniophora quercina</i> (Pers.) Cooke	Branches	Sap rot	++	+
<i>Peziza</i> spp.	Bark	-	+	+
<i>Omphalotus olearius</i> (DC.) Singer	Stump	-	++	+
<i>Fuscoporia torulosa</i> (Pers.) Wagn. & Fisch.	Trunk	Heart rot	+	+++
<i>Schizophyllum commune</i> Fr.	Trunk	Sap rot	+++	++
<i>Stereum hirsutum</i> (Willd.) Pers.	Trunk	Sap rot	+++	++
<i>Trametes hirsuta</i> (Wulf.) Pil.	Trunk	Sap rot	+++	++

Addendum to the table: + Fungus appears rarely or has no significant importance

++ Fungus occurs moderately often and is important

+++ Fungus appears often and has great importance

The largest number of taxa was found on the trunk (Table 1). It was also found that fungi from this group cause the most damage to trees (Table 1, Figure 1). Fungi from this group are categorized as those that directly cause damage to trees (*Fomes fomentarius*, *Inonotus hispidus*, *Inonotus nidus-pici* and *Fuscoporia torulosa*), and others that occur on damaged trees (Table 1, Figure 1).

The fungus *Schizophyllum commune* was found on both uninjured and injured trees (see Figure 1). Description of the damaged wooden substrate type and fungi occurrence is shown in Table 2.

Table 2. *The condition of the tree damage and the presence of fungi*

Damage type	Fungus species
Separated bark	<i>Schizophyllum commune</i> <i>Stereum hirsutum</i>
Widened radial fissures on the bark	
Windbreaks	
Stumps	<i>Fuscoporia torulosa*</i> <i>Irpex lacteus</i> <i>Omphalotus olearius</i>

* The fungus causes tree cutting




Figure 1. The most common fungi on the trunk and stumps of Turkey oak trees in the cities of Serbia: A – *Inonotus nidus-pici*, B – *Inonotus hispidus*, C – *Irpex lacteus*, D – *Peniophora* spp., E-F – *Schizophyllum commune*



Figure 2. The most common fungi on the trunk, leaves, and stumps of Turkey oak trees in the cities of Serbia: A – *Omphalotus olearius*, B – *Alternaria* spp., C – *Fuscoporia torulosa*

The number of fungal species found on Turkey oak in urban areas was lower compared to the number of fungi found in natural stands in Serbia (Karadžić et al., 2017). This could be due to the negative impact of urbanization on the presence of fungi (Meyer et al., 2021). In urban environments, there is a reduction in the diversity of fungi in the air, which is not the case with the fungi in the soil (Abrego et al., 2020). However, it is worth noting that most fungi found on Turkey oak in this study were facultative parasites, indicating that these species are actively dispersed through the air in urban conditions, unaffected by the environment. Additionally, the limited number of fungi found on the roots of Turkey oak does not necessarily mean a decrease in soil fungal diversity, as only the most common and noticeable fungi were examined for Turkey oak.

It is necessary to pay attention to *Inonotus nidus-pici*, which causes the most damage on Turkey oak in natural and urban areas (Karadžić & Milenković, 2015; Milenković et al., 2022). This fungus was recorded in Serbia on *Fraxinus angustifolia* and *Tilia platyphyllos* in urban areas, where infection spreads through wounds, probably due to climate change (Milenković et al., 2023). Most recorded fungi occur in the natural Turkey oak stands (Karadžić et al., 2017). Also, different fungi taxa have been found on previously damaged trees, causing white rot. Therefore, the first null hypothesis was rejected and the alternative hypothesis that different fungi on Turkey oak trees cause the same symptoms was accepted.

The findings are consistent with previous research (Ragazzi et al., 2003), which suggests fewer fungi on Turkey oak trees in urban areas than forests. The study also indicates that urban temperatures and conditions may contribute to the increased pathogenicity of endophytic fungi. The study identified the same fungal species in urban areas as in natural stands. As a result, the hypothesis that the causes of Turkey oak decline in urban areas are not different from those in natural areas has been rejected, and the alternative hypothesis has been accepted. Additionally, the study found no previously recorded individual species that rarely appear on Turkey oak trees (Vemić & Milenković, 2019). The researchers recommend monitoring rarely occurring fungi, such as the recently identified species *Trametes trogii* (Radulović et al., 2023), especially on other oak species to prevent the spread of infection, particularly in the context of globalization.

4. CONCLUSION

For the first time, this study detailed the diversity of the most common parasitic and saprophytic fungi on Turkey oak trees in urban conditions in Serbia. The results obtained are as follows:

- Nineteen taxa were identified, most of which were found for the first time on Turkey oak in urban areas in Serbia.
- The most important were the fungi found on the trunk. From this group, the most important were *Fomes fomentarius*, *Inonotus nidus-pici*, and *Fuscoporia torulosa*, causing white rot.
- The fungi *Schizophyllum commune* and *Stereum hirsutum* are often found on trees mechanically damaged by abiotic factors.

- The obtained results enable better planning of Turkey oak tree protection measures in the urban conditions of Serbia.

Acknowledgment: This research was conducted under the 2024 Ministry of Science, Technology Development, and Innovation financing contract.

REFERENCES

- Abrego, N., Crosier, B., Somervuo, P., Ivanova, N., Abrahamyan, A., Abdi, A., Hämäläinen, K., Junninen, K., Maunula, M., Purhonen, J., & Ovaskainen, O. (2020). Fungal communities decline with urbanization — more in air than in soil. *The ISME Journal*, 14(11), 2806–2815. <https://doi.org/10.1038/s41396-020-0732-1>
- Bercea, I. (2010). Hungarian oak (*Quercus frainetto*) and Turkey oak (*Quercus cerris*) damaging fungi. *Annals of the University of Craiova - Agriculture, Montanology, Cadastre Series* 40(2), 29-38.
- Capretti, P., & Battisti, A. (2007). Water stress and insect defoliation promote the colonization of *Quercus cerris* by the fungus *Biscogniauxia mediterranea*. *Forest Pathology* 37(2), 129-135. <https://doi.org/10.1111/j.1439-0329.2007.00489.x>
- Cvjetičanin, R., Brujić, J., Perović, M., & Stupar V. (2016). Dendrology. Faculty of Forestry, University of Belgrade (Original title: Dendrologija. Univerzitet u Beogradu, Šumarski fakultet).
- de Rigo, D., Enescu, C.M., Houston Durrant, T., & Caudullo, G. (2016). *Quercus cerris* in Europe: distribution, habitat, usage and threats. In: J. San-Miguel-Ayanz, D. de Rigo, G. Caudullo, T. Houston Durrant, & A. Mauri (Eds.), *European Atlas of Forest Tree Species* (pp. e01b479+). Publ. Off. EU, Luxembourg.
- Ellis, M.B., & Ellis, J.P. (1985). *Microfungi on land plants*. Croom Helm, London and Sidney.
- Georgieva, M., Nedjalkov, D., Georgiev, G., Matova, M., Zaemdzhikova, G., Petkov, P., & Mirchev, P. (2018). *Monitoring of health status of Quercus cerris L. in the Eastern Balcan range and the Ludogorie (Bulgaria)*. Annual of Sofia University, Faculty of Biology, Book 4 Scientific sessions of Faculty of Biology / Volume 104, 2019 - International Scientific Conference Kliment's Days 2018 (pp. 154-165).
- Kasper, J., Leuschner, C., Walentowski, H., Petritan, A.M., & Weigel, R. (2022). Winners and losers of climate warming: Declining growth in *Fagus* and *Tilia* vs. stable growth in three *Quercus* species in the natural beech–oak forest ecotone (western Romania). *Forest Ecology and Management* 506, 119892. <https://doi.org/10.1016/j.foreco.2021.119892>
- Kasprzyk, I., Grinn-Gofroń, A., Ćwik, A., Kluska, K., Cariñanos, P. & Wójcik, T. (2021). Allergenic fungal spores in the air of urban parks. *Aerobiologia* 37, 39-51. <https://doi.org/10.1007/s10453-020-09671-7>
- Karadžić, D., & Milenković, I. (2015). Contribution to the knowledge of parasitic fungus *Inonotus nidus-pici* Pilát the agent of canker on broadleaved trees. *Forestry* 1-2, 15-30. (Original title: Prilog poznavanju parazitarne gljive *Inonotus nidus-pici* Pilát uzročnika rak-rana na stablima lišćara. *Šumarstvo* 1-2, 15-30).

- Karadžić, D., Milenković, I., Milanović, S., Golubović-Ćurguz, V., & Tomović, Z. (2017). The most important parasitic and saprophytic fungi in oak forests in the area of Public Enterprise "Vojvodinašume". Public Enterprise "Vojvodinašume", Petrovaradin, Serbia. (Original title: Najznačajnije parazitske i saprofitske gljive u hrastovim šumama na području JP „Vojvodinašume“. JP „Vojvodinašume“, Petrovaradin, Srbija).
- Nobles, M.K. (1948). Studies in Forest Pathology. VI. Identification of cultures of wood-rooting fungi. *Canadian Journal of Research* 26, 281-431. <https://doi.org/10.1139/cjr48c-026>
- Nobles, M.K. (1964). Identification of cultures of wood-inhabiting *Hymenomyces*. *Canadian Journal of Botany*, 43, 1097-1139. <https://doi.org/10.1139/b65-126>
- Mészáros, I., Adorján, B., Nyitrai, B., Kanalas, P., Oláh, V., & Levanič, T. (2022). Long-term radial growth and climate-growth relationships of *Quercus petraea* (Matt.) Liebl. and *Quercus cerris* L. in a xeric low elevation site from Hungary. *Dendrochronologia* 76, 126014. <https://doi.org/10.1016/j.dendro.2022.126014>
- Meyer, S., Rusterholz, H-P., & Baur, B. (2021). Saproxylic insects and fungi in deciduous forests along a rural–urban gradient. *Ecology and Evolution* 11(4), 1634-1652. <https://doi.org/10.1002/ece3.7152>
- Milenković, I., Radulović, Z., Jovanović, D., & Karadžić, D. (2023). Two new hosts of the parasitic fungus *Inonotus nidus-pici* Pilát in Serbia. *Forestry* 3-4, 115-126. (Original title: Dva nova domaćina parazitske gljive *Inonotus nidus-pici* Pilát u Srbiji. *Šumarstvo* 3-4, 115-126).
- Milenković, I., Radulović, Z., Tomšovský, M., Sikora, K., Golubović Ćurguz, V., Jovanović, D., & Karadžić, D. (2022). Distribution and new hosts of the parasitic fungus *Inonotus nidus-pici* in Serbia. *Bulletin of the Faculty of Forestry, University of Banja Luka*, 32, 5-16.
- Muntanola Cvetković, M. (1990). General mycology. Scientific book, Belgrade (Original title: Opšta mikologija. Naučna knjiga, Beograd).
- Radulović, Z., Karadžić, D., & Milenković, I. (2023). *Trametes trogii* Berk. - Bioecological characteristics, importance and possibility of use for medicinal purposes (medicinal properties). *Forestry* 3-4, 83-98. (Original title: *Trametes trogii* Berk. – Bioekološke karakteristike, značaj i mogućnost korišćenja u lekovite svrhe (lekovita svojstva). *Šumarstvo* 3-4, 83-98).
- Ragazzi, A., Moricca, S., Turco, E., Marianelli, L., & Dellavalle, I. (2003). *The mycobiota of healthy and declining oaks in an urban setting*. Second International Symposium on plant health in urban horticulture, Berlin, Germany, 27-29 August, 2003 (pp. 43-47).
- Solomou, A.D., Topalidou, E.T., Germani, R., Argiri, A., & Karetos, G. (2019). Importance, Utilization and Health of Urban Forests: A Review. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 47(1), 10-16. <https://doi.org/10.15835/nbha47111316>
- Stalpers, J.A. (1978). Identification of Wood inhabiting Aphyllophorales in pure culture. *Studies in mycology* 16, 1-248.
- Šimková, M., Vacek, S., Šimůnek, V., Vacek, Z., Cukor, J., Hájek, V., Bílek, L., Prokúpková, A., Štefančík, I., Sitková, Z., et al. (2023). Turkey Oak (*Quercus cerris* L.) Resilience to Climate Change: Insights from Coppice Forests in Southern and Central Europe. *Forests* 14, 2403. <https://doi.org/10.3390/f14122403>

Vannini, A., Lucero, G., Anselmi, N., & Vettrai, A.M. (2009). Response of endophytic *Biscogniauxia mediterranea* to variation in leaf water potential of *Quercus cerris*. *Forest Pathology* 39(1), 8-14. <https://doi.org/10.1111/j.1439-0329.2008.00554.x>

Vemić, A., & Milenković, I. (2019). A contribution to the knowledge of rare lignicolous fungi in the forests of Serbia and Montenegro. *Forestry* 1-2, 155-162. (Original title: Prilog poznavanju retkih lignikolnih gljiva u šumama Srbije i Crne Gore. *Šumarstvo* 1-2, 155-162).

THE MOST COMMON FUNGI ASSOCIATED WITH A DECLINE OF TURKEY OAK (*QUERCUS CERRIS* L.) IN URBAN CONDITIONS IN SERBIA

Aleksandar VEMIĆ, Zlatan RADULOVIĆ, Katarina MLADENOVIĆ, Ljubinko RAKONJAC

Summary

A multi-year study examined the mycoflora associated with symptoms in Turkey oak trees (*Quercus cerris*) in Serbian cities due to the need to preserve trees in urban areas. Nineteen fungi taxa were identified, and their distribution varied on different parts of the Turkey oak trees. The most significant number of taxa, including *Fomes fomentarius*, *Inonotus hispidus*, *Inonotus nidus-pici*, *Fuscoporia torulosa*, *Schizophyllum commune*, *Stereum hirsutum*, and *Trametes hirsuta*, was found on the trunk. Fewer taxa, such as *Alternaria* spp., *Diplodia* spp., and *Gnomonia quercina*, were found on the leaves; *Cytospora ambiens* and *Peziza* spp. were found on the bark; *Alternaria* spp., *Botrytis cinerea*, and *Cladosporium* spp. were found on the fruits; *Peniophora* spp. and *Peniophora quercina* were found on the branches; and *Irpex lacteus* and *Omphalotus olearius* were found on the stumps. The smallest number of taxa was found on the root, specifically *Armillaria mellea*.

The most significant damage was caused by *Fomes fomentarius*, *Inonotus nidus-pici*, and *Fuscoporia torulosa*, which resulted in breaks and premature felling of the affected trees. Long-term monitoring revealed that these species were present on trees with no mechanical damage, indicating their role as the primary cause of damage. Conversely, *Schizophyllum commune* and *Stereum hirsutum* were found on damaged trees, and *Peniophora quercina* was observed on fallen branches.

The number of pathogens on the leaves was not high enough to threaten the health of the trees. *Alternaria* spp. was commonly found on leaves before they fell and on leaves that were damaged by different biotic and abiotic factors. *Botrytis cinerea* was the primary fungus affecting the fruits, causing gray mold in the cupule and nut. Other fungi were also frequently seen, but their significance was not determined.

It is essential to control the number of Turkey oak trees in park forests and parks to prevent mechanical damage, which can lead to the growth of various fungi. All the different types of fungi found should be considered when developing forest protection plans to minimize the spread of their inoculum and reduce damage to Turkey oak trees in urban areas.

The study recorded many fungi for the first time on Turkey oak trees in urban areas in Serbia. These findings provide valuable insights into the ecology of the identified fungi in urban conditions based on their presence on Turkey oak trees.

NAJČEŠĆE GLJIVE POVEZANE SA PROPADANJEM STABALA CERA (*QUERCUS CERRIS* L.) U URBANIM PODRUČJIMA SRBIJE

Aleksandar VEMIĆ, Zlatan RADULOVIĆ, Katarina MLADENović, Ljubinko RAKONJAC

Rezime

Usled potrebe očuvanja stabala u urbanim područjima, izvršeno je višegodišnje istraživanje mikoflore povezane sa simptomima na stablima cera (*Quercus cerris*) u gradovima Srbije. Utvrđeno je devetnaest taksona gljiva, čija distribucija je različita na različitim delovima stabala cera. Najveći broj taksona: *Fomes fomentarius*, *Inonotus hispidus*, *Inonotus nidus-pici*, *Fuscoporia torulosa*, *Schizophyllum commune*, *Stereum hirsutum*, *Trametes hirsuta* je konstatovan na deblu. Manji broj taksona: *Alternaria* spp., *Diplodia* spp. i *Gnomonia quercina* su konstatovane na listovima, na kori: *Cytospora ambiens* i *Peziza* spp., plodovima: *Alternaria* spp., *Botrytis cinerea* i *Cladosporium* spp., granama: *Peniophora* spp. i *Peniophora quercina*, panjevima: *Irpex lacteus* i *Omphalotus olearius*. Najmanji broj taksona je konstatovan na korenu (*Armillaria mellea*).

Najveća oštećenja su zabeležena od *Fomes fomentarius*, *Inonotus nidus-pici* i *Fuscoporia torulosa*, koja su dovela do lomova i prevremene seče zahvaćenih stabala. Takođe, višegodišnji monitoring je pokazao da su ove vrste konstatovane na pojedinačnim stablima na kojima nije bilo mehaničkih oštećenja, ukazujući na njihov značaj kao glavnih uzročnika oštećenja. Sa druge strane, *Schizophyllum commune* i *Stereum hirsutum* su konstatovane na oštećenim stablima, a *Peniophora quercina* na opalim granama.

Patogeni listova nisu zabeleženi u većoj brojnosti koja bi ugrozila vitalnost stabala. Najčešće je konstatovana *Alternaria* spp., na listovima pred opadanjem, kao i na listovima oštećenim usled različitih abiotičkih i biotičkih uzročnika. Na plodovima je bila najznačajnija *Botrytis cinerea* koja je izazivala sivu plesan kupula i orašica. Ostale gljive su takođe često konstatovane ali njihov značaj nije utvrđen.

Mere zaštite treba da budu usmerene u regulisanju gustine stabala cera u park šumama i parkovima, iz razloga smanjenja mehaničkog oštećenja koje pogoduje razvoju velikog broja konstatovanih gljiva. Sve konstatovane vrste gljiva treba uključiti u šire strategije zaštite šuma u cilju ograničenja prenosa njihovog inokuluma, kako bi se umanjile štete na ceru u urbanim uslovima.

Većina zabeleženih gljiva u ovoj studiji je prvi put konstatovana na ceru u urbanim uslovima Srbije. Dobijeni rezultati omogućavaju bolje poznavanje ekologije konstatovanih gljiva bazirano na njihovim nalazima na ceru u urbanim uslovima.