

---

Mirjana Šumarac Dumanović<sup>1,2</sup>

## DA LI JE GOJAZNOST BOLEST?

**Sažetak:** Gojaznost je kompleksan entitet koji može da nastane zbog različitih uzroka, kao što su endokrini (u disfunkciji štitaste žlezde ili hiperfunkciji nadbubrežne žlezde – Kušingov sindrom), ali je najčešće gojaznost kombinacija neaktivnosti i prekomernog unosa hrane. S druge strane posmatrano, postoje i genetski faktori koji stvaraju sklonost ka nastanku prekomerne težine čak i kada je unos hrane onoliki koliko kalorija bi bio odgovarajući broj za većinu ljudi. Da li su uzroci gojaznosti genetski, hormonalni ili su negde u mozgu (sistem nagrade i zadovoljstva ili pak centralno uslovljeni ustaljeni obrasci ponašanja kad su u pitanju navike u ishrani, veličina porcija, izbor hrane ...) je teško izdvojiti. Zagovornici stava da je gojaznost bolest smatraju da zadovoljava kriterijume da se definiše kao bolest. Gojaznost smanjuje očekivanu dužinu života i oštećuje normalno funkcionisanje tela, može da bude uslovljena i genetskim nasleđem. Oponenti ove ideje smatraju da gojaznost nije bolest već stanje koje je faktor rizika za druge bolesti i koje može da se prevenira. Gojaznost je rezultat unosa previše hrane uz nedovoljnu aktivnost. Gojaznost je udružena sa različitim bolestima, kao što su dijabetes tipa 2, ateroskleroza, kardiovaskularne bolesti i neki karcinomi, pa se može smatrati odgovornom za visoke stope morbiditeta i mortaliteta. Razumevanje patofiziologije aterogeneze je značajno poraslo tokom poslednjih nekoliko decenija. Patogenetski mehanizmi u gojaznosti i u nastanku i razvoju komorbiditeta, koji prate gojaznost, pokazuju mnoge karakteristike inflamatornih procesa. Ključnu ulogu i u patogenezi gojaznosti mogao bi da igra imunski sistem. Uprkos identifikovanju brojnih kritičnih elemenata u ovim procesima i pronalaženja novih terapijskih modaliteta u borbi protiv gojaznosti, lečenje gojaznosti je još uvek veliki izazov i uglavnom sa ne tako uspešnim ishodima.

**Ključne reči:** gojaznost, inflamacija, komorbiditeti, belo masno tkivo, braon masno tkivo

---

<sup>1</sup> Medicinski fakultet Univerziteta u Beogradu

<sup>2</sup> Klinika za endokrinologiju, dijabetes i bolesti metabolizma, Multidisciplinarni centar za lečenje gojaznosti, Klinički centar Srbije, email: msumaracdumanovic@gmail.com.

## *Uvod*

Gojaznost je povezana sa različitim bolestima, kao što su dijabetes tipa 2, ateroskleroza, kardiovaskularne bolesti i neki karcinomi, pa se može smatrati odgovornom za visoke stope morbiditeta i mortaliteta u ovoj populaciji. Prevalenca gojaznosti u svetu je, prema najnovijim statistikama Svetske zdravstvene organizacije, skoro udvostručena od 1980. godine. Svake godine zbog gojaznosti umire najmanje 2,8 miliona ljudi, a očekuje se dodatno povećanje u narednih par dekada (1). Gojaznost je kompleksan entitet koji može da nastane zbog različitih uzroka kao što su endokrini (u disfunkciji štitaste žlezde ili hiperfunkciji nadbubrežne žlezde – Kušingov sindrom), ali je najčešće gojaznost kombinacija neaktivnosti i prekomernog unosa hrane. S druge strane posmatrano, postoje i genetski faktori koji stvaraju sklonost ka nastanku prekomerne težine čak i kada je unos hrane onoliki koliko kalorija bi bio odgovarajući broj za većinu ljudi. Da li su uzroci gojaznosti genetski, hormonalni ili su negde u mozgu (sistem nagrade i zadovoljstva ili pak centralno uslovljeni ustaljeni obrasci ponašanja kad su u pitanju navike u ishrani, veličina porcija, izbor hrane ...) je teško izdvojiti. Zagovornici stava da je gojaznost bolest smatraju da zadovoljava kriterijume da se definiše kao bolest. Gojaznost smanjuje očekivanu dužinu života i oštećuje normalno funkcionisanje tela, može da bude uslovljena i genetskim nasleđem. Oponenti ove ideje smatraju da gojaznost nije bolest već stanje koje je faktor rizika za druge bolesti i koje može da se prevenira. Gojaznost je rezultat unosa previše hrane uz nedovoljnu aktivnost. Patogenetski mehanizmi u gojaznosti i u nastanku i razvoju komorbiditeta koji prate gojaznost pokazuju mnoge karakteristike inflamatornih procesa (2). Ključnu ulogu u patogenezi gojaznosti bi mogao da igra imunski sistem. U toku su obimna istraživanja radi identifikovanja kritičnih igrača u ovim procesima i pronalaženja novih terapijskih modaliteta u borbi protiv gojaznosti. Uprkos tome, lečenje gojaznosti je još uvek veliki izazov i uglavnom sa ne tako uspešnim ishodima. Zato se danas govori i istražuje sličnost i povezanost između gojaznosti i ateroskleroze u cilju lečenja i prevencije oba ova stanja. Zanimljivo je da novije studije pokazuju da aktivacija bež ili braon adipocita i posledično povećanje potrošnje energije može da smanji masu masnog tkiva u telu i potencijalno smanji inflamaciju masnog tkiva (3). Osim toga, postoji sve više dokaza da je funkcija braon i bež masnog tkiva regulisana imunskim procesima (4). Sadašnja strategija je povećanje potrošnje energije od strane braon masnog tkiva (BAT) sa ciljem da se smanje preterani energetske depoi u gojaznosti. Intervencije u imunskim putevima mogu da prethode razvoju novih strategija koje bi trebalo da povećaju aktivnost braon i bež masnog tkiva kao terapijski target za lečenje gojaznosti.

## ***Uloga hronične niskostepene inflamacije u patogenezi gojaznosti i udruženih komorbiditeta***

Prvi dokaz da je inflamacija važna za patogenezu gojaznosti i da dovodi do opšte metaboličke disfunkcije pružio je Hotamisligil i sar. (5). Proinflamatorni citokin faktor nekroze tumora (TNF) je pronađen u WAT-u i korelirao je sa insulinskom rezistencijom kod ljudi i miševa (5). Postalo je jasno da infiltracija proinflamatornih makrofaga u WAT-u igra centralnu ulogu u inflamatornom odgovoru kao dominantan izvor TNF-a (5). U WAT normalno uhranjenih miševa, 10–15% ćelija su makrofagi, dok u WAT gojaznih sadrže 45–60% makrofaga (6). Rezydentni makrofagi u WAT negojaznih imaju dominantno antiinflamatorni fenotip, dok su u gojaznosti inflamatorni monociti regrutovani u WAT, gde podležu diferencijaciji i postaju proinflamatorni ili M1 fenotip i čine većinu makrofaga (7). Antiinflamatorni ili M2 makrofagi zavise od citokina IL-4 i IL-13 i zahtevaju STAT6 da održe svoje alternativno aktiviranje (8). Druge mijeloidne ćelije koje igraju ulogu u WAT-u su neutrofil i eozinofil. Neutrofil su veoma kratkoživeće ćelije koje su već prisutne u WAT-u u roku od 3 dana od unosa visokomasne hrane (high fat diet -HFD) (9). Nasuprot tome, broj eozinofila je u obrnutoj korelaciji sa gojaznošću. Iscrpljenost eozinofila kod miševa rezultira povećanjem telesne mase, oštećenom tolerancijom glukoze i insulinskom rezistencijom (10). Za oba tipa 2 urođenih limfoidnih ćelija (ILC2s) i eozinofila je tek nedavno pokazano da su važna populacija ćelija u WAT-u i predstavljaju dominantan izvor IL-4 i IL-13, citokina potrebnih za indukciju polarizacije M2 makrofaga (10). T- ćelije su takođe komponenta repertoara imunskih ćelija u WAT. Deset procenata stromalne vaskularne frakcije WAT-a u negojaznih čine T-ćelije. Veliki deo čine CD4 T-helper ćelije, od kojih su oko 50% regulatorne T-ćelije (Tregs). Kod ljudi je broj T-ćelija u WAT-u u korelaciji sa BMI (11). Kod miševa se količina T-ćelija u WAT-u povećava u roku od 2 nedelje od HFD-a. Postoji samo nekoliko CD8+ citotoksičnih T-ćelija i CD4+ efektornih T-ćelija u WAT negojaznih, ali obe populacije rastu drastično u gojaznih, dok se CD4 + Tregs smanjuju (13). Slično se i odnos M1 i M2 makrofaga povećava u WAT-u kod gojaznih, kao i odnos Th1 i Th2 T-ćelija. Ovo dovodi do smanjenja u Th2 indukovanim citokinima kao što su IL-4 i IL-13, čime se smanjuje polarizacija M2 makrofaga. Porast Th1 T-ćelija i citotoksičnih T-ćelija dovodi do prekomerne sekrecije TNF i IFN $\gamma$ , koja polarizuje makrofage u proinflamatorno stanje, dovodeći do povećane inflamacije u WAT gojaznih (10,12). Hronična niskostepena inflamacija u WAT gojaznih takođe uključuje i regrutovanje B-ćelija, ćelija prirodnih ubica (NK) i mast-ćelija (13). NK- ćelije se aktiviraju prepoznavanjem lipidnih antigena, a mast-ćelije sadrže granule koje mogu da oslobađaju različite medijatore kao što je histamin, serotonin i citokini, koji takođe promovišu regrutovanje inflamatornih ćelijskih populacija (13).

### ***Uloga braon i bež masnog tkiva***

Za razliku od jasne uloge različitih ćelija imunskog sistema u WAT-u, doprinos imunskog sistema u razvoju, funkciji i aktivnosti BAT-a je još uvek nepoznat. Međutim, znamo da gojazne osobe imaju smanjenu količinu aktivnog BAT-a, mereno preko preuzimanja glukoze (FDG), koje je u korelaciji sa njihovim niskostepenim inflamatornim stanjem. Štaviše, neaktivni braon adipociti akumuliraju lipide, slično belim adipocitima. Ablacija noradrenergičnog stimulusa selektivnom simpatičkom denervacijom BAT-a rezultuje prelaskom braon u bele adipocite sa velikim intraćelijskim vakuolama. Pošto je regrutovanje makrofaga u WAT u korelaciji sa lipolizom uskladištenih triglicerida (14), verovatno je da oslobađanje masnih kiselina takođe izaziva regrutovanje imunskih ćelija u BAT. Međutim, da li je to zaista tako još uvek je nepoznato. U gojaznosti indukovanoj ishranom (DIO), termoneutralno okruženje dovodi do dodatnog povećanja inflamacije u WAT-u i vaskulaturi u poređenju sa normalnim uslovima stanovanja. Iako ne izaziva povećanje insulinske rezistencije, povećanje vaskularne inflamacije dovodi do progresije ateroskleroze (14), ukazujući da BAT štiti protiv ateroskleroze indukovane gojaznošću. U jednoj od studija imunokompromitovani miševi doživljavaju hladno kao stres kada su smešteni na 23 °C što modulira homeostazu energije i telesne mase i tako ih štiti od razvoja gojaznosti. Međutim, u termoneutralnim uslovima (33 °C), oni razvijaju gojaznost, povećanu akumulaciju triglicerida u jetri, povećanje markera zapaljenja i netoleranciju glukoze (15). Ovo ukazuje da BAT štiti od metaboličkog nereda i inflamacije u masnom tkivu. Pored temperature okoline, drugi podsticaji, kao što su biološki sat (16), hormoni (17) i unos hrane ne samo da moduliraju potrošnju energije putem hipotalamusa, već utiču i na inflamaciju. Na primer, vremenski ograničen unos hrane sa većim sadržajem masti (HFD) na 8 sati dnevno povećava BAT aktivnost i smanjuje hipertrofiju adipocita i upalu u odnosu na Ad libitum HFD hranjenih miševa (18). Hormoni digestivnog trakta poput GLP-1 posreduju efekte na unos hrane, potrošnju energije i inflamaciju. GLP-1 receptorski signaling aktivira BAT i promoviše braonizaciju WAT-a (19, 20), dok GLP-1 takođe smanjuje infiltraciju makrofazima i inflamatorni signaling u belim adipocitima i makrofagima (21, 22). Druge hormonalne promene, kao što je menopauza, takođe utiču na energetske metabolizam. Estradiol inhibira AMPK u hipotalamusu, koja aktivira termogenezu u BAT-u (18). Zaista, ovarijektomisani miševi sa smanjenim estradiolom dobijaju više u telesnoj masi, i imaju veći stepen inflamacije u WAT-u (23).

### ***Kliničke implikacije***

Prekomeran unos energije dovodi do povećanog skladištenja lipida u belim i braon adipocitima, što dovodi do disfunkcije ovih ćelija. Imunske ćelije i signali u

belom (WAT) i braon masnom tkivu (BAT) su neophodni za homeostazu ovih tkiva. One doprinose ulasku lipida koji se nalaze u belim masnim ćelijama i visokoj stopi oksidacije u braon i bež masnom tkivu. Imune ćelije, uključujući eozinofile i alternativno aktivirane makrofage, imaju regulatornu ulogu u metaboličkoj homeostazi kako WAT-a tako i BAT-a. Brojna istraživanja se bave identifikacijom preciznih imunoloških aktera. Imunološki procesi predstavljaju intrigantan terapijski cilj – smanjenje dobitanja na telesnoj masi povećanjem potrošnje energije. Broj imunih ćelija u normalno uhranjenih osoba, kao i u BAT gojaznih je mnogo manji nego u WAT. To ukazuje da su BAT rezistentnije na inflamaciju izazvanu dijetom, ali kod pozitivnog energetskog balansa inflamacija se ipak dešava i u BAT-u. Veliki značaj bi imala identifikacija metaboličke povezanosti između imunih ćelija i braon, bež i belih adipocita i redosleda događaja tokom razvoja gojaznosti. Važna pitanja koja treba rešiti se odnose na detektovanje regulatornih efektornih molekula koje sekretuju braon adipociti (ili bež ili beli adipociti koji bi da postanu bež): da bi privukli ili regulisali imune ćelije? Kako je BAT aktivnost regulisana u gojaznosti? Koja je uloga simpatičkog nervnog sistema? I kako se menja aktivnost BAT-a tokom starenja? Na kraju, prisustvo BAT-a kod odraslih ljudi i njihovo potencijalno aktiviranje ili indukovanje prelaska bež masnih ćelija u WAT je zanimljivo za lečenje ili čak sprečavanje poremećaja koji su povezani sa gojaznošću. Hladnoća je još uvek daleko najjači simpatički signal za aktiviranje BAT-a. Međutim, deo opsežnih istraživanja je traganje za identifikovanjem biohemijskih i imunoloških puteva koji su odgovorni za aktivaciju BAT-a, a da to nije prolongirano izlaganje hladnoći. Cilj je da se nađe način povećanja aktivnosti BAT-a i promene energetskog metabolizma uticajem na imunološke odgovore. Razumevanje ovih složenih interakcija će možda da doprinese definisanju novih potencijalnih biomarkera kardiovaskularnog i potencijalnih terapijskih meta.

---

Mirjana Šumarac Dumanović<sup>1,2</sup>

## IS OBESITY A DISEASE?

**Abstract:** Obesity is a complex entity that can have many causes, such as endocrine (like thyroid dysfunction or hyperfunctioning of the suprarenal gland–Cushing’s syndrome) but often obesity is from a combination of inactivity and overeating. On the other side, there are genetic factors that produce a tendency to overweight even with the consumption of what would be for most people an appropriate number of calories. Whether the causes are hormonal, genetic or reside in the brain (its reward system or the circuitry that underlies habit, perception of portion size, the choice of food...) is often difficult to sort out. Proponents contend that obesity is a disease because it meets the definition of disease. Obesity decreases life expectancy and impairs the normal body functions, also it can be caused by genetic factors. Opponents contend that obesity is not a disease because it is a preventable risk factor for other diseases. Obesity is the result of eating too much as well as it is caused by exercising too little. Formally disease or condition obesity is associated with a variety of diseases such as type 2 diabetes, atherosclerosis, cardiovascular diseases and certain cancers, and may also be responsible for high rates of morbidity and mortality. Understanding the pathophysiology of obesity has grown significantly over the last few decades. Pathogenetic mechanisms in obesity and in the development of comorbidities that accompany obesity exhibit many of the characteristics of inflammatory processes. A key role in the pathogenesis of obesity could play the immune system. Despite identifying many critical players in these processes and finding new therapeutic modalities in the fight against obesity, treatment of obesity is still a great challenge and mostly with not-so-successful outcomes.

**Key words:** obesity, inflammation, comorbidity, white adipose tissue, brown adipose tissue

---

<sup>1</sup> Medical faculty, University in Belgrade

<sup>2</sup> Clinic for endocrinology, diabetes and metabolic diseases, Multidisciplinary center for treatment Obesity, Clinical Center of Serbia, email: msumaracdumanovic@gmail.com.

## ***Introduction***

Obesity is associated with a variety of medical conditions such as type 2 diabetes, atherosclerosis, cardiovascular diseases and some cancers and is therefore responsible for high morbidity and mortality rates in these population. According to the latest statistics of the world health organization, the worldwide prevalence of obesity, has nearly doubled since 1980 and at least 2.8 million people die each year as a result of obesity. This number is expected to further increase over the next decade (1). Obesity is a complex entity that can have many causes, such as endocrine (like thyroid dysfunction or hyperfunctioning of the suprarenal gland-Cushing's syndrome) but often obesity is from a combination of inactivity and overeating. On the other side, there are genetic factors that produce a tendency to overweight even with the consumption of what would be for most people an appropriate number of calories. Whether the causes are hormonal, genetic or reside in the brain (its reward system or the circuitry that underlies habit, perception of portion size, the choice of food...) is often difficult to sort out. Proponents contend that obesity is a disease because it meets the definition of disease. Obesity decreases life expectancy and impairs the normal body functions, also it can be caused by genetic factors. Opponents contend that obesity is not a disease because it is a preventable risk factor for other diseases. Obesity is the result of eating too much as well as it is caused by exercising too little.

Pathogenetic mechanisms in obesity and in the development of comorbidities that accompany obesity exhibit many of the characteristics of inflammatory processes (2). A key role in the pathogenesis of obesity could play the immune system. Despite identifying many critical players in these processes and finding new therapeutic modalities in the fight against obesity, treatment of obesity is still a great challenge and mostly with not-so-successful outcomes. Hence the talk and explore similarities and links between obesity and atherosclerosis exist in order to treat and prevent both conditions. Interestingly, recent studies have shown that activation of beige or brown adipocytes, and, consequently, increase in energy consumption can reduce the mass of fat tissue in the body and potentially reduce the inflammation of body fat (3). There is evidence that the function of brown and beige fat tissue is regulated by immune processes (4). The current strategy is to increase the energy consumption of brown adipose tissue (BAT) in order to reduce excessive energy depots in obesity. Interventions in immune pathways could precede the development of new strategies to increase the activity of brown and beige adipose tissue as a therapeutic target for obesity.

### ***The role of low grade inflammation in pathogenesis of obesity and associated comorbidity***

The first proof that inflammation is important in the pathogenesis of obesity and the resulting metabolic dysfunction was provided by Hotamasligil et al (5). The pro-inflammatory cytokine tumor necrosis factor- $\alpha$  (TNF) was found to be present in WAT and correlated with insulin resistance in humans and mice (5). It became evident that the infiltration of proinflammatory macrophages in WAT plays a central role in the inflammatory response as main source of TNF (6). In WAT of lean mice, 10-15% of the cells are macrophages, whereas obese WAT contains 45-60% macrophages (6). Resident macrophages in lean WAT have a predominant anti-inflammatory phenotype, whereas in obesity, inflammatory monocytes are recruited to WAT, where they differentiate, acquire a proinflammatory or M1 phenotype and form the majority of macrophages (7). Antiinflammatory or M2 macrophages depend on the cytokines IL-4 and IL-13 and require STAT6 to maintain their alternative activation condition (8). Other myeloid cells that play a role in WAT include neutrophils and eosinophils. Neutrophils are very short-lived cells that are already present in WAT within 3 days of HFD (9). In contrast, the number of eosinophils is inversely correlated with adiposity, and exhaustion of eosinophils in mice results in increased body weight, glucose intolerance and insulin resistance (10). Both type 2 innate lymphoid cells (ILC2s) and eosinophils have only recently been shown to be an important cell population in WAT and are a predominant source of IL-4 and IL-13, the cytokines required for the induction of M2 macrophage polarization (10). Lymphoid cell infiltration in WAT T cells are also a component of the repertoire of immune cells found in WAT. Ten percent of the stromal vascular fraction of lean WAT consists of T cells. A large part of these are CD4<sup>+</sup> T-helper cells, of which approximately 50% are regulatory T cells (Tregs). In humans, the number of T cells in WAT correlates with BMI (11). In mice, the amount of T cells in WAT increases within 2 weeks of HFD. There are only a few CD8<sup>+</sup> cytotoxic T cells and CD4<sup>+</sup> effector T cells in lean WAT, but both populations increase drastically in an obese state, whereas CD4<sup>+</sup> Tregs decrease (12). Similarly, as the ratio between M1 and M2 macrophages increases in obese WAT, the Th1 and Th2 T cell ratio does as well. This results in a decrease in Th2 derived cytokines such as IL-4 and IL-13, thereby reducing M2 macrophage polarization. An increase in Th1 T cells and cytotoxic T cells results in excessive secretion of TNF and IFN $\gamma$ , which polarizes macrophages to a pro-inflammatory state, resulting in increased inflammation in obese WAT (10,12). The chronic low-grade inflammation in obese WAT also includes the recruitment of B cells, natural killer (NK) cells and mast cells (13). NK cells are activated by recognition of lipid antigens and mast cells contain granules that can release

a variety of mediators, including histamine, serotonin and cytokines, which also promote recruitment of inflammatory cell types (13).

### ***The role of brown and beige adipose tissue***

In contrast to the established role of the different immune cells in WAT, the contribution of the immune system to the development, function and activity of BAT is still largely unknown. However, we do know that obese individuals have a decreased amount of active BAT, based on FDG uptake, which is related to their low-grade inflammatory state. Moreover, an inactive brown adipocyte accumulates lipids, similar to a white adipocyte and ablation of noradrenergic input by selective sympathetic denervation of BAT indeed results in a ‘whitened’ appearance of brown adipocytes with large intracellular vacuoles. Since the recruitment of macrophages into WAT is correlated with lipolysis of stored triglycerides (14), it is likely that release of fatty acids also induces recruitment of immune cells in BAT. However, whether this is indeed the case is still unknown. In diet-induced obesity (DIO), thermoneutral housing leads to an additive increase in inflammation in white adipose tissue and in the vasculature compared to normal housing conditions. Although not causing increased insulin resistance, the increase in vascular inflammation does cause enhanced progression of atherosclerosis (14), indicating that BAT protects against obesity-induced atherosclerosis. In another study, a similar phenomenon was observed. Immune compromised nude mice experience cold stress when housed at 23°C which modulates energy and body weight homeostasis and are therefore protected from DIO. However, at thermoneutrality (33°C), they do develop DIO with increased adiposity, hepatic triglyceride accumulation, increased inflammatory markers and glucose intolerance (15), showing that BAT activity protects against metabolic disarray and adipose tissue inflammation. Besides environmental temperature, other incentives such as the biological clock (16), hormones (17) and food intake not only modulate energy expenditure via the hypothalamus but also affect inflammation. For instance, time-restricted feeding of HFD for 8 hours per day increases BAT activity and reduces adipocyte hypertrophy and inflammation in WAT compared to ad libitum HFD-fed mice (18). Gut hormones such as GLP-1 mediate effects on food intake, energy expenditure and inflammation. GLP-1 receptor signaling activates BAT and promotes beiging of WAT (19,20) while GLP-1 also reduces macrophage infiltration and inflammatory signalling in white adipocytes and macrophages (21,22). Other hormonal changes such as menopause also affect energy metabolism. Estradiol inhibits AMPK in the hypothalamus, which activates thermogenesis in BAT (18). Indeed, ovariectomised mice with reduced estradiol levels gain more weight than sham operated mice and have reduced energy expenditure and increased WAT inflammation (23).

### *Clinical implications*

Excessive energy intake results in increased storage of lipids in both white and brown adipocytes, which challenges the function of these cells. Immune cells and signals in WAT (white adipose tissue) and BAT are indispensable for the homeostasis of the tissue and contribute to the efflux of lipids stored in white adipocytes and to high rates of oxidation in brown and beige adipocytes. Immune cells, including eosinophils and alternatively activated macrophages, have regulatory roles in metabolic homeostasis of both WAT and BAT. Research to identify immunological players is ongoing. If the mechanism is unravelled in detail, immune regulation is an intriguing therapeutic target in increasing energy expenditure to reduce weight gain. Notably, the numbers of immune cells in lean as well as obese BAT are much lower than in WAT, indicating that BAT is relatively more resistant to diet induced inflammation, but increased tissue inflammation in BAT does occur upon a positive energy balance. The challenge is to identify the metabolic crosstalk between immune cells and brown, beige and white adipocytes and the order of events that occur during obesity development. Important questions to further address include: What are the immune regulatory effector molecules that are secreted by brown adipocytes (or a pre-beige adipocyte or a white-wanting-to-become beige adipocyte) to attract or regulate immune cells? How is BAT activity regulated in obesity? What is the role of the sympathetic nerve system? And how does BAT activity change during aging? The presence of BAT in humans and the potential activation of resident BAT or induction of beige adipocytes in WAT is an interesting target to treat or even prevent obesity-related disorders. Cold is still by far the strongest sympathetic signal to activate BAT, but the quest for identifying biochemical and immunological pathways that are responsible for BAT activation, and thereby can bypass prolonged cold exposure, is ongoing. The recent finding on the role for immune cells in brown and beige adipocyte development and physiology harbours a great potential to increase BAT activity and beneficially alter energy metabolism by interfering in immune responses.

### *Literatura*

1. Berrington de Gonzalez A, Hartge P et al. Body-Mass Index and Mortality among 1.46 Million White Adults. *NEJM* 2010; 363: 2211–2219.
2. Rocha V Z, Libby P. Obesity, inflammation, and atherosclerosis. *Nat Rev Cardiol* 2009; 6, 399–409.
3. Yoneshiro T, Aita S, Matsushita M et al. Recruited brown adipose tissue as an antiobesity agent in humans. *J Clin Invest* 2013; 123: 3404–3408.
4. Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *Clin Invest* 2011; 121: 2111–2117.

5. Hotamisligil G, Shargill N, Spiegelman B. Adipose expression of tumor necrosis factor- $\alpha$ : direct role in obesity-linked insulin resistance. *Science* 1993; 259: 87–91.
6. Gregor MF, Hotamisligil GS. Inflammatory Mechanisms in Obesity. *Ann Rev Immunol* 2011; 29: 415–445.
7. Chawla A, Nguyen KD, Goh YPS. Macrophage-mediated inflammation in metabolic disease. *Nature Rev Immunol* 2011; 11: 738–749.
8. Brestoff JR, Kim BS, Saenz SA et al. Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature* 2015; 519: 242–246.
9. Talukdar S, Oh DY, Bandyopadhyay G et al. Neutrophils mediate insulin resistance in mice fed a high-fat diet through secreted elastase. *Nature Med* 2012; 18: 1407–1412.
10. Wu D, Molofsky AB, Liang H-E, Ricardo-Gonzalez RR, Jouihan HA, Bando JK, Chawla A, Locksley RM. Eosinophils Sustain Adipose Alternatively Activated Macrophages Associated with Glucose Homeostasis. *Science* 2011; 332: 243–247.
11. Chatzigeorgiou A, Karalis KP, Bornstein SR, Chavakis T. Lymphocytes in obesity related adipose tissue inflammation. *Diabetologia* 2012; 55: 2583–2592.
12. Winer S, Winer DA. The adaptive immune system as a fundamental regulator of adipose tissue inflammation and insulin resistance. *Immun Cell Biol* 2012; 90: 755–762.
13. Liu J, Divoux A, Sun J, Zhang J, Clement K, Glickman JN, Sukhova GK, Wolters PJ, Du J, Gorgun CZ, Doria A, Libby P, Blumberg RS, Kahn BB, Hotamisligil GS, Shi G-P. Genetic deficiency and pharmacological stabilization of mast cells reduce diet-induced obesity and diabetes in mice. *Nature Med* 2009; 15: 940–945.
14. Tian Xiao Y, Ganeshan K, Hong C, Nguyen Khoa D, Qiu Y, Kim J, Tangirala Rajendra K, Tonotonoz P, Chawla A. Thermoneutral Housing Accelerates Metabolic Inflammation to Potentiate Atherosclerosis but Not Insulin Resistance. *Cell Metabol* 2016; 23: 165–178.
15. Stemmer K, Kotzbeck P, Zani F, Bauer M, Neff C, Muller TD, Pfluger PT, Seeley RJ, Divanovic S. Thermoneutral housing is a critical factor for immune function and diet-induced obesity in C57BL/6 nude mice. *Int Obes* 2015; 39: 791–797.
16. Orozco-Solis R, Aguilar-Arnal L, Murakami M, Peruquetti R, Ramadori G, Coppari R, Sassone-Corsi P. The Circadian Clock in the Ventromedial Hypothalamus Controls Cyclic Energy Expenditure. *Cell Metab* 2016; 23: 467–478.
17. Martínez de Morentin Pablo B, González-García I, Martins L, Lage R, Fernández-Mallo D, Martínez-Sánchez N, Ruíz-Pino F, Liu J, Morgan Donald A, Pinilla L, Gallego R, Saha Asish K, Kalsbeek A, Fliers E, Bisschop Peter H, Diéguez C, Nogueiras R, Rahmouni K, Tena-Sempere M, López M. Estradiol Regulates Brown Adipose Tissue Thermogenesis via Hypothalamic AMPK. *Cell Metab* 2014; 20: 41–53.
18. Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong Eric A, Gill S, Leblanc M, Chaix A, Joens M, Fitzpatrick James AJ, Ellisman Mark H, Panda S. Time-Restricted Feeding without Reducing Caloric Intake Prevents Metabolic Diseases in Mice Fed a High-Fat Diet. *Cell Metab* 2012; 15: 848–860.
19. Kooijman S, Wang Y, Parlevliet ET, Boon MR, Edelschaap D, Snaterse G, Pijl H, Romijn JA, Rensen PCN. Central GLP-1 receptor signalling accelerates plasma clearance

- of triacylglycerol and glucose by activating brown adipose tissue in mice. *Diabetologia* 2015; 58: 2637–2646.
20. Beiroa D, Imbernon M, Gallego R, Senra A, Herranz D, Villarroya F, Serrano M, Fernø J, Salvador J, Escalada J, Dieguez C, Lopez M, Frühbeck G, Nogueiras R. GLP-1 Agonism Stimulates Brown Adipose Tissue Thermogenesis and Browning Through Hypothalamic AMPK. *Diabetes* 2014; 63: 3346.
  21. Lee YS, Park MS, Choung JS, Kim SS, Oh HH, Choi CS, Ha SY, Kang Y, Kim Y, Jun HS. Glucagon-like peptide-1 inhibits adipose tissue macrophage infiltration and inflammation in an obese mouse model of diabetes. *Diabetologia* 2012; 55: 2456–2468.
  22. Wang Y, Parlevliet ET, Geerling JJ, van der Tuin SJL, Zhang H, Bieghs V, Jawad AHM, Shiri-Sverdlov R, Bot I, de Jager SCA, Havekes LM, Romijn JA, Willems van Dijk K, Rensen PCN. Exendin-4 decreases liver inflammation and atherosclerosis development simultaneously by reducing macrophage infiltration. *British J Pharmacol* 2014; 171: 723–734.
  23. Rogers NH, Perfield JW, Strissel KJ, Obin MS, Greenberg AS. Reduced Energy Expenditure and Increased Inflammation Are Early Events in the Development of Ovariectomy-Induced Obesity. *Endocrinology* 2009; 150: 2161–2168.