Mihajlovski Mirjana¹, Perišić Zlatko¹, Raspopović Miloš¹,
Petrović Nataša²

SERUMSKI BIOMARKERI SEPSE

Sažetak: Sepsa je imunološki odgovor na sistemsku infekciju i širom sveta predstavlja sve veći problem u pogledu hospitalnog morbiditeta i mortaliteta. Brzi molekularni testovi, razvijeni da odgovore na potrebu brze i adekvatne dijagnostike, nisu uvek kadri da pruže jednoznačan odgovor. Izazovi sa kojim se suočavaju istraživači relevantnih biomarkera sepse leže u nedostatku jedinstvenog protokola i metodologije i nivoa dokaza. Pod sepsom se podrazumeva ceo skup poremećaja funkcije vitalnih organa i organskih sistema, koji dovode do multiorganske disfunkcije, otkazivanja funkcije osovine vitalnih organa i na kraju do smrti. Jedan od osnovnih problema u sepsi predstavlja činjenica da vrlo često nedostaju tipične i rane kliničke manifestacije. Za sada, ipak, zlatni standard dijagnoze sepse predstavlja tradicionalna upotreba mikrobnih kultura krvi (hemokultura).

Glavno ograničenje upotrebne vrednosti kulture krvi je u dužini vremena potrebnom za detekciju uzročnika sepse.

Biomarkeri bakterijske sepse: C-reaktivni protein je opšti protein reaktanta akutne faze **Prokalcitonin** je prekursor hormona kalcitonina i moćan je biomarker za bakterijske infekcije, **serumski amiloid A (SAA)** je važan apolipoprotein.

Biomarkeri za infekcije izazvane gljivicama: Antitela mannan i anati-mannan koriste se za otkrivanje gljivičnih infekcija zbog prisustva manna u ćelijskim zidovima invazivnih gljivičnih patogena. Glavni nedostatak dijagnostike upotrebom samo Mannan i Antimannan antitela je visoka stopa lažno pozitivnih i negativnih rezultata.

Biomarkeri za virusne infekcije: interferon Gamma inducible protein 10 (IP-10) proinflamatorni hemokin je važan biomarker za dijagnostiku virusnih infekcija.

Drugi biomarkeri (na raspolaganju ili u istraživanju): IL-10, proteinski vezani lipopolisaharid,

¹ Mihajlovski Mirjana, Klinika za urgentnu hirurgiju Urgentnog centra KCS

² Klinika za anesteziologiju i reanimatologiju KCS

rastvorljivi pokretački receptor (Soluble triggering receptor), tool-like receptors (TLRs), TLR-2 i neutrofilni CD64 receptori, CD163, mikro RNA (miRNA), SNP polimorfizam sa jednim nukleotidom, TNF alfa, IL-1 kinaza.

Dobar biomarker može da ima ključni značaj u dijagnostici i diferencijalnoj dijagnostici i ishodu sepse.

Ključne reči: sepsa, rana dijagnoza, biomarkeri, mortalitet, morbiditet

Uvod

Sepsa je imunološki odgovor na sistemsku infekciju i širom sveta predstavlja sve veći problem u pogledu hospitalnog morbiditeta i mortaliteta. Septično stanje zahteva trenutnu agresivnu terapiju, što podrazumeva pravovremenu, što raniju tačnu dijagnostiku.¹ Brzi molekularni testovi, razvijeni da odgovore na ovu potrebu, nisu uvek kadri da pruže jednoznačan odgovor. Najčešće se koriste serumski biomarkeri, sa različitom dijagnostičkom tačnošću. Najkorišćeniji biomarkeri su C reaktivni protein i nivo prokalcitonina (PCT); takođe je, mada nešto ređe, zastupljen nivo serumskog amiloida.² Već decenijama traje potraga za, po mogućnosti, pojedinačnim biomarkerom sepse. Za sada se ispostavlja da ne postoji idealan biomarker, pa se istraživanja nastavljaju u smislu pronalaženja grupe biomarkera, koji bi bili potentni u dijagnostici i praćenju toka septičnog stanja. Izazovi sa kojim se suočavaju istraživači relevantnih biomarkera sepse leže u nedostatku jedinstvenog protokola i metodologije, standardizacije ispitivanja i nivoa dokaza. Otkrivanje pouzdanog biomarkera ili grupe biomarkera imalo bi ogroman dijagnostički i prognostički značaj u lečenju sepse.²

Sepsa se smatra drugim najvažnijim uzrokom oboljevanja i smrtnosti u hospitalnim uslovima. Pod sepsom se ne podrazumeva samo infekcija krvi (pozitivna hemokultura), već ceo skup poremećaja funkcije vitalnih organa i organskih sistema, koji dovode do multiorganske disfunkcije, otkazivanja funkcije osovine vitalnih organa i na kraju do smrti. Jedan od osnovnih problema u sepsi predstavlja činjenica da vrlo često nedostaju tipične i rane kliničke manifestacije, tako da je potrebno pronalaženje markera ili seta markera koji bi bili potentni za pretkliničku dijagnozu sepse. Imunološki, sepsa predstavlja posledicu poremećaja regulacije urođenog imunog odgovora. Biomarkeri su stoga važni signalizatori prisustva i ozbiljnosti sepse u pretkliničkoj fazi, te bi njihovo protokolisanje imalo ogromnu prognostičku vrednost i značaj.^{1,2}

Veoma je važno napomenuti da sepsa nije jedno oboljenje i samo fiziološka manifestacija sistemskog imunog odgovora. Većina referentnih studija i publikacija definiše sepsu kao sistemski upalni odgovor na noksu koja je izaziva (primarna infekcija, trauma, oboljenje...). Za sada ne postoji jednoznačni biomarker za sepsu,

već se vrednosti serumskih nivoa poznatih biomarkera tumače u kontekstu simptomatologije. Uobičajena terapija septičnih pacijenata teži, s jedne strane, lečenju sistemske infekcije korišćenjem antimikrobnih sredstava širokog spektra, a s druge, sprečavanju otkazivanja vitalnih organa i organskih sistema. Za sada, ipak, zlatni standard dijagnoze sepse predstavlja tradicionalna upotreba mikrobnih kultura krvi (hemokultura).

Glavno ograničenje upotrebne vrednosti kulture krvi je u dužini vremena potrebnom za detekciju uzročnika sepse (za rezultat Bactek metode potrebno je čekati oko 48 h). Za to vreme sprovodi se terapija *ex iuvantibus*. Ova činjenica ilustruje važnost i potrebu za otkrivanjem ranih serumskih biomarkera. Treba napomenuti, takođe, da i kulture mogu biti negativne pod određenim okolnostima, npr. kada se radi o spororastućim izazivačima ili potentnim izazivačima u niskoj koncentraciji.

Dakle, u svetlu ovog problema razvijene su alternativne dijagnostičke metode pomoću metoda upotrebljivanih u molekularnoj biologiji da bi dijagnostika sepse na bazi potencijalno otkrivenih markera mogla da se radi brzo i automatizovano. Ovi testovi su u osnovi ELISA testovi, citometrijska detekcija specifičnih markerskih antigena, imunokolorimetrijski testovi, FISH metodologije i PCR i pojačana PCR tehnika. Dakle, sve metode imaju kao osnovni ishod detekciju proteina markera.³

Paralelno sa razvojem biomarkera, koji omogućava brže i osetljivije dijagnostičke metode za detekciju samog infektivnog agensa, razvijaju se istraživanja posleđičnih promena serumskih proteina, koji bi, što se brže menjaju I što im je koncentracija veća, bili bolji kandidati za serumske biomarkere sepse.

Biomarker se najbolje definiše kao merljiva komponenta, obično proteina, koja se može objektivno evaluirati kao indikator normalnog biološkog odgovora ili patološkog procesa. Ovaj rad je fokusiran na širok spektar biomarkera koji se koriste na bazi dijagnostičke validnosti (DA-diagnostic accuracy), koja se zasniva na osetljivosti, specifičnosti i prediktivnoj vrednosti.

Da bi bio klinički relevantan, biomarkeri treba da imaju veoma visok DA, odnosno osetljivost, specifičnost i prediktivne vrednosti (kako pozitivne – positive predictive values, PPV, tako i negativne – negative predictive values, NPV). U nekim studijama publikovani su načini izračunavanja koeficijenta verovatnoće, koji bi mogli da determinišu rezultat koji je posledica oboljenja ili je, pak, do njega dovela slučajnost. Vrednosti ovih koeficijenata su pomogle da se odrede granice dijagnostičkih preseka. Pošto je sepsa stanje uzrokovano bakterijskim, gljivičnim ili virusnim agensima, čijom kolonizacijom nastaje hemoinfekcija, takođe treba razmišljati o biomarkerima za specifične patogene, naročito kada postoji klinička sumnja na određeni agens. Za sada ne postoje apsolutni i specifični pojedinačni marker (koji zadovoljavaju ranije navedene kriterijume), ali se neki merljivi proteini široko koriste npr. za određivanje klase izazivača sepse, da bi se započela što više ciljana terapija antimikrobnim lekovima pre egzaktnog izdvajanja izazivača.⁴

Biomarkeri bakterijske seps

C-reaktivni protein

Ovo je opšti protein reaktanta akutne faze, čija koncentracija kao odgovor na infekciju i prateći inflamatorni proces u krvi raste do 1.000 puta.

Osetljivosti za CRP nalazi se u rasponu 30–97% sa vrednostima specifičnosti 75–100%, a PPV 31–100%. Upotreba različitih testova uzrokovala je varijaciju rezultata, pošto je svakom testu dodeljeno jedinstveno funkcionalno ograničenje. Nivoi CRP u serumu takođe variraju sa godinama starosti i kod zdravih pojedinaca. Granična vrednost od 10 mg/L koristi se za pacijente >65 godina starosti, dok se granična vrednost od 5 mg/L koristi za odrasle i novorođenčad od 3 do 7 dana starosti. Nivo CRP-a je viši kod starije, nego kod mlade dece.^{3,5} Studije su pokazale da je CRP vrlo osetljiv i veoma specifičan za infekciju. Pomoću CRP koncentracije u serumu mogu se razlikovati gram-pozitivne i gram-negativne, kao i bakterijske i gljivične infekcije, gde su nivoi iznad 100 mg/L specifični za bakterijske, a niži nivoi označavaju gljivične infekcije. Nivo koncentracije CRP takođe se može koristiti kao prognostički marker za nehospitalnu pneumoniju (stečenu u zajednici). Koncentracija nivoa 110mg/L sugeriše potrebu za hospitalizacijom. Međutim, dijagnostika virusne infekcije je nepouzdana.⁶

Prokalcitonin

PCT je prekursor hormona kalcitonina i moćan je biomarker za bakterijske infekcije. U serumu zdravih pojedinaca beleži se niska koncentracija od oko 0,33 ng/ml, koja se povećava oko 1.000 puta u uslovima inflamacije. Koncentracija raste u roku 2–4 sata od infekcije i dostiže vrhunac za 6–8 sati. Stalni povećani nivoi ukazuju na kontinuirano prisustvo bakterijske infekcije. Brza regulacija i održavanje seruma čini ga jedinstvenim biomarkerom. Kako je ovaj molekul veoma stabilan, on ostaje prisutan u uzorcima krvi i prilikom zamrzavanja, čak i nakon dugotrajnog skladištenja.⁷

PCT ima visoku DA i osetljivosti od oko 75–100, a vrednosti specifičnosti 70–100%. PPV vrednosti su 55–100. Većina referentnih studija predlaže CV PCT nivoa od 0,5 ng/mL. Većina autora smatra PCT egzaktnijim inflamatornim markerom od ostalih proinflamatornih markera. PCT, takođe, ima veoma prediktivnu prognostičku vrednost i njegovi nivoi su distinktivni za gljivične i virusne infekcije. Tokom virusnih infekcija vrednosti nivoa serumskih koncentracija PCT ostaju niske, često i u normalnim granicama. PCT se takođe koristi u laboratorijskoj diferencijalnoj dijagnostici bakterijskog i virusnog meningitisa. Suprotno tome, gljivične infekcije imaju tendenciju da izazivaju blago povišenje koncentracije PCT u poređenju sa nivoima koji se vide kod bakterijskih infekcija.

Istraživanja invazivne aspergiloze i invazivne kandidijaze su pokazale da postoji značajno viši nivo PCT kod pacijenata sa bakterijskom, nego sa gljivičnom sepsom.^{5,8}

PCT se takođe pokazao kao potencijalni biomarker za pneumonije, infekcije donjih disajnih puteva, abdominalnu sepsu, urosepsu, infarkt miokarda i kao biomarker za praćenje uspešnosti antibiotske terapije, naročito kada je data spekulativno, a ne na bazi antibiograma.

Serumski amiloid A (SAA)

Amiloid A je važan apolipoprotein, za koji u referentnoj literaturi postoje ozbiljni dokazi da ima potencijalnu vrednost u ranoj dijagnostici sepse. Nivo SAA je oko 1.000 puta višim nakon 8–24 sata od početka sepse. U poređenju sa nivoima CRP, izveštava se da nivo SAA raste brže i više nakon početka sepse i duže ostaje na višim relativnim vrednostima. Nivoi SAA u serumu ispod 15 mg/L za starije od 65 godina i manje od 10 mg/L za odrasle i novorođenčad u dobi 3–7 dana su granične vrednosti za utvrđivanje dijagnostičkih kriterijuma. Iako se SAA uglavnom koristi kao marker za bakterijske infekcije, u koncentraciji iznad 10 mg/L pokazao se potentnim i u dijagnostici virusnih infekcija. SAA nema veliku osetljivost u poređenju sa nekim markerima, pa je stoga vrednost njegove kliničke primene još uvek diskutabilna i, možda, precenjena od strane nekih autora.⁸

Biomarkeri za infekcije izazvane gljivicama

Antitela mannan i anitimannan koriste se za otkrivanje gljivičnih infekcija zbog prisustva manna u ćelijskim zidovima invazivnih gljivičnih patogena. Nivoi mannana i anitimannana ostaju povišeni u uslovima infekcije poput aspergiloze i kandidijaze, što ih čini korisnim markerima kod gljivične sepse. Postoji puno varijacija u DA, PPV vrednosti osetljivosti i specifičnosti, zbog različitih protokola kako u ispitivanju tako i u upotrebi markera u dijagnostici.⁹ Glavni nedostatak dijagnostike upotrebom samo Mannan i Antimannan antitela je visoka stopa lažno pozitivnih i negativnih rezultata testiranja. Ova činjenica zahteva da se upotreba ovih testova kombinuje sa drugim dijagnostičkim testovima, kao pr. sa beta-D glukan testovima, jer je beta-D glukan veoma osetljiv i specifičan za invazivnu mikozu, a nije specifičan za vrste.¹⁰

Biomarkeri za virusne infekcije

Interferon Gamma inducible protein 10 (IP-10)

IP-10 proinflamatorni hemokin je važan biomarker za dijagnostiku virusnih infekcija zbog njegove uloge u odgovoru domaćina na viremiju. Studija o bakterijskim infekcijama kod novorođenčadi sa vrlo niskom težinom na rođenju pokazala je da je IP-10 dobar biomarker tokom početnih merenja, kod koga se beleži značajno i

merljivo povišen nivo kod zaraženih pacijenata, inicijalno i tokom prva 24 sata, bez „preklapanja” ovih nivoa. Pokazano je da su nivoi IP-10 u korelaciji sa težinom sepse. IP-10 se oslobađa kao odgovor na viruse poput rinovirusa, respiratornog sincitijalnog virusa, hepatitisa B i virusa gripa, a zatim i hepatitisa C i H5N1. Takođe se pokazalo da predstavlja vrlo dobar prognostički marker kod pacijenata sa hepatitisom C.¹¹

Drugi biomarkeri (na raspolaganju ili u istraživanju)

IL-10

IL-10 je u studijama označen kao koristan biomarker sepse, posebno u dijagnostici u ranom (prekliničkom) i kasnom (ranom kliničkom) ispoljavanju sepse novorođenčadi. Vrednosti CV više od 17,3 pg/mL procenjene su kao “cut-off” vrednosti DA. Ovaj marker pokazuje osetljivost i specifičnost u rasponu od 80%. Glavni problem u njegovoj upotrebi je činjenica da cirkulišući citokini imaju vrlo kratak poluživot. Ovo dovodi do značajnog broja lažno negativnih rezultata, što ograničava njihovu vrednost DA, pa tako i upotrebnu vrednost.¹²

Proteinski vezani lipopolisaharid

Proteinski vezani lipopolisaharid takođe je koristan biomarker, specifičan za bakterijske infekcije. Poznato je da se nivo ovog proteina u serumu povećava tokom bakterijskih i gljivičnih, ali ne i kod virusnih infekcija.¹³

Rastvorljivi pokretački receptor (Soluble triggering receptor)

Pokretački receptori, koji pokazuju ekspresiju na mijeloidnim ćelijama – 1 TREM, su grupa receptora ćelijske površine koji pripadaju superfamiliji imunoglobulina. TREM-1 se pojavljuje uglavnom na makrofazima i neutrofilima i identifikovan je kao pojačavač imunog odgovora koji snažno osnažuje aktivaciju leukocita u prisustvu mikrobnih proizvoda. Poslednji rezultati sugeriraju da sTREM-1 može biti oslobođen širokim spektrom inflamatornih podražaja. Prvi obećavajući rezultat upotrebe sTREM-1 u plazmi u dijagnostici kod sepse kod pacijenata u Jedinici intenzivnog lečenja (JIL) pokazao je da bi sTREM-1 mogao biti odavno traženi savršen dijagnostički biomarker sepse. Rezultati se kreću u rasponu tačnosti od gotovo 100% (ROC-AUC od 0,97) za dijagnozu sepse u pacijenata koji se nalaze u JIL.¹⁴

Tool-like receptori (TLRs)

Rezultati referentnih studija sugerišu da TLR i drugi receptori koji se nalaze na ćelijskoj membrane mogu biti pobuđeni od strane neutrofila i imaju dinamičku modulaciju u toku sepse.

Pojavljjujući se na makrofazima, dendritskim ćelijama i ostalim ćelijama, TLR igraju važnu ulogu u urođenim imunskim odgovorima na infekciju unakrsnom reakcijom i prepoznavanjem specifičnih bakterijskih antigena. Ekspresija TLR gena u mononuklearnim ćelijama naglo opada u težim oblicima sepse. TLR se dinamički modifikuju u raznim fazama sepse. TLR-4 je specifičan za prepoznavanje LPS, dok je TLR-2 koristan za prepoznavanje Gram-pozitivnih bakterijskih komponenti.

TLR-2 i neutrofilni CD64 receptori

(TLR-2) i neutrofilni CD64 receptori su dva važna biomarkera prisutna na ćelijskim površinama. Tokom infekcije pojačava se odgovor na stimulus na površini monocita i neutrofila od strane oba ova biomarkera.¹⁵ Ovaj odgovor je značajno pojačan kako tokom bakterijskih tako i virusnih infekcija. Vrednosti su značajno povišene u uslovima infekcije poput gripa A i B, respiratorno sincitijalnog virusa ili mumpsa, virusa herpes zoster, citomegalovirusne infekcije, što ukazuje da TLR-2 može biti potencijalni biomarker za virusne infekcije.^{5,15} Za sada se CD64 koristi kao marker bakterijskih infekcija. Na ekspresiju CD64 ne utiču blagi inflamatorni procesi i nema razlike u nivoima kod sistemskih i lokalnih bakterijskih infekcija, vrednosti DA, vrednosti osetljivosti i specifičnosti kod PPV u rasponu 71–100%, što ga čini značajnim biomarkerom.

CD163 je transmembranski molekul koji se nalazi na membrani mononuklearnih fagocita. Budući da je specifičan receptor za uklanjanje hema iz hemoglobina u organizmu, vrlo je specifičan za identifikaciju kompleksa haemoglobin-haptoglobin. CD163 reguliše ekspresiju antiinflamatornih molekula, poput citokina IL-10 i heme-oksigenaze-1. Rastvorljivi CD163 potiče iz molekula „zguljenih” sa ovih površina fagocita i veoma su dobri markeri sepse i teške sepse. Rezultati istraživanja pokazuju da su superiorniji od nivoa PCT i CRP.

Mikro RNA (miRNA) je vrsta endogenih nekodirajućih malih RNK dužine oko 22 nukleotida i igraju važnu ulogu u posttranskripcionoj regulaciji ekspresije gena. Značajna količina miRNA registrovana je ekstraćelijski, unutar različitih telesnih tečnosti. Slobodne miRNA u ćelijama su stabilne čak i u teškim uslovima. MiR-150 je naznačen kao biomarker za sepsu, a odnos plazme miR-150/interleukin 18 ispituje se kao pokazatelj težine sepse.

SNP Polimorfizam sa jednim nukleotidom

Polimorfizam pojedinačnih nukleotida (SNP) najčešći je izvor genetskih varijacija u humanom genomu i obuhvata oko 90% svih DNK polimorfizama.

Poznato je da je C-reaktivni protein (CRP) važan biomarker sepse. Nekoliko nukleotidnih polimorfizama (SNP) gena CRP može odrediti nivo CRP u plazmi, pa tako SNP postaje marker sepse. On se, takođe, istražuje i kao faktor rizika za mnoge bolesti (maligne, artritis i dijabetes). Najviše se upotrebljava kao marker infantilne sepse. Ne zna se, međutim, da li je njegov nivo povezan sa ishodom infantilne sepse.

TNF alfa – Faktor tumorske nekroze alfa

Ovo je pleiotropni citokin nastao aktivacijom makrofaga, koji ima ključnu ulogu kako u inflamatornom odgovoru tako i u progresiji autoimunih bolesti. Uočena je povezanost sepse sa ekspresijom genotipa TNF-2.¹⁶

IL-1 kinaza, povezana sa receptorima interleukina 1, važna je za TLR2 i TLR4 indukovanu aktivaciju kappa beta faktora jezgra, ključnog u regulaciji transkripcije mnogih proinflamatornih citokina u septičnim stanjima.

Sepsa je sve značajniji uzrok morbiditeta i mortaliteta kritično obolelih pacijenata u jedinicama intenzivnog lečenja. Ovo stanje zahteva brzo, ako je moguće pretkliničko postavljanje dijagnoze, kao i brzo ordiniranje adekvatne terapije, što značaj biomarkera stavlja u prvi plan. Dobar biomarker može da ima ključni značaj u dijagnostici i diferencijalnoj dijagnostici i ishodu sepse.¹⁶

Literatura

1. Kramer MS, Demissie K, Yang H, Platt RW, Sauve R, Liston R. The contribution of mild and moderate preterm birth to infant mortality. Fetal and Infant Health Study Group of the Canadian Perinatal Surveillance System. *JAMA*. 2000; 284: 843–9.
2. Vincent JL, Sakr Y, Sprung CL, et al. Sepsis occurrence in acutely ill patients investigators: sepsis in European intensive care units: results of the SOAP Study. *Crit Care Med*. 2006; 34: 344–53.
3. Kaus IAM, Grion CMC, Cardoso LTQ *et al*. The epidemiology of sepsis in a Brazilian teaching hospital. *Brazilian J. Infect. Dis*. 2010; 14(3), 264–270.
4. Wang P, Yang Z, He Y, Shu C. Pitfalls in the rapid diagnosis of positive blood culture. *Rev. Med. Microbiol*. 2010; 21(3), 39–43.
5. Papachristou GI, Clermont G, Sharma A *et al*. Risk and markers of severe acute pancreatitis. *Gastroenterol Clin N Am* 2007; 36: 277–96.
6. Rego MAC, Martinez FE, Elias J, Mussi-Pinhata MM. Diagnostic value of interleukin-6 and C-reactive protein on early onset bacterial infection in preterm neonates with respiratory distress. *J. Perinat. Med*. 2010; 38(5), 527–533.

7. Morris AC, Simpson AJ, Walsh TS. Hyperinflammation and mediators of immunosuppression in critical illness. In: Vincent JL, editor. Annual update in intensive care and emergency medicine. Berlin Heidelberg: Springer- Verlag; 2013; 135–44.
8. Cetinkaya M, Ozkan H, Koksall N, Celebi S, Hacimustafaoglu M. Comparison of serum amyloid A concentrations with those of C-reactive protein and procalcitonin in diagnosis and follow-up of neonatal sepsis in premature infants. *J. Perinatol.* 2009; 29(3), 225–231.
9. Méan M, Marchetti O, Calandra T. Bench-to-bedside review: Candida infections in the intensive care unit. *Crit Care.* 2008; 12: 204. doi: 10.1186/cc6212.
10. Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, Moreno R, Lipman J, Gomersall C, Sakr Y, et al. EPIC II Group of Investigators International study of the prevalence and outcomes of infection in intensive care units. *JAMA.* 2009; 302: 2323–9. doi: 10.1001/jama.2009
11. Grebely J, Feld JJ, Applegate T, Matthews GV, Hellard M, Sherker A, Petoumenos K, Zang G, Shaw I, Yeung B, George J, Teutsch S, Kaldor JM, Cherepanov V, Bruneau J, Shoukry NH, Lloyd AR, Dore GJ. Plasma interferon-gamma-inducible protein-10 (IP-10) levels during acute hepatitis C virus infection. *Hepatology.* 2013 Jun; 57(6): 2124–34. doi: 10.1002/hep.26263.
12. Mehr S, Doyle LW. Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J.* 2000; 19: 879–87.
13. Villar J, Perez-Mendez L, Espinosa E et al. Serum lipopolysaccharide binding protein levels predict severity of lung injury and mortality in patients with severe sepsis. *PLoS One* 2009; 4: e6818.
14. Barati M, Bashar FR, Shahrami R, et al. Soluble triggering receptor expressed on myeloid cells 1 and diagnosis of sepsis. *J Crit Care.* 2010; 25: 362. e1–6.
15. Tsujimoto H, Ono S, Efron PA, Scumpia PO, Moldawer LL, Mochizuki H. Role of Toll-like receptors in the development of sepsis. *Shock.* 2008 Mar; 29(3): 315–21. doi: 10.1097/SHK.0b013e318157ee55.
16. De JW, te VH, Prakken BJ, Kuis W, Rijkers GT. Simultaneous detection of 15 human cytokines in a single sample of stimulated peripheral blood mononuclear cells. *Clin Diagn Lab Immunol.* 2003; 10: 133–9.

Mihajlovski Mirjana¹, Perišić Zlatko¹, Raspopović Miloš¹,
Petrović Nataša²

SERUM BIOMARKERS OF SEPSIS

Abstract: Sepsis is an innate immunological response of the systemic inflammatory pathway to infection, and is growing problem worldwide in terms of mortality rates. Rapid molecular based tests have been developed to address this need but are not always accurate in this respect. The most commonly used biomarkers specially the serum markers are reviewed for their current uses and the diagnostic accuracies. Challenges that are faced during biomarker research lie in the lack of uniform protocol and methodology. Define sepsis to be a systemic inflammatory response to a bacteria, virus or fungi. In clinical setting several other physiological symptoms must be presented to properly diagnose sepsis. The gold standard of sepsis diagnosis has been traditionally the use of microbial cultures to identify the source of illness. The major limitation of using cultures is the length of the time required to develop cultures to identifiable quantities to detect sepsis. The gold standard of sepsis diagnosis has been traditionally the use of microbial cultures to identify the source of illness. The major limitation of using cultures is the length of the time required to develop cultures to identifiable quantities to detect sepsis.

Biomarkers for the Infections caused by Bacteria: C-reactive protein is the general acute phase reactant protein, **Procalcitonin** is a precursor of the hormone calcitonin and is another potent biomarker for bacterial infections, **Serum Amyloid A** is an important apolipoprotein,

Biomarkers for the Infections caused by Fungi. Mannan and Antimannan antibodies are used to detect the fungal infections due to the presence of mannan in the cell walls of the invasive fungal pathogens. The main disadvantage Mannan and Antimannan antibodies test alone is the high rate of false positives and negatives.

Biomarkers for Viral Infections: Interferon Gamma inducible protein 10 (IP-10)

¹ Mihajlovski Mirjana, Clinic for Emergency Surgery, Emergency Center, Clinical Center of Serbia

² Center for Anesthesiology and Resuscitation, Clinical Center of Serbia

IP-10 a proinflammatory chemokine is an important biomarker for diagnosing viral infections.

Other Biomarkers on Avenue: IL10, Lipopolysaccharide Binding Protein, Soluble triggering receptor, TLRs, Toll like Receptors 2 (TLR-2) and Neutrophil CD64 receptors, CD163, Micro RNAs, SNP, TNF alpha, IL-1.

Sepsis, as an increasing cause of mortality in the group of multiple injured and critically ill patients in the ICU, requires rapid diagnosis and treatment, and hence the importance of biomarkers is highly important.

Key words: sepsis, early diagnosis, biomarkers, morbidity, mortality

Introduction

Sepsis is an innate immunological response of the systemic inflammatory pathway to infection, and is growing problem worldwide in terms of mortality rates. Immediate treatment is always required which necessitates the need of early and accurate diagnosis.¹ Rapid molecular based tests have been developed to address this need but are not always accurate in this respect. The most commonly used biomarkers specially the serum markers are reviewed for their current uses and the diagnostic accuracies including the c reactive proteins and the PCT levels and also serum amyloid and others.² It is clear that a single biomarker which is ideal though has not been identified but a series specific studies on individual pathogen mediated sepsis has been studied extensively. Challenges that are faced during biomarker research lie in the lack of uniform protocol and methodology, standardization of the assays and the detection levels to be fixed age wise from population to population. The reliable detection of Biomarkers can be useful to have an immense diagnostic and prognostic relevance in relation to the management of sepsis.²

Sepsis is considered as the most important cause of morbidity and mortality. By sepsis we mean infection of the blood and if it is caused by bacteria, we term it as bacteremia and if by virus it is termed viremia. As such the infection in the blood is not so dangerous but as blood perfuses the important tissues and organs infections can be carried by blood to the kidneys or lungs which can result in the end organ failures of a patient leading to death in severe conditions. The main issue is that sepsis often lacks clinical manifestations so if the specific and the sensitive indicators of the infections are collected and analyzed then it will create superior outcomes in the management of the disease. Immunologically it can be considered as the dysregulation of the response of the innate immune system. Biomarkers are hence important to signal the presence and severity of sepsis in early conditions for effective prognostic values and importance.^{1,2}

It is very important to note that sepsis is not a true disease and just a physiological manifestation of the systemic immune response in innate immunity majority of the studies define sepsis to be a systemic inflammatory response to a bacteria, virus or fungi. In clinical setting several other physiological symptoms must be presented to properly diagnose sepsis. The common treatments of patients suspected with sepsis aims to eradicate the bacteria in the blood or reduce its growth by using broad spectrum antimicrobials. The gold standard of sepsis diagnosis has been traditionally the use of microbial cultures to identify the source of illness. The major limitation of using cultures in the length of the time required to develop cultures to identifiable quantities to detect sepsis and the possible microbes to start the treatment with antimicrobials and this with Bactek culture takes around 48 hours and hence there is a need of early detection through the detection of serum biomarkers which thus becomes important. Even the cultures might be insensitive under several conditions including the slow growing and non cultivatable organisms present at very low concentration. So, in the light of this problem alternative diagnostic methods using molecular biology tool kits have been developed to address the situation for rapid and automated detection of sepsis. These tests are basically the ELISA tests, flow cytometric detection of the specific marker antigens, immunocolorimetric tests, FISH methodologies and PCR and amplified PCR techniques and the main ideology was to detect the marker proteins.³

In parallel to the development of biomarkers which provides a faster and more sensitive diagnostic methods for infectious microbes thus relies on the monitoring of the changes in specific serum protein biomarker concentrations.

Biomarker is best defined as a factor or feature that can be measured and evaluated objectively as an indicator of normal biological responses or pathological processes. This article will focus on the wide range of biomarkers that is being in use and are destined to be launched in the market on the basis of their diagnostic accuracy (DA) which takes into account parameters like sensitivity, specificity and the predictive values.

For the purpose of being clinically relevant the biomarkers should have a very high DA that is it must have a very high sensitivity, specificity and positive predictive values and at the same time having negative predictive values. Some studies also calculated the likelihood ratios to determine the results which were actual cause of the disease or just occurred due to random chance. These values helped to assign the cut off limits for the purpose of correct diagnosis. As sepsis is a condition caused by bacterial, fungal or viral infections and thus leading to the infection of blood, the biomarkers will also be discussed for the specific pathogens in relation to sepsis. The individual markers discussed below may not be an absolute and specific for a particular diseased microbe but can be used to specify the class of the microbe responsible for sepsis so as to design target methods for antibiotic initiation or other antimicrobials for the management of this life-threatening critical condition.⁴

Biomarkers for the Infections caused by Bacteria

C-reactive protein

This is the general acute phase reactant protein rises in concentration up to 1000 times in the blood in response to the inflammation and infection. The sensitivity values have been reported in the range of 30-97% with specificity values of 75-100% and PPV of 31-100%. The use of different assays caused the variation of the results, as because each assay had unique functional limits assigned. Serum CRP levels also varied with age and in healthy individuals. A CV of 10mg/l was used for elderly patients over 65 years of age, while a CV of 5mg/ml was used for adults and infants in the range of 3-7 days old. CRP levels are higher in older children than in younger ones.^{3,5} Studies have indicated that CRP is highly sensitive and is highly specific for infection. This is the reason it can differentiate between gram positive and Gram negative infections. It can also be used as a prognostic marker for community acquired pneumonia and suggesting a need for hospitalization with a CV value of 110mg/l. It also differentiates between bacterial and fungal infections where the levels above 100mg/l are indicated for bacterial infections and below that as fungal infections. CRP is however unreliable for viral infections.⁶

Procalcitonin

PCT is a precursor of the hormone calcitonin and is another potent biomarker for bacterial infections. It has been reported to be in low concentrations of 0.33ng/ml in the serum of healthy individuals and increases around 1000 times under the inflammatory conditions. It is reported to rise within 2 to 4 hours of infection and peak at 6-8hr. Persistent increased levels will indicate continuous presence of bacterial infections. The rapid upregulation and sustenance in the serum makes it a unique biomarker. And as this molecule is highly stable it remains stable in blood preparation methods and freezing procedures even after long term storage.⁷

PCT has high DA and the sensitivity values around 75-100% and specificity values of 70-100%. The PPV values are 55-100%. Many studies have CV of PCT in the range of 0.5ng/ml a range of CVs has been proposed. As a biomarker for the bacterial infections most studies find PCT to be accurate biomarker than other inflammatory markers. It has very high prognostic values. PCT is also distinguish fungal and viral infections. During the viral infections the PCT values remain low often like the values found in the normal individuals. PCT is also used to make difference between bacterial and viral meningitis. In contrast fungal infections tend to cause mild elevations in the PCT concentration compared with the levels seen in the bacterial infections. Studies on invasive Aspergillosis and invasive candidiasis has shown high levels of PCT values in patients with bacterial sepsis than with fungal sepsis.^{5,8} PCT has also shown to be

a potential biomarker for pneumonia, abdominal infections, urinary tract infections, lower respiratory infections, myocardial infarctions and as a biomarker for the speculated therapy with antibiotics in patients with community acquired pneumonia.

Serum Amyloid A

Serum Amyloid A is an important apolipoprotein which is reported to have potential value for diagnosis of sepsis. SAA is expressed in levels around 1000times higher after 8-24 hours from the onset of sepsis. Compared to the levels of CRP, SAA levels are reported to rise faster and higher after the onset of sepsis and remain at higher relative elevations. Serum SAA levels of less than 15mg/l for the elderly over 65 years and less than 10mg/l for adults and newborns aged 3-7days has been indicated for fixing diagnostic criteria.

Though SAA is mainly used as markers for bacterial infections but it can be used to report for viral infections above the levels of 10mg/l among the infected patients. SAA might not be too sensitive and hence its clinical values may be overestimated.⁸

Biomarkers for the Infections caused by Fungi

Mannan and Antimannan antibodies are used to detect the fungal infections due to the presence of mannan in the cell walls of the invasive fungal pathogens. Both the levels of Mannan and Antimannan remain elevated high in conditions like aspergillosis and candidiasis making them useful markers in fungal sepsis. There are lot of variations in the DA, PPV, sensitivity and specificity values due to the variation of the assay protocols.⁹

The main disadvantage Mannan and Antimannan antibodies test alone is the high rate of false positives and negatives that the assays produce requiring these tests to be used as markers in conjunction with other diagnostic tests. Beta -D glucan tests have been used in combination with these tests because Beta-D glucan is highly sensitive and specific for invasive mycosis and are not species specific.¹⁰

Biomarkers for Viral Infections

Interferon Gamma inducible protein 10 (IP-10)

IP-10 a proinflammatory chemokine is an important biomarker for diagnosing viral infections due to its role in host response to viral infections. Study on bacterial infections at very low birth weight infants concluded that IP-10 is a good biomarker during initial measurements recording elevated levels in infected patients at both initial and 24-hour measurements with no overlap of the ranges. IP-10 levels can correlate also with the severity of sepsis. IP-10 is released in response to viruses like rhinovirus,

respiratory syncytial virus, hepatitis B and hepatitis C and H5N1 influenza virus. It has also shown to be a very good marker for prognostic treatment in hepatitis C patients.¹¹

Other Biomarkers on Avenue

IL-10

IL-10 has been indicated as a useful biomarker for sepsis specially in the diagnostic values in early and late onset of sepsis amongst the neonates where CV values of greater than 17.3 pg/ml has been estimated as significant DA values. This marker is also sensitive and specific to the tune of 80%. The main problem is that in general circulating cytokines have very short half life thus resulting in false negatives hence limiting their DA values.¹²

Lipopolysaccharide Binding Protein

Lipopolysaccharide Binding Protein is also a useful biomarker specific to bacterial infections. Serum Lipopolysaccharide Binding Protein levels are known to increase during bacterial and fungal infections but not in viral infections.¹³

Soluble triggering receptor

Triggering receptor expressed on myeloid cells-1 TREM are a group of cell-surface receptors that belong to the immunoglobulin superfamily [26]. TREM-1 is expressed mainly on macrophages and neutrophils, and has been identified as an amplifier of the immune response that strongly enhances leukocyte activation in the presence of microbial products [26, 27]. The latter results suggest that sTREM-1 can be released by a broad spectrum of inflammatory stimuli. The first promising result of the use of sTREM-1 in plasma to diagnose sepsis in ICU patients indicated that sTREM-1 might be that perfect diagnostic sepsis biomarker that everybody had been looking for. Results are ranging from an accuracy of almost 100 % (a ROC-AUC of 0.97) for the diagnosis of sepsis in ICU patients.¹⁴

TLRs

The results obtained by our group show that TLR and other cellular surface receptors may be differently regulated on mononuclear cells and neutrophils, and that they are dynamically modulated across the stages of sepsis. Toll-like receptor signaling gene expression in mononuclear cells is decreased in more severe forms of the disease. In contrast, up-regulated genes are seen along the clinical spectrum of sepsis in neutrophils.¹² Expressed by the macrophages, dendritic cells, and other cells plays

an important role in the innate immune responses to response to infection by cross reacting and recognizing specific bacterial antigens.¹⁵ TLR 4 is critical for recognition of LPS while the tLR-2 is useful to recognize gram positive bacterial components.^{5,12}

Toll like Receptors 2 (TLR-2) and Neutrophil CD64 receptors

Toll like Receptors 2 (TLR-2) and Neutrophil CD64 receptors are two important biomarkers present on the cell surfaces, during infection both these biomarkers are upregulated on the surface of the monocytes and neutrophils¹⁵. These receptors are significantly upregulated during both bacterial and viral infections. They are increased in conditions like influenza A and B, respiratory syncytial virus, mumps, Herpes zoster virus, cytomegalovirus infections which indicates that TLR-2 could be potential biomarkers for viral infections^{5,15}. Up to now, the CD64 is used for the marking of bacterial infections. CD64 expression is not affected by the underlying inflammatory diseases and is not differentially between systemic and local bacterial infections, Da values, sensitivity values and specificity values with PPV range from 71-100% justifying it as a significant biomarker.

CD163

CD163 is transmembrane molecule found on the membrane of mononuclear phagocytes. Being a specific scavenging receptor for hemoglobin heme inside the body it is very capable to identify hemoglobin-haptoglobin complex. CD163 regulates the expression of anti-inflammatory molecules like IL-10 and hemeoxygenase-1. Soluble CD163 comes from the molecules peeled from these phagocyte surfaces they becomes very good markers of sepsis and severe sepsis and was found to be superior than the PCT and CRP levels.

Micro RNAs

They are the type of endogenous non coding small RNAs about 22 nucleotides in length and play important roles by inhibiting the expression of the messenger RNAs. A significant amount of the miRNAs has been observed outside the cells within the various body fluids. These cell free miRNAs are stable under harsh conditions. miR-150 has been indicated as a biomarker for sepsis and the plasma ratio of miR-150/interleukin 18 can be used to study sepsis severity.

SNP

C-reactive protein (CRP) is an important biomarker of sepsis. Several single-nucleotide polymorphisms (SNPs) in the CRP gene can determine plasma CRP levels and

also are risk factors in many diseases, such as cancer, arteritis, and diabetes. However, it is unknown whether polymorphisms in CRP are associated with susceptibility to and outcome of infantile sepsis.

TNF alpha

Tumor necrosis factor alpha a pleiotropic cytokine produced by the activated macrophages plays a key role in inflammatory response and in the progression of the autoimmune diseases. An association with sepsis and expression of TNF-2 genotype was found.¹⁶

IL-1

The interleukin 1 receptor associated kinase is important for the TLR2 and TLR4 induced activation of nuclear factor kappa beta which is a critical factor in the transcriptional regulation of many sepsis associated proinflammatory mediators.¹⁶

Sepsis, as an increasing cause of mortality in patients with infectious disease specially in the critically ill patients in the ICU requires rapid diagnosis and treatment, and hence the importance of biomarkers is highly important.

References

1. Kramer MS, Demissie K, Yang H, Platt RW, Sauve R, Liston R. The contribution of mild and moderate preterm birth to infant mortality. Fetal and Infant Health Study Group of the Canadian Perinatal Surveillance System. *JAMA*. 2000; 284: 843–9.
2. Vincent JL, Sakr Y, Sprung CL, et al. Sepsis occurrence in acutely ill patients investigators: sepsis in European intensive care units: results of the SOAP Study. *Crit Care Med*. 2006; 34: 344–53.
3. Kauss IAM, Grion CMC, Cardoso LTQ *et al*. The epidemiology of sepsis in a Brazilian teaching hospital. *Brazilian J. Infect. Dis*. 2010; 14(3), 264–270.
4. Wang P, Yang Z, He Y, Shu C. Pitfalls in the rapid diagnosis of positive blood culture. *Rev. Med. Microbiol*. 2010; 21(3), 39–43.
5. Papachristou GI, Clermont G, Sharma A et al. Risk and markers of severe acute pancreatitis. *Gastroenterol Clin N Am* 2007; 36: 277–96.
6. Rego MAC, Martinez FE, Elias J, Mussi-Pinhata MM. Diagnostic value of interleukin-6 and C-reactive protein on early onset bacterial infection in preterm neonates with respiratory distress. *J. Perinat. Med*. 2010; 38(5), 527–533.
7. Morris AC, Simpson AJ, Walsh TS. Hyperinflammation and mediators of immunosuppression in critical illness. In: Vincent JL, editor. *Annual update in intensive care and emergency medicine*. Berlin Heidelberg: Springer- Verlag; 2013; 135–44.

8. Cetinkaya M, Ozkan H, Koksall N, Celebi S, Hacimustafaoglu M. Comparison of serum amyloid A concentrations with those of C-reactive protein and procalcitonin in diagnosis and follow-up of neonatal sepsis in premature infants. *J. Perinatol.* 2009; 29(3), 225–231.
9. Méan M, Marchetti O, Calandra T. Bench-to-bedside review: Candida infections in the intensive care unit. *Crit Care.* 2008; 12: 204. doi: 10.1186/cc6212.
10. Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, Moreno R, Lipman J, Gomersall C, Sakr Y, et al. EPIC II Group of Investigators International study of the prevalence and outcomes of infection in intensive care units. *JAMA.* 2009; 302: 2323–9. doi: 10.1001/jama.2009
11. Grebely J, Feld JJ, Applegate T, Matthews GV, Hellard M, Sherker A, Petoumenos K, Zang G, Shaw I, Yeung B, George J, Teutsch S, Kaldor JM, Cherepanov V, Bruneau J, Shoukry NH, Lloyd AR, Dore GJ. Plasma interferon-gamma-inducible protein-10 (IP-10) levels during acute hepatitis C virus infection. *Hepatology.* 2013 Jun; 57(6): 2124–34. doi: 10.1002/hep.26263.
12. Mehr S, Doyle LW. Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J.* 2000; 19: 879–87.
13. Villar J, Perez-Mendez L, Espinosa E et al. Serum lipopolysaccharide binding protein levels predict severity of lung injury and mortality in patients with severe sepsis. *PLoS One* 2009; 4: e6818.
14. Barati M, Bashar FR, Shahrami R, et al. Soluble triggering receptor expressed on myeloid cells 1 and diagnosis of sepsis. *J Crit Care.* 2010; 25: 362. e1–6.
15. Tsujimoto H, Ono S, Efron PA, Scumpia PO, Moldawer LL, Mochizuki H. Role of Toll-like receptors in the development of sepsis. *Shock.* 2008 Mar; 29(3): 315–21. doi: 10.1097/SHK.0b013e318157ee55.
16. De JW, te VH, Prakken BJ, Kuis W, Rijkers GT. Simultaneous detection of 15 human cytokines in a single sample of stimulated peripheral blood mononuclear cells. *Clin Diagn Lab Immunol.* 2003; 10: 133–9.