
Marijana C. Jandrić-Kočić¹, Mario Mizdrarić^{2, 3}

HROMOGRANIN A

Sažetak: Hromogranin A (engl. Chromogranin A, CgA) je najvažniji član porodice granina. Skladišti se s hormonima i neurotransmiterima u velike vezikule gustog jezgra ili hromafine granule i zajedno s njima oslobađa konstitutivno ili u kontrolisanom putu.

U okviru ovog preglednog rada prikazana je uloga CgA i iz njega izvedenih biološki aktivnih peptida, kao i njegova upotreba kao markera određenih bolesti.

CgA ima značajnu ulogu u sekretornoj granulogenezi (sortiranju i obradi sekretornog proteinskog sadržaja, sazrijevanju i kondenzaciji sekretornih granula) i održavanju homeostaze kalcijuma. Glikozilacijom, fosforilacijom, sulfatacijom i/ili proteolitičkom obradom CgA nastaju biološki aktivni peptidi kojim ispoljavaju hormonsku, autokrinu i parakrinu aktivnost. U kardiovaskularnom, metaboličkom, imunološkom i difuznom neuroendokrinom sistemu CgA i iz njega izvedeni biološki aktivni peptidi imaju mnogobrojne, ponekad oprečne uloge. CgA je identifikovan kao važan imunohistohemijski tumorski marker neuroendokrinih tumora, karcinoma prostate i mikrocelularnog karcinoma pluća s neuroendokrinom diferencijacijom. Njegova vjerodostojnost je uslovljena samim tumorom (lokalizacijom, veličinom, gustoćom sekretornih granula i sekretornom aktivnošću), diseminacijom tumora i primjenjem dijagnostičkom tehnikom. Povišene vrijednosti CgA se ne mogu koristiti u procjeni progresije tumora koji nisu neuroendokrinog podrijetla. U hroničnoj insuficijenciji srca i akutnom koronarnom sindromu CgA predstavlja nezavisni prognostički faktor bolesti.

Ključne riječi: hromogranin A, difuzni neuroendokrini sistem, tumor, bolest

Summary: Chromogranin A (CgA) is the most important member of the granin family. It is stored with hormones and neurotransmitters in large dense core vesicles or chromaffin granules and together with them it is released constitutively or in a controlled way.

¹ Marijana C. Jandrić-Kočić, Dom zdravlja Krupa na Uni, marijanajandrickocic@gmail.com

² Univerzitetski klinički centar Republike Srpske

³ Medicinski fakultet Univerziteta u Banjaluci

In this review are presented the role of CgA and biologically active peptides derived from it, as well as its use as a marker of certain diseases.

CgA plays a significant role in sectorial granulogenesis (sorting and processing of secretory protein content, maturation and condensation of secretory granules) and maintaining calcium homeostasis. Glycosylation, phosphorylation, sulfation, and/or proteolytic processing of CgA results in biologically active peptides that exhibit hormonal, autocrine, and paracrine activity. In the cardiovascular, metabolic, immune and diffuse neuroendocrine system, CgA and biologically active peptides derived from it have numerous, sometimes conflicting roles. CgA has been identified as an important immunohistochemical tumor marker of neuroendocrine tumors, prostate cancer and microcellular lung cancer with neuroendocrine differentiation. Its credibility is conditioned by the tumor itself (localization, size, density of secretory granules and secretory activity), dissemination of the tumor and application of the diagnostic technique. Elevated CgA values cannot be used to assess the progression of tumors that are not of neuroendocrine origin. In chronic heart failure and acute coronary syndrome, CgA is an independent prognostic factor of the disease.

Key words: chromogranin A, diffuse neuroendocrine system, tumor, disease

UVOD

Porodicu granina čine hromogranin A, hromogranin B, sekretogranin II (ili hromogranin C), sekretogranin III (ili produkt gena 1B1075), sekretogranin IV (ili antigen HSL-19), sekretogranin V (ili antigen 7B2), sekretogranin VI (ili neuroendokrini sekretorni protein 55) i sekretogranin VII (ili inhibitor proprotein konvertaze subtilizin/keksin tipa 1) (1, 2). Granini su proteini molekulske mase od 27 do 10 kilodaltona (engl. kilodalton, kDa) (1). Izgrađeni su od jednostrukih polipeptidnih lanaca s najmanje 10% kiselinskih ostataka (glutaminske ili asparaginske kiseline) i višestrukih pojedinačnih i dvobaznih aminokiselinskih ostataka (3). Uskladišteni su u sekretornim granulama endokrinih, neuroendokrinih i neuronskih ćelija (1). Granini imaju značajnu ulogu u sekretornoj granulogenezi, sortiranju i obradi sekretornog proteinskog sadržaja, sazrijevanju i kondenzaciji sekretornih granula (3). Oni održavaju homeostazu kalcijuma i ispoljavaju hormonsku, autokrinu i parakrinu aktivnost (1, 4, 5).

Granini se proizvode u granuliranom endoplazmatskom retikulumu i preko signalnog peptida na N terminalnom kraju molekula ulaze u unutrašnji prostor retikuluma (3, 4). Otuda, putem transportnih vezikula, oni ulaze u Goldžijev kompleks (3). Potom se, s preostalim sadržajem vezikula, u trans mreži Goldžijevog kompleksa pakuju u

nezrele sekretorne granule (1, 3, 4). Tu se djelimično prerađuju u različite biološki aktivne peptide (3). Nezrele granule sazrijevaju naknadnom acidifikacijom sadržaja granula i uklanjanjem klatrinske ovojnice, sinaptotagmina IV i membranskog proteina 4 (1, 3). Zrele sekretorne granule s gustim jezgrom (engl. Dense-core secretory granules, DCG), poznate kao velike vezikule gustog jezgra (engl. Large dense-core vesicles, LDCV) ili hromafine granule (engl. Chromaffin granules, CG) skladište se u srži nadbubrežne žlijezde i njihov sadržaj se oslobađa nakon stimulacije (3).

Granini olakšavaju stvaranje DCG kada se nezrela granula oslobađa iz trans mreže Goldžijevog kompleksa (4, 5). Oni omogućuju regrutovanje i kondenzaciju topivog proteinskog sadržaja sekretornih granula u guste jezgre (5). Prisustvo granina u sekretornim granulama, kao i njihov visok kapacitet vezivanja kalcijuma pri niskom pH, osigurava skladištenje kalcijuma u ćelije (4). Proteolitičkom razgradnjom granina nastaju proteini koji imaju hormonsku, autokrinu i parakrinu aktivnost (2, 4, 5).

Granini i iz njih izvedeni bioaktivni fragmenti doprinose oslobađanju određenih hormona, kao i homeostazi glukoze, neuronskoj ekscitabilnosti, autoimunosti i kontraktilnosti glatkih mišića (4). Na taj način oni moduliraju metabolizam, puteve bola, raspoloženje i krvni pritisak (4). Granini predstavljaju markera specifičnih bolesti, uključujući neurodegenerativne bolesti, psihijatrijske bolesti i karcinome (4).

METODE

Literatura je pretražena korišćenjem ključnih reči: hromogranin A, difuzni neuroendokrini sistem, tumor, bolest. Pretraživanje je sprovedeno u okviru sledećih baza podataka: PubMed, Emabase, Scopus, SCIn-dex i Hrčak. Zbog ograničenog broja dostupnih studija u pretraživanju baza nisu korišteni dostupni filteri. Nakon iščitavanja sažetaka, radovi su detaljnije proučeni, te su isključeni oni koji ne odgovaraju postavljenom cilju istraživanja.

HROMOGRANIN A

Hromogranin A (engl. Chromogranin A, CgA) je prvi identifikovani predstavnik granina (6). Njegovo ime potiče od mjesta primarne identifikacije, hromafinih vezikula srži nadbubrežne žlijezde koje sadrže katecholamine (engl. Adrenal medulla chromaffin vesicles containing catecholamines) (6).

CgA je protein molekularne težine 48 kDa, izgrađen od 439 aminokiselina (1). Hromafina ćelija srži nadbubrežne žlijezde predstavlja glavni izvor cirkulišućeg CgA (2). Adrenergički nervni završeci i neuroendokrine ćelije izlučuju CgA u perifernim tkivima (2). CgA je otkriven u adenohipofizi, plućima, pankreasu, želucu, tankom crijevu (jejunumu i ileumu), frontalnom cerebralnom korteksu,

štitnoj žlijezdi, paraštitnim žlijezdama, miokardu i polimorfonuklearnim leukocitima (2, 7–10).

Proizvodnju CgA reguliše nekoliko unutarćelijskih glasničkih sistema, koji uključuju protein kinazu A, protein kinazu C, kao i unutarćelijski kalcijum (8). Histamin, nikotin, bradikinin, angiotenzin II, prostaglandin E2 i kalijum mogu olakšati proizvodnju CgA (8). Zajedno s hormonima ili neurotransmiterima, CgA se oslobađa konstitutivno ili u kontrolisanom putu (8).

Gen CgA nalazi se na hromosomu 14q32.12 i obuhvata 12.192 baznih parova (engl. base pair, bp), organizovanih u osam egzona i sedam introna (1). Primarni transkript od 2 kilobaza (engl. kilobase, kb) provodi se u protein molekularne težine 52 kDa izgrađen od 457 aminokiselina koji prolazi kroz posttranslacijske modifikacije, odnosno endoproteolizu prohormonskim konvertazama (1). Proteolitičkim cijepanjem CgA nastaje nekoliko biološki aktivnih peptida: vazostatini 1 i 2, prohromacin, hromacini 1 i 2, hromofungin, pankreastatin, katestatin, parastatin, serpinin, WE14, EA-92, SS-18, LF-19, GR-44, SS-18, AL-11, ER-37, ES-43, GV 19, WA-8 (1, 7).

FIZIOLOŠKE ULOGE HROMOGRANINA A I PRODUKATA NJEGOVOG CIJEPANJA

CgA ima mnogobrojne fiziološke uloge u ćeliji i izvan nje (Tabela 1) (1, 7, 9). Inhibicijom razgradnje granula posredovane serpininom on modulira biogenezu DCG (1). Takođe, ima važnu ulogu u obnavljanju ćelija sekretornih granula nakon egzocitoze (1). Obilje (~2-4 mM) CgA unutar DCG omogućava pohranu velike količine kalcijuma u njima (1). U isto vrijeme, CgA omogućava brzu razmjenu slobodnog i vezanog kalcijuma unutar DCG, kao i mobilizaciju kalcijuma u citoplazmu (kroz aktivaciju inozitol trifosfatnih receptora kalcijumovih kanala na membranama DCG) (1).

Uloga CgA izvan ćelije nije u potpunosti jasna (1, 7). CgA inhibira izlučivanje proopiomelanokortina u prednjem režnju hipofize (15). Indukovanjem proizvodnje faktora nekroze tumora alfa i difuzibilnih neurotoksičnih supstanci u ćelijama mikroglije uzrokuje apoptozu neurona (1). Značajan broj uloga, prvobitno smatranim ulogama CgA, pripisane su biološki aktivnim peptidima koji nastaju glikozilacijom, fosforilacijom, sulfatacijom i/ili proteolitičkom obradom CgA (7). Većina njih se sastoji od direktne ili indirektne inhibitorne regulacije putem autokrinih, parakrinih ili hormonskih puteva (7). Do sada je otkrivena uloga katestatina, prohromacina, hromacina 1 i 2, hromofungina, pankreastatina, parastatina, vazostatina 1 i 2 i WE 14 (1, 7). Uloga preostalih biološki aktivnih peptida izvedenih iz CgA ostaje nepoznata (1, 7).

Tabela 1. Uloga hromogranina A i iz njega izvedenih biološki aktivnih peptida (7)

U ćeliji	Izvan ćelije
Granulogena aktivnost	Inhibiranje izlučivanja želučane kiseline
Skladištenje proteina	Inhibiranje peristaltike debelog crijeva
Skladištenje i regulisanje protoka kalcijuma	Inhibiranje vazokonstrikcije arterija
Prohormon	Negativan inotropni efekat
	Regulisanje metabolizma ugljenih hidrata, masti i proteina
	Inhibiranje izlučivanja: insulina, paratiroidnog hormona, kateholamina, pankreasne amilaze, proopiomelanokortina
	Antimikrobni učinak
	Antiupalna aktivnost
	Obnavljanje tkiva: regulisanje adhezije ćelija
	Regulisanje nocicepcije
	Aktivacija mikroglije u centralnom nervnom sistemu
	Apoptotička degeneracija kortikalnih neurona

Vazostatini 1 i 2 predstavljaju N-terminalne fragmente CgA (1). Oni inhibiraju vazokonstrikciju arterija posredovanu endotelinom-1 i imaju negativan inotropan i luzitropan učinak na srce, predominantno u prisustvu intenzivnih adrenergičkih stimulusa (1, 7). Nadalje, inhibiraju proliferaciju i migraciju endotelnih ćelija indukovanu vaskularnim endotelnim faktorom rasta (1). Zahvaljujući kardiotropnim i vazoaktivnim osobinama, u uslovima prekomjerne simpatičke stimulacije, oni mogu imati ulogu homeostatskog stabilizatora kardiovaskularnog sistema (1). Vazostatin 1 i 2 inhibiraju lučenje želučane kiseline i paratiroidnog hormona i imaju antimikrobni i antiupalni učinak (7). Naime, oni inhibiraju rast kvasaca, bakterija i gljivica i stimulišu migraciju makrofaga (1). Štite integritet endotelne barijere od proupalnih agenasa i moduliraju adheziju fibroblasta (7). U konačnici, vazostatini 1 i 2 indukuju neuronsku apoptozu i vrše kontrolu nocicepcije (7).

Pankreastatin je prvi identifikovani biološki aktivan peptid izveden iz CgA (1). On inhibira egzokrinu aktivnost pankreasa i želuca, kao i oslobađanje paratiroidnog hormona (1, 7). Modifikuje omjer inzulina:glukagon stimulišući izlučivanje glukagona i inhibirajući izlučivanje insulina uslovljeno fiziološkim stimulusima (7). Povećava proizvodnju azotnog oksida koji inhibira insulin (9). Oslobođen s kateholaminima iz simpatičkog nervnog sistema u stresnim situacijama učestvuje u modulaciji energetskog metabolizma (1). Naime, pri neuravnoteženoj simpatičkoj aktivaciji, višak kateholamina zajedno s povišenim vrijednostima pankreastatina smanjuje unos glukoze

(za ~50%) i povećava oslobađanje slobodnih masnih kiselina (za 4,5 do 6,4 puta) (1). Pojedinačna istraživanja ukazuju da pankreastatin (mehanizmom koji nije u potpunosti jasan) ima značajnu ulogu u patogenezi različitih oblika dijabetes melitusa (9).

Katestatin djeluje na nikotinske acetilholinske receptore kao snažan autokrini inhibitor lučenja kateholamina (1, 7). Smatra se da nedostatak katestatina igra značajnu ulogu u razvoju hipertenzije uslovljene dugotrajnom prekomjernom aktivnošću simpatičkog nervnog sistema (1). Indukovanjem oslobađanja histamina iz mastocita on ostvaruje vazorelaksanti i antihipertenzivni učinak (1). Katestatin posjeduje izraženu angiogenu i vaskulogenu aktivnost, budući da indukuje migraciju i proliferaciju endotelnih i glatkih mišićnih ćelija krvnih sudova (1). On stimulira hemotaksu aktivnost i aktivnost proupalnih citokina, te inhibira rast gljivica, kvasaca i bakterija (Gram-pozitivnih i Gram-negativnih) (1).

Parastatin je otkriven u paratiroidnoj žlijezdi svinje (1). U prisustvu niskih vrijednosti kalcijuma u krvi on inhibira lučenja paratiroidnog hormona (1).

Prohromacin i iz njega izvedeni biološki aktivni peptidi, hromacini 1 i 2, inhibiraju rast gljivica i bakterija (7, 10).

WE 14 može modulirati vrijednost gonadotropnih hormona (7). Osim toga, istraživanja eminentnih autora identifikovali su WE 14 kao autoantigen pankreasa koji ima značajnu ulogu u patogenezi različitih oblika dijabetes melitusa (9).

METODE ODREĐIVANJA VRIJEDNOSTI HROMOGRANINA A U KRV

Postoji niz komercijalno dostupnih setova za određivanje CgA u krvi (6). Oni su zasnovani na tri dijagnostičke tehnike: enzimski imunosorbentni test (engl. Enzyme linked immunosorbent assay, ELISA), imunoradiometrijski test (engl. Immunoradiometric assay, IRMA) i radioimunološki test (engl. Radioimmunoassay, RIA) (6). Iako značajan broj istraživanja preporučuje RIA, ne postoji univerzalno prihvaćena dijagnostička tehnika, kao ni standardizacija niti jedne od navedenih dijagnostičkih tehnika (6, 11). Samim tim, preporučuje se oprez pri pokušaju poređenja rezultata različitih dijagnostičkih centara (6). CgA se može određivati u plazmi ili serumu (6). Vrijednost CgA u plazmi je nerijetko značajno viša od vrijednosti utvrđene u serumu (6). Istraživanja određenih autora preporučuju određivanje CgA u pljuvački (1).

HROMOGRANINA A KAO DIJAGNOSTIČKI I PROGNOСТИČKI MARKER

Porast vrijednosti CgA u krvi nastaje u određenim benignim bolestima i stanjima, malignim bolestima, kao i jatrogeno (Tabela 2) (12).

CgA je povišen u osoba koje koriste inhibitore protonске pumpe, antagoniste histaminskih H2 receptora i selektivne inhibitore ponovnog preuzimanja serotonina (12). Samim tim, preporučuje se prekid upotrebe lijekova (antagonisti histaminskih H2 receptora 24 sata, inhibitora protonске pumpe 14 dana) prije određivanja vrijednosti CgA u krvi (12). Intenzivna fizička aktivnost, kao i unos hrane neposredno prije testiranja rezultuju porastom vrijednosti CgA u krvi (12). Lagano povišene vrijednosti CgA utvrđene su u menopauzi (usljed povišenog simpatičkog tonusa) i trudnoći (1).

Tabela 2. Uzroci porasta vrijednosti CgA u krvi (1, 12)

Benigne bolesti i stanja	Jatrogeni uzroci	Maligne bolesti
Trudnoća	Inhibitori protonске pumpe	Karcinom želuca
Menopauza	Antagonisti histaminskih H2 receptora	Karcinom pankreasa
Hronični atrofični gastritis	Selektivni inhibitori ponovnog preuzimanja serotonina	Hepatocelularni karcinom
Helicobacter pylori infekcija	Intenzivna fizička aktivnost	Karcinom dojke
Pankreatitis	Skorašnji unos hrane	Karcinom jajnika
Hronični hepatitis		Karcinom prostate
Ciroza jetre		Neuroblastom
Sindrom iritabilnog crijeva		Mikrocelularni karcinom pluća
Ulcerozni kolitis		Neuroendokrini tumori
Chronova bolest		
Arterijska hipertenzija		
Srčana insuficijencija		
Akutni koronarni sindrom		
Infarkt miokarda		
Giantcell arteritis		
Bubrežna insuficijencija		
Hronični bronhitis		
Opstruktivne bolesti pluća		
Reumatoidni artritis		
Sistemiški eritematozni lupus		
Hipertiroidizam		
Hiperparatiroidizam		
Hiperkortizolemija		

Hronični atrofični gastritis i gastritis uzrokovan *Helicobacter pylori* mogu uzrokovati porast vrijednosti CgA, usljed hroničnog porasta vrijednosti gastrina u serumu (1). U osoba s ovim bolestima, predominantno u onih liječenih inhibitorima protonske pumpe, mjerenje vrijednosti CgA u krvi može doprinijeti praćenju hiperplazije ćelija želuca sličnih enterohromafinu (1). Ulcerozni kolitis i Crohnova bolest uzrokuju diskretan porast vrijednosti CgA u krvi (1).

Kod disfunkcije organa, poput insuficijencija bubrega, jetre i srca, vrijednosti CgA u krvi također mogu biti značajno povećane (1). Veoma visoke vrijednosti CgA u osoba s insuficijencijom bubrega (usporedive s vrijednostima CgA u osoba s neuroendokrinim tumorom) govore u prilogu značajne uloge bubrega u uklanjanju CgA iz tijela (2). Njegove umjereno povišene vrijednosti u osoba s insuficijencijom jetre ukazuju ili na metabolizam CgA u jetri ili na aktivaciju neuroendokrinog sistema u ovih osoba (2). Aktivacija neuroendokrinog sistema prisutna je i u osoba s insuficijencijom srca, akutnim koronarnim sindromom i hipertenzijom (1, 2). Visoke vrijednosti CgA u krvi ukazuju na visok rizik od smrti nakon infarkta miokarda, akutnog koronarnog sindroma i insuficijencijom srca (1).

Postoje visoke vrijednosti CgA u određenim upalnim bolestima, poput reumoidnog artritisa i sistemskog eritematoznog lupusa (1). Endokrini poremećaji povezani s povišenim vrijednostima CgA u krvi su hiperkortizolemija, hipertireoidizam i hiperparatireoidizam (1).

CgA predstavlja važan marker neuroendokrinih tumora, uključujući karcinoid tumor, feokromocitom, medularni karcinom tiroidne žlijezde, adenom paratiroidne žlijezde, neuroblastom, gastroenteropankreatične neuroendokrine tumore, sindrom multiple endokrine neoplazije tip 1 i bronhopulmonalne neuroendokrine tumore (13). Najveća vjerodostojnost ovog tumorskog markera utvrđena je u diseminovanim i dobro diferentovanim neuroendokrinim tumorima koji pokazuju intenzivnu sekretornu aktivnost (13).

Senzitivnost CgA u osoba s neuroendokrinim tumorima iznosi između 10 i 100%, a njegova specifičnost između 68 i 100% (13). Kod malih tumora (osim gastrinoma), poput insulinoma, paraganglioma, karcinoid tumora bronha i tumora hipofize, vrijednosti CgA mogu biti u referentnom rasponu (13). Najveća osjetljivost CgA utvrđena je kod gastrinoma (100%), feohromocitoma (89%), karcinoid tumora (80%), nefunkcionalnih tumora endokrinog pankreasa (69%) i medularnog karcinoma tiroidne žlijezde (50%) (13). U osoba s metastatskom bolešću, CgA posjeduje osjetljivost od 60 do 100% i njegova vrijednost u krvi može porasti čak 100–1000 puta (13). Najviše vrijednosti utvrđene su u karcinoid tumora s metastazama u jetri (13). Osim u postavljanju dijagnoze neuroendokrinih tumora, CgA je važan u praćenju odgovora na liječenje (13).

U osoba oboljelih od mikrocelularnog karcinoma pluća s neuroendokrinom diferencijacijom, CgA može pomoći u praćenju odgovora na liječenje i otkivanju

recidiva bolesti (13). Iako vrijednost CgA u krvi ukazuje na vjerojatni stepen neuroendokrine diferencijacije u karcinomu, njegova niska osjetljivost čini ga nepouzdanim u otkrivanju bolesti (13).

U osoba s karcinomom prostate određivanje vrijednosti CgA u krvi može pomoći u postavljanju dijagnoze i određivanju prognoze bolesti (2). Naime, vrijednosti CgA u krvi oboljelih mogu biti povišene čak i ako je vrijednost prostata specifičnog antigena s normalna (13). Visoke vrijednosti CgA u krvi mogu predvidjeti izostanak odgovora karcinoma na hormonsku terapiju, odnosno lošu prognozu bolesti (13).

Imunoreaktivnost CgA ustanovljena je u ćelijama kolorektalnih karcinoma i karcinoma dojke, ali njegova dijagnostička vrijednost nije utvrđena (2).

ZAKLJUČAK

CgA i iz njega izvedeni biološki aktivni peptidi imaju mnogobrojne, ponekad oprečne uloge u kardiovaskularnom, metaboličkom, imunološkom i difuznom neuroendokrinom sistemu. CgA je identifikovan kao važan imunohistohemijski tumorski marker neuroendokrinih tumora, karcinom prostate i mikrocelularnog karcinoma pluća s neuroendokrinom diferencijacijom. Njegova vjerodostojnost je uslovljena samim tumorom (lokalizacijom, veličinom, gustoćom sekretornih granula i sekretornom aktivnošću), diseminacijom tumora i primjenjenom dijagnostičkom tehnikom. Povišene vrijednosti CgA se ne mogu koristiti u procjeni progresije tumora koji nisu neuroendokrinog podrijetla. U hroničnoj insuficijenciji srca i akutnom koronarnom sindromu CgA predstavlja nezavisni prognostički faktor bolesti.

LITERATURA

1. D'amico MA, Ghinassi B, Izzicupo P, Manzoli L, Di Baldassarre A. Biological function and clinical relevance of chromogranin A and derived peptides. *Endocr Connect*. 2014 Apr 29; 3(2): R45–54. <https://pmc.ncbi.nlm.nih.gov/articles/PMC5395093/>
2. Taupenot L, Kimberly HL, O'Connor DT. The Chromogranin–Secretogranin Family. *NEJM*. 2003; 348(12): 1134–1149. <https://www.nejm.org/doi/full/10.1056/nejmra021405>
3. Feldman SA, Eiden LE. The chromogranins: Their roles in secretion from neuroendocrine cells and as markers for neuroendocrine neoplasia. *Endocr Pathol*. 2003; 14: 3–23. <https://doi.org/10.1385/EP:14:1:3>
4. Bartolomucci A, Possenti R, Mahata SK, Fischer-Colbrie R, Loh YP, Salton SR. The extended granin family: structure, function, and biomedical implications. *Endocr Rev*. 2011 Dec; 32(6): 755–97. <https://pmc.ncbi.nlm.nih.gov/articles/PMC3591675/>
5. Estevez-Herrera J, Pardo MR., Dominguez N, Pereda D, Machado JD, Borges R. The role of chromogranins in the secretory pathway. *BioMolecular Concepts*. 2013; 4(6): 605–609. <https://www.degruyter.com/document/doi/10.1515/bmc-2013-0020/html?sr->

- sltid=AfmBOoqMHFufhcNduEZBODJaF4PL_ddiPBhRUJL34a8zEZqRqx-ZKya2
6. Gut P, Czarnywojtek A, Fischbach J, Bączyk M, Ziemnicka K, Wrotkowska E et al. Chromogranin A – unspecific neuroendocrine marker. Clinical utility and potential diagnostic pitfalls. Arch Med Sci. 2016 Feb 1; 12(1): 1–9. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4754364/>
7. Díaz Pérez JÁ, Currás Freixes M. Cromogranina A y tumores neuroendocrinos. Endocrinol Nutr. 2013; 60: 386–395. <https://www.elsevier.es/en-revista-endocrinologia-nutricion-412-articulo-chromogranin-a-neuroendocrine-tumors-S2173509313001372#bib0100>
8. Louthan O. Chromogranin A in Physiology and Oncology. Folia Biologica (Praha). 2011; 57: 173–181. <https://fb.cuni.cz/file/5597/FB2011A0026.pdf>
9. Herold Z, Doleschall M, Kovesdi A, Patocs A, Somogyi A. Chromogranin-A and its role in the pathogenesis of diabetes mellitus. Endokrynologia Polska. 2018; 69(5): 598–610. https://journals.viamedica.pl/endokrynologia_polska/article/view/EP.a2018.0052/45437
10. Scavello F, Kharouf N, Lavalley P, Haikel Y, Schneider F, Metz-Boutigue MH. The antimicrobial peptides secreted by the chromaffin cells of the adrenal medulla link the neuroendocrine and immune systems: From basic to clinical studies. Front Immunol. 2022 Aug 25; 13: 977175. <https://pmc.ncbi.nlm.nih.gov/articles/PMC9452953/#B23>
11. Stridsberg M, Eriksson B, Oberg K, Janson ET. A comparison between three commercial kits for chromogranin A measurements. J Endocrinol. 2003 May; 177(2): 337–41. <https://pubmed.ncbi.nlm.nih.gov/12740022/>
12. Di Giacinto P, Rota F, Rizza L, Campana D, Isidori A, Lania A et al. Chromogranin A: From Laboratory to Clinical Aspects of Patients with Neuroendocrine Tumors. Int J Endocrinol. 2018 Jul 2; 2018: 8126087. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6051263/>
13. Glinicki P, Jeske W. Chromogranin A (CgA) – the influence of various factors *in vivo* and *in vitro*, and existing disorders on it's concentration in blood. Endokrynologia Polska/Polish Journal of Endocrinology 2010; 61(4): 384–397. [file:///C:/Users/W10/Downloads/25370-31124-1-PB%20\(1\).pdf](file:///C:/Users/W10/Downloads/25370-31124-1-PB%20(1).pdf)

Marijana C. Jandric-Kocic¹, Mario Mizdraric^{2, 3}

CHROMOGRANIN A

Summary: Chromogranin A (CgA) is the most important member of the granin family. It is stored with hormones and neurotransmitters in large dense core vesicles or chromaffin granules and together with them it is released constitutively or in a controlled way.

In this review are presented the role of CgA and biologically active peptides derived from it, as well as its use as a marker of certain diseases.

CgA plays a significant role in sectorial granulogenesis (sorting and processing of secretory protein content, maturation and condensation of secretory granules) and maintaining calcium homeostasis. Glycosylation, phosphorylation, sulfation, and/or proteolytic processing of CgA results in biologically active peptides that exhibit hormonal, autocrine, and paracrine activity. In the cardiovascular, metabolic, immune and diffuse neuroendocrine system, CgA and biologically active peptides derived from it have numerous, sometimes conflicting roles. CgA has been identified as an important immunohistochemical tumor marker of neuroendocrine tumors, prostate cancer and microcellular lung cancer with neuroendocrine differentiation. Its credibility is conditioned by the tumor itself (localization, size, density of secretory granules and secretory activity), dissemination of the tumor and application of the diagnostic technique. Elevated CgA values cannot be used to assess the progression of tumors that are not of neuroendocrine origin. In chronic heart failure and acute coronary syndrome, CgA is an independent prognostic factor of the disease.

Key words: chromogranin A, diffuse neuroendocrine system, tumor, disease

INTRODUCTION

The granin family consists of chromogranin A, chromogranin B, secretogranin II (or chromogranin C), secretogranin III (or 1B1075 gene product), secretogranin IV (or

¹ Marijana C Jandric-Kocic, Health care Center Krupa na Uni, Republic of Srpska, Bosnia and Herzegovina, marijanajandrickocic@gmail.com

² University Clinical Center of the Republic of Srpska

³ University of Banja Luka, Faculty of Medicine

HISL-19 antigen), secretogranin V (or 7B2 antigen), secretogranin VI (or neuroendocrine secretory protein 55) and secretogranin VII (or subtilisin/kexin type 1 proprotein convertase inhibitor) (1,2). Granins are proteins with a molecular weight of 27 to 10 kilodaltons (kDa) (1). They are built from single polypeptide chains with at least 10% acid residues (glutamic or aspartic acid) and multiple single and dibasic amino acid residues (3). They are stored in secretory granules of endocrine, neuroendocrine and neuronal cells (1). Granins play a significant role in sectorial granulogenesis, sorting and processing of secretory protein content, maturation and condensation of secretory granules (3). They maintain calcium homeostasis and exhibit hormonal, autocrine and paracrine activity (1,4,5).

Granins are produced in the granular endoplasmic reticulum and enter the inner space of the reticulum via the signal peptide at the N terminal end of the molecule (3,4). From there, via transport vesicles, they enter the Golgi complex (3). Then, with the remaining contents of the vesicles, they are packed into immature secretory granules in the trans network of the Golgi complex (1,3,4). There, they are partially processed into various biologically active peptides (3). Immature granules mature by subsequent acidification of the granule contents and removal of the clathrin envelope, synaptotagmin IV and membrane protein 4 (1,3). Mature dense-core secretory granules (DCG), known as large dense-core vesicles (LDCV) or chromaffin granules (CG) are stored in the medulla of the adrenal gland and their contents are released after stimulation (3).

Granins facilitate DCG formation when an immature granule is released from the trans network of the Golgi complex (4,5). They enable the recruitment and condensation of the soluble protein content of the secretory granules into dense nuclei (5). The presence of granins in secretory granules, as well as their high calcium binding capacity at low pH, ensures calcium storage in cells (4). Proteolytic degradation of granins produces proteins that have hormonal, autocrine and paracrine activity (2,4,5).

Granins and bioactive fragments derived from them contribute to the release of certain hormones, as well as glucose homeostasis, neuronal excitability, autoimmunity and smooth muscle contractility (4). In this way, they modulate metabolism, pain pathways, mood and blood pressure (4). Granins represent a marker of specific diseases, including neurodegenerative diseases, psychiatric diseases and cancers (4).

METHODS

The literature was searched using keywords: chromogranin A, diffuse neuroendocrine system, tumor, disease. The search was conducted within the following databases: PubMed, Emabase, Scopus, SCIn-dex and Hrčak. Due to the limited number of available studies, no available filters were used in the database search.

After reading the abstracts, the papers were studied in more detail, and those that did not correspond to the research objective were excluded.

CHROMOGRANIN A

Chromogranin A (CgA) is the first identified representative of granins (6). Its name comes from the place of primary identification, adrenal medulla chromaffin vesicles containing catecholamines (6).

CgA is a protein with a molecular weight of 48 kDa, made of 439 amino acids (1). Chromaffin cells of the medulla of the adrenal gland represent the main source of circulating CgA (2). Adrenergic nerve endings and neuroendocrine cells secrete CgA in peripheral tissues (2). CgA has been detected in the adenohypophysis, lung, pancreas, stomach, small intestine (jejunum and ileum), frontal cerebral cortex, thyroid gland, parathyroid glands, myocardium and polymorphonuclear leukocytes (2,7-10).

The production of CgA is regulated by several intracellular messenger systems, which include protein kinase A, protein kinase C, as well as intracellular calcium (8). Histamine, nicotine, bradykinin, angiotensin II, prostaglandin E2 and potassium can facilitate the production of CgA (8). Together with hormones or neurotransmitters, CgA is released constitutively or in a controlled pathway (8).

The CgA gene is located on chromosome 14q32.12 and includes 12,192 base pairs (bp) organized in eight exons and seven introns (1). The primary transcript of 2 kilobases (kb) is made into a protein with a molecular weight of 52 kDa, made of 457 amino acids, which undergoes post-translational modifications, i.e. endoproteolysis by prohormone convertases (1). Proteolytic cleavage of CgA produces several biologically active peptides: vasostatins 1 and 2, prochromacin, chromacins 1 and 2, chromofungin, pancreastatin, catestatin, parastatin, serpinin, WE14, EA-92, SS-18, LF-19, GR-44, SS-18, AL-11, ER-37, ES-43, GV 19, WA-8 (1,7).

PHYSIOLOGICAL ROLE OF CHROMOGRANIN A AND ITS CLEAVAGE PRODUCTS

CgA has numerous physiological roles inside and outside the cell (Table 1) (1,7,9). By inhibiting granule degradation mediated by serpinin, it modulates DCG biogenesis (1). Also, it plays an important role in the recovery of secretory granule cells after exocytosis (1). The abundance (~2-4 mM) of CgA within DCG enables the storage of a large amount of calcium in them (1). At the same time, CgA enables the rapid exchange of free and bound calcium within the DCG, as well as the mobilization of calcium into the cytoplasm (through the activation of inositol triphosphate receptors of calcium channels on the DCG membranes) (1).

The role of CgA outside the cell is not completely clear (1,7). CgA inhibits the secretion of proopiomelanocortin in the anterior lobe of the pituitary gland (15). By inducing the production of tumor necrosis factor alpha and diffusible neurotoxic substances in microglia cells, it causes neuronal apoptosis (1). A significant number of roles, originally considered roles of CgA, have been attributed to biologically active peptides resulting from glycosylation, phosphorylation, sulfation and/or proteolytic processing of CgA (7). Most of them consist of direct or indirect inhibitory regulation via autocrine, paracrine or hormonal pathways (7). So far, the role of catestatin, prochromacin, chromacin 1 and 2, chromofungin, pancreastatin, parastatin, vasostatin 1 and 2 and WE 14 has been discovered (1,7). The role of the remaining biologically active peptides derived from CgA remains unknown (1,7).

Table 1. Role of chromogranin A and biologically active peptides derived from it (7)

In the cell	Outside the cell
Granulogenic activity	Inhibition of gastric acid secretion
Protein storage	Inhibition of colonic peristalsis
Storage and regulation of calcium flow	Inhibition of arterial vasoconstriction
Prohormone	Negative inotropic effect
	Regulation of carbohydrate, fat and protein metabolism
	Inhibition of secretion of: insulin, parathyroid hormone, catecholamine, pancreatic amylase, proopiomelanocortin
	Antimicrobial effect
	Anti-inflammatory activity
	Tissue regeneration: regulation of cell adhesion
	Regulation of nociception
	Activation of microglia in the central nervous system
	Apoptotic degeneration of cortical neurons

Vasostatins 1 and 2 represent N-terminal fragments of CgA (1). They inhibit vasoconstriction of arteries mediated by endothelin-1 and have a negative inotropic and lusitropic effect on the heart, predominantly in the presence of intense adrenergic stimuli (1,7). Furthermore, they inhibit proliferation and migration of endothelial cells

induced by vascular endothelial growth factor (1). Thanks to their cardiotropic and vasoactive properties, in conditions of excessive sympathetic stimulation, they can play the role of a homeostatic stabilizer of the cardiovascular system (1). Vasostatin 1 and 2 inhibit the secretion of gastric acid and parathyroid hormone and have an antimicrobial and anti-inflammatory effect (7). Namely, they inhibit the growth of yeasts, bacteria and fungi and stimulate the migration of macrophages (1). They protect the integrity of the endothelial barrier against pro-inflammatory agents and modulate the adhesion of fibroblasts (7). Ultimately, vasostatins 1 and 2 induce neuronal apoptosis and control nociception (7).

Pancreastatin is the first identified biologically active peptide derived from CgA (1). It inhibits the exocrine activity of the pancreas and stomach as well as the release of parathyroid hormone (1,7). It modifies the insulin:glucagon ratio by stimulating the secretion of glucagon and inhibiting the secretion of insulin conditioned by physiological stimuli (7). It increases the production of nitric oxide which inhibits insulin (9). Released with catecholamines from the sympathetic nervous system in stressful situations, it participates in the modulation of energy metabolism (1). Namely, with unbalanced sympathetic activation, excess catecholamines together with increased values of pancreastatin reduce glucose intake (by ~50%) and increase the release of free fatty acids (by 4.5 to 6.4 times) (1). Individual studies indicate that pancreastatin (by a mechanism that is not entirely clear) plays a significant role in the pathogenesis of various forms of diabetes mellitus (9).

Catestatin acts on nicotinic acetylcholine receptors as a strong autocrine inhibitor of catecholamine secretion (1,7). It is believed that catestatin deficiency plays a significant role in the development of hypertension caused by long-term overactivity of the sympathetic nervous system (1). By inducing the release of histamine from mast cells, it achieves a vasorelaxant and antihypertensive effect (1). Catestatin has pronounced angiogenic and vasculogenic activity, since it induces the migration and proliferation of endothelial and smooth muscle cells of blood vessels (1). It stimulates chemotaxis activity and the activity of pro-inflammatory cytokines, and inhibits the growth of fungi, yeasts and bacteria (Gram-positive and Gram-negative) (1).

Parastatin was discovered in the parathyroid gland of a pig (1). In the presence of low levels of calcium in the blood, it inhibits the secretion of parathyroid hormone (1).

Prochromacin and the biologically active peptides derived from it, chromacins 1 and 2 inhibit the growth of fungi and bacteria (7,10).

WE 14 can modulate the value of gonadotropic hormones (7).

In addition, research by eminent authors identified WE 14 as an autoantigen of the pancreas that plays a significant role in the pathogenesis of various forms of diabetes mellitus (9).

METHODS OF DETERMINING THE VALUE OF CHROMOGRANIN A IN THE BLOOD

There are a number of commercially available kits for the determination of CgA in blood (6). They are based on three diagnostic techniques: enzyme linked immunosorbent assay (ELISA), immunoradiometric assay (IRMA) and radioimmunoassay (RIA) (6). Although a significant number of studies recommend RIA, there is no universally accepted diagnostic technique and no standardization of any of the mentioned diagnostic techniques (6,11). Therefore, caution is recommended when trying to compare the results of different diagnostic centers (6). CgA can be determined in plasma or serum (6). The value of CgA in plasma is often significantly higher than the value determined in serum (6). Research by certain authors recommends the determination of CgA in saliva (1).

CHROMOGRANIN A AS A DIAGNOSTIC AND PROGNOSTIC MARKER

An increase in the value of CgA in the blood occurs in certain benign diseases and conditions, malignant diseases as well as iatrogenic (Table 2) (12).

CgA is elevated in individuals using proton pump inhibitors, histamine H2 receptor antagonists, and selective serotonin reuptake inhibitors (12). Therefore, it is recommended to stop using drugs (histamine H2 receptor antagonists for 24 hours, proton pump inhibitors for 14 days) before determining the CgA value in the blood (12). Intense physical activity as well as food intake immediately before testing results in an increase in CgA values in the blood (12). Slightly elevated CgA values were found in menopause (due to increased sympathetic tone) and pregnancy (1).

Table 2. Causes of increased CgA values in the blood (1,12)

Benign diseases and conditions	Iatrogenic causes	Malignant diseases
Pregnancy	Proton pump inhibitors	Gastric cancer
Menopause	Antagonists of histamine H2 receptors	Pancreatic cancer
Chronic atrophic gastritis	Selective serotonin reuptake inhibitors	Hepatocellular carcinoma
Helicobacter pylori infection	Intense physical activity	Breast cancer
Pancreatitis	Recent food intake	Ovarian cancer
Chronic hepatitis		Prostate cancer
Liver cirrhosis		Neuroblastoma
Irritable bowel syndrome		Small cell lung cancer

Ulcerative colitis		Neuroendocrine tumors
Chron's disease		
Arterial hypertension		
Heart failure		
Acute coronary syndrome		
Myocardial infarction		
Giant cell arteritis		
Renal insufficiency		
Chronic bronchitis		
Obstructive lung diseases		
Rheumatoid arthritis		
Systemic lupus erythematosus		
Hyperthyroidism		
Hyperparathyroidism		
Hypercortisolemia		

Chronic atrophic gastritis and gastritis caused by *Helicobacter pylori* can cause an increase in the value of CgA, due to a chronic increase in the value of gastrin in the serum (1). In people with these diseases, predominantly in those treated with proton pump inhibitors, the measurement of CgA values in the blood can contribute to the monitoring of enterochromaffin-like gastric cell hyperplasia (1). Ulcerative colitis and Crohn's disease cause a discrete increase in CgA values in the blood (1).

With organ dysfunction, such as kidney, liver and heart failure, CgA values in the blood can also be significantly increased (1). Very high CgA values in people with kidney failure (comparable to CgA values in people with neuroendocrine tumors) support the important role of the kidneys in removing CgA from the body (2). Its moderately elevated values in people with liver failure indicate either the metabolism of CgA in the liver or the activation of the neuroendocrine system in these people (2). Activation of the neuroendocrine system is also present in people with heart failure, acute coronary syndrome and hypertension (1,2). High CgA values in the blood indicate a high risk of death after myocardial infarction, acute coronary syndrome and heart failure (1).

There are high values of CgA in certain inflammatory diseases, such as rheumatoid arthritis and systemic lupus erythematosus (1). Endocrine disorders associated with elevated CgA values in the blood are hypercortisolemia, hyperthyroidism and hyperparathyroidism (1).

CgA is an important marker of neuroendocrine tumors including carcinoid tumor, pheochromocytoma, medullary carcinoma of the thyroid gland, adenoma of the parathyroid gland, neuroblastoma, gastroenteropancreatic neuroendocrine tumors, multiple endocrine neoplasia syndrome type 1 and bronchopulmonary neuroendocrine tumors (13). The highest reliability of this tumor marker was found in disseminated and well-differentiated neuroendocrine tumors that show intense secretory activity (13).

The sensitivity of CgA in people with neuroendocrine tumors is between 10 and 100%, and its specificity is between 68 and 100% (13). In small tumors (except gastrinoma), such as insulinoma, paraganglioma, carcinoid tumor of the bronchus and pituitary tumor, CgA values can be in the reference range (13). The highest sensitivity of CgA was found in gastrinomas (100%), pheochromocytomas (89%), carcinoid tumors (80%), non-functioning tumors of the endocrine pancreas (69%) and medullary carcinoma of the thyroid gland (50%) (13). In patients with metastatic disease, CgA has a sensitivity of 60 to 100% and its blood level can increase by as much as 100-1000 times (13). The highest values are found in carcinoid tumors with liver metastases (13). In addition to diagnosing neuroendocrine tumors, CgA is important in monitoring response to treatment (13).

In patients with small cell lung cancer with neuroendocrine differentiation, CgA can help monitor response to treatment and detect disease recurrence (13). Although the CgA level in the blood indicates the likely degree of neuroendocrine differentiation in the cancer, its low sensitivity makes it unreliable in detecting the disease (13).

In patients with prostate cancer, determining the CgA level in the blood can help in diagnosing and determining the prognosis of the disease (2). Namely, the CgA level in the blood of patients with they can be elevated even if the value of prostate specific antigen s is normal (13). High CgA values in the blood can predict the absence of cancer response to hormone therapy, that is, a poor prognosis of the disease (13).

CgA immunoreactivity has been established in colorectal and breast cancer cells, but its diagnostic value has not been established (2).

CONCLUSION

CgA and the biologically active peptides derived from it have numerous, sometimes conflicting roles in the cardiovascular, metabolic, immune and diffuse neuroendocrine systems. CgA has been identified as an important immunohistochemical tumor marker of neuroendocrine tumors, prostate cancer and microcellular lung cancer with neuroendocrine differentiation. Its credibility is conditioned by the tumor itself (localization, size, density of secretory granules and secretory activity), dissemination of the tumor and application of the diagnostic technique. Elevated CgA values cannot be used to assess the progression of tumors that are not of neuroendocrine origin. In

chronic heart failure and acute coronary syndrome, CgA is an independent prognostic factor of the disease.

LITERATURE

1. D'amico MA, Ghinassi B, Izzicupo P, Manzoli L, Di Baldassarre A. Biological function and clinical relevance of chromogranin A and derived peptides. *Endocr Connect*. 2014 Apr 29;3(2):R45-54. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC5395093/>
2. Taupenot L, Kimberly HL, O'Connor DT. The Chromogranin–Secretogranin Family. *NEJM*. 2003;348(12):1134-1149. Available from: <https://www.nejm.org/doi/full/10.1056/nejmra021405>
3. Feldman SA, Eiden LE. The chromogranins: Their roles in secretion from neuroendocrine cells and as markers for neuroendocrine neoplasia. *Endocr Pathol*. 2003;14:3–23. Available from: <https://doi.org/10.1385/EP:14:1:3>
4. Bartolomucci A, Possenti R, Mahata SK, Fischer-Colbrie R, Loh YP, Salton SR. The extended granin family: structure, function, and biomedical implications. *Endocr Rev*. 2011 Dec;32(6):755-97. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC3591675/>
5. Estevez-Herrera J, Pardo MR., Dominguez N, Pereda D, Machado JD, Borges R. The role of chromogranins in the secretory pathway. *BioMolecular Concepts*. 2013;4(6):605-609. Available from: https://www.degruyter.com/document/doi/10.1515/bmc-2013-0020/html?srsId=AfmBOoqMHFufhcNduEZBODJaF4PL_ddiPBhRUJL34a8zE-ZqRqx-ZKya2
6. Gut P, Czarnywojtek A, Fischbach J, Bączyk M, Ziemnicka K, Wrotkowska E et al. Chromogranin A - unspecific neuroendocrine marker. Clinical utility and potential diagnostic pitfalls. *Arch Med Sci*. 2016 Feb 1;12(1):1-9. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC4754364/>
7. Díaz Pérez JÁ, Currás Freixes M. Cromogranina A y tumores neuroendocrinos. *Endocrinol Nutr*. 2013;60:386–395. Available from: <https://www.elsevier.es/en-revista-endocrinologia-nutricion-412-articulo-chromogranin-a-neuroendocrine-tumors-S2173509313001372#bib0100>
8. Louthan O. Chromogranin A in Physiology and Oncology. *Folia Biologica (Praha)*. 2011;57:173-181. Available from: <https://fb.cuni.cz/file/5597/FB2011A0026.pdf>
9. Herold Z, Doleschall M, Kovesdi A, Patocs A, Somogyi A. Chromogranin-A and its role in the pathogenesis of diabetes mellitus. *Endokrynologia Polska*. 2018;69(5):598-610. Available from: https://journals.viamedica.pl/endokrynologia_polska/article/view/EP.a2018.0052/45437
10. Scavello F, Kharouf N, Lavalley P, Haikel Y, Schneider F, Metz-Boutigue MH. The antimicrobial peptides secreted by the chromaffin cells of the adrenal medulla link the neuroendocrine and immune systems: From basic to clinical studies. *Front Immunol*.

- 2022 Aug 25;13:977175. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9452953/#B23>
11. Stridsberg M, Eriksson B, Oberg K, Janson ET. A comparison between three commercial kits for chromogranin A measurements. *J Endocrinol.* 2003 May;177(2):337-41. Available from: <https://pubmed.ncbi.nlm.nih.gov/12740022/>
 12. Di Giacinto P, Rota F, Rizza L, Campana D, Isidori A, Lania A et al. Chromogranin A: From Laboratory to Clinical Aspects of Patients with Neuroendocrine Tumors. *Int J Endocrinol.* 2018 Jul 2;2018:8126087. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC6051263/>
 13. Glinicki P, Jeske W. Chromogranin A (CgA) — the influence of various factors *in vivo* and *in vitro*, and existing disorders on it's concentration in blood. *Endokrynologia Polska/ Polish Journal of Endocrinology* 2010; 61(4):384-397. Available from: [file:///C:/Users/W10/Downloads/25370-31124-1-PB%20\(1\).pdf](file:///C:/Users/W10/Downloads/25370-31124-1-PB%20(1).pdf)