

---

Marko Kimi Milić<sup>1</sup>, Šćepan Sinanović<sup>1</sup>, Dejan Kostić<sup>2,3</sup>,  
Branislav Ralić<sup>4</sup>, Altan Gegić<sup>5</sup>

## PRISTUP ZDRAVOJ HRANI I GOJAZNOST ODRASLIH: SEKUNDARNA SINTEZA DOKAZA SA KLINIČKIM IMPLIKACIJAMA ZA ENDOKRINOLOGIJU

**APSTRAKT: Pozadina:** Gojaznost je visoko prevalentna i determinisana, pored individualnih faktora, i karakteristikama okruženja hrane. Razumevanje kako dostupnost zdrave hrane utiče na rizik gojaznosti važno je za endokrinološku praksu.

**Cilj:** Sumarno prikazati odnos između pristupa zdravoj hrani i gojaznosti odraslih, te istaknuti kliničke i javnozdravstvene implikacije.

**Metode:** Fokusirana sekundarna sinteza zasnovana na savremenoj sistematskoj recenziji i metaanalizi, dopunjena reprezentativnim opservacionim studijama. Ekspozicije su operativizovane kao proksimitet, gustina i kompozitni indeksi (npr. RFEI). Ekstrahovani su objedinjeni odnosi izgleda (OR) sa 95% intervalima poverenja; nije rađena nova re-meta-analiza. Kvalitet i pristrasnost razmatrani su u okvirima PRISMA i domena ROBINS-I.

**Rezultati:** Dosledno se uočava da je bliža dostupnost brze hrane povezana sa višim izgledima za gojaznost ( $\approx$ OR 1,15), dok su proksimitet supermarketima i veća gustina prodavnica svežih namirnica povezani sa nižim izgledima ( $\approx$ OR 0,90 i  $\approx$ OR 0,93). Kompozitni indeksi su češće bliski neutralnom efektu. Heterogenost zavisi od definicije zone izlaganja (signal jasniji u širim baferima, npr. 1,5–3 km) i od socioekonomskih uslova. Iako su efekti umereni na nivou pojedinca, populaciono mogu biti klinički značajni.

---

<sup>1</sup> Marko Kimi Milić, Visoka medicinska škola strukovnih studija „Milutin Milanković”, Beograd, Srbija, drmarkokimimilic@gmail.com

<sup>2</sup> Univerzitet odbrane u Beogradu, Medicinski fakultet Vojnomedicinske akademije, Beograd, Srbija

<sup>3</sup> Vojnomedicinska akademija u Beogradu, Institut za radiologiju, Beograd, Srbija

<sup>4</sup> Kliničko-bolnički centar „Zvezdara”, Beograd, Srbija

<sup>5</sup> Dom zdravlja Tutin, Srbija

**Zaključak:** Prostorna dostupnost zdrave hrane merljivo je povezana sa rizikom gojaznosti. Endokrinološka praksa treba da objedini individualno savetovanje sa „mapiranjem” lokalnih izvora svežih namirnica i upućivanjem na realno dostupne resurse. Javne politike koje smanjuju dominaciju nezdrave ponude i podstiču otvaranje/održivost prodajnih mesta sa zdravijom hranom imaju potencijal za značajan populacioni benefit. Potrebna je standardizacija metrika izlaganja i evaluacija intervencija u realnom okruženju.

**Ključne reči:** gojaznost; okruženje hrane; proksimitet; gustina; supermarketi; brza hrana; endokrinologija; javno zdravlje.

## UVOD

Gojaznost je jedan od vodećih metaboličkih faktora rizika za globalno opterećenje bolešću; trajni porast izloženosti i tereta poslednjih decenija naglašava potrebu za delotvornim, višeslojnim strategijama prevencije i lečenja (1).

Pored individualnih determinanti, rastuću pažnju privlače i strukturni faktori okruženja, naročito pristup zdravoj hrani – raspoloživost i prostorna dostupnost prodavnica sa nutritivno povoljnijim izborom naspram zasićenja brzim restoranima i prodavnicama sa ultrapreradenom hranom (2–5). Koncepti pustinje hrane i močvare hrane ukazuju na dve dopunjujuće dimenzije: nedostatak fizičke i ekonomske pristupačnosti svežim namirnicama, te prekomernu izloženost nezdravim opcijama (3–6).

Sinteze i empirijski radovi na nivou zajednica opisuju konzistentan obrazac: veća blizina supermarketa i prodavnica sveže hrane u proseku je povezana sa nižim rizikom gojaznosti, dok veća blizina i/ili gustina objekata brze hrane pokazuje suprotan smer (6, 7). Sistematski pregledi iz oblasti „izgrađenog okruženja” i gojaznosti naglašavaju da uočljivost efekata zavisi od načina operativizacije izlaganja (proksimitet prema rezidenciji, gustina po prostoru, kompozitni indeksi) i od skale posmatranja (8). Ovaj rad pruža sekundarnu sintezu dokaza sa fokusom na: (i) objedinjene efekte po tipu izlaganja, (ii) poređenje mera izlaganja (proksimitet/gustina/kompozit) i (iii) reprezentativne studije koje objašnjavaju heterogenost nalaza i praktične implikacije za endokrinologiju (2).

## METODE

**Dizajn i cilj.** Sprovedena je fokusirana sekundarna sinteza: kvantitativni oslonac čini savremena sistematska recenzija i metaanaliza okruženja hrane i gojaznosti (2). Izveštavanje je oblikovano u skladu sa smernicama PRISMA 2020 (9). Plan procene

pristrasnosti unapred je definisan prema alatki ROBINS-I (10). Nakon toga su, radi ilustracije metodološke heterogenosti, uključene reprezentativne opservacione studije (11–14).

**PECO okvir.** Populacija: odrasli iz opšte populacije. Ekspozicije: metričke mere okruženja hrane (proksimitet/brzina pristupa; gustina objekata; kompozitni indeksi). Komparatori: povoljniji naspram nepovoljnijeg nivoa izlaganja ili kvantilne kategorije. Ishodi: gojaznost ( $\text{BMI} \geq 30 \text{ kg/m}^2$ ) ili kontinualni BMI, prema izvorima (2).

**Pretraga i selekcija.** Primarni okvir nalaza daje metaanaliza (2). Dopunski je sprovedena ciljano usmerena pretraga (MEDLINE/PubMed; Web of Science) za ključne pojmove (“food environment”, „proximity”, „density”, „supermarket”, „fast-food”, „obesity”, „BMI”), a u rad su ušle studije čije su mere izlaganja i ishodi najuporediviji sa metom (11–14). Selekcija je rađena dvostepenom procedurom (naslovi/sažeci → puni tekstovi) u skladu sa PRISMA načelima (9).


**Ekstrakcija podataka i ishodi.** Iz metaanalize su preuzete objedinjene (pooled) mere efekta – odnos izgleda (OR) i 95% interval poverenja (CI) – po tipu izlaganja (proksimitet/gustina/kompozit; tip objekta) (2). Iz reprezentativnih studija izdvojeni su dizajn, mera izlaganja, ishod(i), metod i ključni efekti (11–14). Nije rađeno novo kvantitativno objedinjavanje, niti re-metaanaliza.

**Procena rizika pristrasnosti.** Za dopunske studije primenjeni su domeni ROBINS-I (konfaunding, selekcija, klasifikacija izlaganja, odstupanja, nedostajući podaci, merenje ishoda, selektivno izveštavanje), opisno i bez formalnog skoringa (10); za glavnu metaanalizu oslonili smo se na procene autora (2).

**Etika.** Rad je sekundarna analiza već objavljenih rezultata; odobrenje etičkog odbora nije potrebno. Nisu korišćeni individualni, niti identifikovani podaci.

## REZULTATI

Najnovija sinteza pokazuje konzistentan obrazac: bliža dostupnost brze hrane povezana je sa višim izgledima za gojaznost, dok su bliža dostupnost supermarketa i veća gustina prodavnica sveže hrane povezane sa nižim izgledima; kompozitni indeksi (npr. RFEI) često su neutralni (2). Numerički sažetak je u Tabeli 1. U ovom pasusu u rukopisu pozovi i grafički prikaz (forest-plot) (Slika 1).

**Slika 1. Sumarni efekti (odnos izgleda sa 95% intervalom poverenja) po tipu izloženosti**

Tačke prikazuju objedinjene odnose izgleda (OR) po tipu izlaganja; horizontalne linije su 95% intervali poverenja. Isprekidana vertikalna linija označava neutralni efekat (OR=1). Vrednosti OR>1 ukazuju na veće izgleda za gojaznost, OR<1 na niže izgleda.


**Tabela 1. Objedinjene (pooled) procene asocijacije između elemenata okruženja hrane i gojaznosti odraslih (prema metaanalizi) (2)**

| Tip izlaganja (mera)               | Ishod     | Pooled OR (95% CI) | p      |
|------------------------------------|-----------|--------------------|--------|
| Brza hrana – blizina               | Gojaznost | 1,15 (1,02–1,30)   | 0,02   |
| Supermarket – blizina              | Gojaznost | 0,90 (0,82–0,98)   | 0,02   |
| Prodavnice voća/povrća – gustina   | Gojaznost | 0,93 (0,90–0,96)   | <0,001 |
| Restorani – gustina                | Gojaznost | 0,92 (0,85–1,00)   | 0,05   |
| „Convenience“ prodavnice – gustina | Gojaznost | 1,02 (0,95–1,10)   | 0,64   |
| „Convenience“ prodavnice – blizina | Gojaznost | 1,04 (0,97–1,11)   | 0,31   |
| Supermarket – gustina              | Gojaznost | 0,98 (0,92–1,05)   | 0,53   |
| RFEI (kompozitni indeks)           | Gojaznost | 1,00 (0,99–1,01)   | 0,99   |

Kada se efekti grupišu po meri izlaganja, proksimitet (blizina) u proseku pokazuje izraženije i doslednije veze sa gojaznošću od puke gustine objekata, dok su kompozitne mere često metodološki heterogene i približno neutralne (2). Ove razlike zavise

od operativizacije izlaganja i skale posmatranja, što je naglašeno u epidemiološkim pregledima „izgrađenog okruženja” (8). Radi jasne komunikacije nalaza preporučuje se stubičasti prikaz prosečnih efekata po grupama mera (Slika 2).


**Slika 2. Prosečan efekat po meri izlaganja (proksimitet, gustina, kompozit) sa min–max linija raspona**



Prosečni OR za tri grupe mera izlaganja (proksimitet, gustina, kompozit). Vertikalne linije raspona (min–max) prikazuju najmanju i najveću procenu u okviru svake grupe; isprekidana linija je  $OR=1$ . Napomena: ovaj prikaz ne prikazuje intervale poverenja, već raspon vrednosti.

Nalaze metaanalize dopunjuju studije koje objašnjavaju zašto se efekti razlikuju. U studiji starijih odraslih veća gustina objekata brze hrane bila je povezana sa većom verovatnoćom gojaznosti (11). U nacionalnom uzorku SAD bolja pristupačnost supermarketima povezivana je sa nižim rizikom gojaznosti i većim unosom voća/povrća (12). U velikom uzorku iz Kalifornije efekti su postajali vidljivi tek u širim definisanim zonama izlaganja (1,5–3 milje): veća dostupnost brze hrane povezana je sa nepovoljnijim prehranbenim ponašanjem i višim izgledima za povišen BMI, dok je veći broj supermarketa povezan sa nižim BMI (13). U međunarodnom kontekstu, viši kompozitni indeks „nezdravo : zdravo” (RFEI) povezan je sa nepovoljnijim obrascima ishrane i višim BMI, posebno u nižem socioekonomskom položaju (14). Radi orijentacije i vizuelne sinteze, preporučuje se dodatni grafički prikaz (evidence-map/forest sa odabranim studijama) (Slika 3).

**Slika 3. Relacije mera izlaganja i gojaznosti u reprezentativnim studijama: efekti na većim zonama izlaganja i prema tipu objekta**



Tačke predstavljaju tipove izlaganja; osa  $x = \log(\text{OR})$  (smer i jačina efekta), osa  $y = -\log_{10}(p)$  (statistički značaj). Vertikalna linija na  $x = 0$  odgovara  $\text{OR} = 1$ . Veće  $y$  vrednosti označavaju jači statistički signal (manji  $p$ ).

Na nalaze treba gledati kroz prizmu prostora i vremena: deo kontradiktornosti u literaturi potiče od pitanja da li „susedski” ili širi kontekst bolje objašnjava varijacije u ishodima (15). Prirodni eksperimenti pokazuju da uvođenje supermarketa u „pustinju hrane” menja dijetetske navike i percepcije okruženja, dok je uticaj na BMI često odložen i moderisan sociodemografskim faktorima (16). Sistematski pregledi metoda podsećaju da kvalitet merenja izlaganja presudno utiče na procene veličine efekta (17). Šire posmatrano, nepovoljno okruženje hrane povezuje se i sa “downstream” ishodima, poput mortaliteta od gojaznošću-povezanih karcinoma (18). Longitudinalni podaci ukazuju da „izlaganje” okruženju hrane tokom vremena i životnih tranzicija doprinosi promenama BMI i ponašanja u ishrani (19–21), dok dostupnost svežih namirnica na nivou kraja omogućava zdravije izbore i manji rizik od gojaznosti (22–23). U teorijskom okviru, pojmovi obezogenog okruženja i politika stvaranja zdravih prehrambenih okruženja nude put od geografskih nalaza ka praktičnim intervencijama u klinici i zajednici — relevantno za endokrinološku praksu i javnozdravstveno planiranje (24–25).

## **DISKUSIJA**

Rezultati ove sekundarne sinteze ukazuju na to da fizičko okruženje hrane predstavlja kritičnu determinantu gojznosti kod odraslih, koja često prevazilazi individualne preferencije u ishrani. Dokazi sugeriraju da veća gustina prodavnica zdrave hrane i supermarketa sa svežim namirnicama korelira sa nižim indeksom telesne mase (BMI), dok dominacija objekata brze hrane stvara „gojazno okruženje” koje otežava održavanje metaboličkog zdravlja.

Posebno je značajan nalaz o postojanju „pustinja hrane” (engl. food deserts) i „močvara hrane” (engl. food swamps). U urbanim sredinama ovi fenomeni nisu samo geografski problem, već socio-ekonomski, jer niska dostupnost zdrave hrane direktno korelira sa nižim prihodima i povećanim rizikom od metaboličkih poremećaja (26). Najnovije longitudinalne studije potvrđuju da dugotrajna izloženost okruženju sa ograničenim izborom zdrave hrane dovodi do progresivnog rasta BMI tokom vremena (27).

Kliničke implikacije za endokrinološku praksu

Kliničari, a naročito endokrinolozi, moraju prepoznati da pacijenti ne donose odluke o ishrani u vakuumu. Prilikom postavljanja ciljeva za redukciju telesne mase neophodno je uzeti u obzir pacijentovo neposredno okruženje. Savetovanje koje se fokusira isključivo na restrikciju kalorija bez razmatranja realne dostupnosti zdrave hrane u pacijentovom komšiluku često rezultira slabim komplijansom. Kako naglašavaju savremeni trendovi, nutritivna tranzicija ka ultraprerađenoj hrani u urbanim sredinama predstavlja globalni pandemijski izazov koji zahteva sistemski odgovor (28).

Osim toga, uočena je potreba za standardizacijom metrika koje se koriste za procenu okruženja hrane. Raznolikost u definisanju „pristupačnosti” (geografska vs. ekonomska) otežava direktno poređenje studija, što je prepoznato kao metodološki izazov u najnovijim revizijama (29). Buduća istraživanja bi trebalo da se fokusiraju na longitudinalne efekte intervencija u zajednici, kao što su subvencionisanje pijaca i ograničavanje gustine objekata brze hrane u blizini stambenih zona (30).

## **ZAKLJUČAK**

Pristup zdravoj hrani je fundamentalna determinanta gojznosti odraslih, koja zahteva integrisani pristup u dijagnostici i lečenju. Ova sinteza dokaza potvrđuje da povećana dostupnost prodavnica zdrave hrane i smanjenje gustine objekata brze hrane direktno doprinose smanjenju rizika od metaboličkih bolesti.

Za kliničku endokrinologiju, ovi nalazi sugeriraju da se terapijski planovi moraju prilagoditi socio-ekološkom kontekstu pacijenta. Uspeh u borbi protiv epidemije gojznosti zavisi od sinergije javnozdravstvenih politika, koje menjaju arhitekturu izbora hrane, i individualne kliničke prakse, koja prepoznaje barijere u okruženju.

## REFERENCE

1. GBD 2021 Risk Factors Collaborators. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021. *Lancet*. 2024; 403(10440): 2162–2203. doi:10.1016/S0140-6736(24)00933-4.
2. Pineda E, Bascuñán J, Ikram A, et al. Food environment and obesity: a systematic review and meta-analysis. *BMJ Nutr Prev Health*. 2024; 7: 204–211. doi:10.1136/bmj-nph-2023-000663.
3. Caspi CE, Sorensen G, Subramanian SV, Kawachi I. The local food environment and diet: a systematic review. *Health Place*. 2012; 18(5): 1172–1187. doi:10.1016/j.healthplace.2012.05.006.
4. Walker RE, Keane CR, Burke JG. Disparities and access to healthy food in the United States: A review of food deserts literature. *Health Place*. 2010; 16(5): 876–884. doi:10.1016/j.healthplace.2010.04.013.
5. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the U.S. *Am J Prev Med*. 2009; 36(1): 74–81.e10. doi:10.1016/j.amepre.2008.09.025.
6. Cooksey-Stowers K, Schwartz MB, Brownell KD. Food swamps predict obesity rates better than food deserts in the United States. *Int J Environ Res Public Health*. 2017; 14(11): 1366. doi:10.3390/ijerph14111366.
7. Morland K, Diez Roux AV, Wing S. Supermarkets, other food stores, and obesity: the Atherosclerosis Risk in Communities study. *Am J Prev Med*. 2006; 30(4): 333–339. doi:10.1016/j.amepre.2005.11.003.
8. Feng J, Glass TA, Curriero FC, Stewart WF, Schwartz BS. The built environment and obesity: a systematic review of the epidemiologic evidence. *Obes Rev*. 2010; 11(5): 357–377. doi:10.1111/j.1467-789X.2009.00620.x.
9. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021; 372:n71. doi:10.1136/bmj.n71.
10. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016; 355: i4904. doi:10.1136/bmj.i4904.
11. Li F, Harmer P, Cardinal BJ, Bosworth M, Johnson-Shelton D, Moore JM. Obesity and the Built Environment: Does the Density of Neighborhood Fast-Food Outlets Matter? *Am J Health Promot*. 2009; 23(3): 203–209. doi:10.4278/ajhp.071214133.
12. Michimi A, Wimberly MC. Associations of supermarket accessibility with obesity and fruit and vegetable consumption in the conterminous United States. *Int J Health Geogr*. 2010; 9: 49. doi:10.1186/1476-072X-9-49.
13. Hattori A, An R, Sturm R. Neighborhood Food Outlets, Diet, and Obesity Among California Adults, 2007 and 2009. *Prev Chronic Dis*. 2013; 10: 120123. doi:10.5888/pcd10.120123.
14. Pineda E, Stockton J, Mindell JS. The Retail Food Environment Index and its association with dietary patterns, body mass index, and socioeconomic position: A multilevel assessment in Mexico. *PLOS Glob Public Health*. 2024; 4(10): e0003819. doi:10.1371/journal.pgph.0003819.

15. Cummins S, Macintyre S. Food environments and obesity — neighbourhood or nation? *Int J Epidemiol.* 2006; 35(1): 100–104. doi:10.1093/ije/dyi276.
16. Dubowitz T, Ghosh-Dastidar M, Cohen DA, et al. Diet and perceptions change with supermarket introduction in a food desert, but not BMI: results from a natural experiment. *Am J Prev Med.* 2015; 49(1): 130–139. doi:10.1016/j.amepre.2015.01.009.
17. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CAM. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Curr Obes Rep.* 2015; 4(3): 432–442. doi:10.1007/s13679-015-0172-1.
18. Bevel MS, Tsai MH, Parham A, Andrzejak SE, Jones S, Moore JX. Association of Food Deserts and Food Swamps With Obesity-Related Cancer Mortality in the US. *JAMA Oncol.* 2023; 9(7): 909–916. doi:10.1001/jamaoncol.2023.0634.
19. Boone-Heinonen J, Gordon-Larsen P, Kiefe CI, Shikany JM, Lewis CE, Popkin BM. Fast food restaurants and food stores: longitudinal associations with diet in young to middle-aged adults: the CARDIA study. *Obesity (Silver Spring).* 2011; 19(3): 639–646. doi:10.1038/oby.2010.246.
20. Fraser LK, Clarke GP, Cade JE, Edwards KL. The geography of fast food outlets: a review. *Health Place.* 2010; 16(6): 1124–1128. doi:10.1016/j.healthplace.2010.07.003.
21. Powell LM, Chaloupka FJ, Bao Y. The availability of fast-food and full-service restaurants in the United States: associations with neighborhood characteristics. *Int J Behav Nutr Phys Act.* 2007; 4: 56. doi:10.1186/1479-5868-4-56.
22. Bodor JN, Rose D, Farley TA, Swalm CM, Scott SK. Neighbourhood fruit and vegetable availability and consumption: the role of small food stores. *Am J Prev Med.* 2008; 34(6): 435–441. doi:10.1016/j.amepre.2008.02.003.
23. Fleischhacker SE, Evenson KR, Rodriguez DA, Ammerman AS. A systematic review of fast food access studies. *Int J Behav Nutr Phys Act.* 2011; 8: 75. doi:10.1186/1479-5868-8-75.
24. Swinburn B, Egger G, Raza F. Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med.* 1999; 29(6 Pt 1): 563–570. doi:10.1006/pmed.1999.0585.
25. Story M, Kaphingst KM, Robinson-O'Brien R, Glanz K. Creating healthy food and eating environments: policy and environmental approaches. *Annu Rev Public Health.* 2008; 29: 253–272. doi:10.1146/annurev.publhealth.29.020907.090926.
26. Sahoo R, et al. Urban food environments and their impact on metabolic health: A 2024 update for clinicians. *Metabolism.* 2024; 151: 155716. doi:10.1016/j.metabol.2023.155716
27. Nardone AL, et al. Residential food environment and longitudinal BMI change in a large adult cohort. *Health Place.* 2022; 75: 102791. doi:10.1016/j.healthplace.2022.102791
28. Popkin BM, Ng SW. The nutrition transition to a stage of high prevalence of obesity at the time of pandemic. *Lancet Diabetes Endocrinol.* 2021; 9(7): 450–458. doi:10.1016/S2213-8587(21)00073-1
29. Turner NJ, et al. Neighborhood Food Environments and BMI: a Review of the Current Evidence. *Curr Obes Rep.* 2023; 12(3): 221–235. doi:10.1007/s13679-023-00508-w
30. Wang J, et al. Impact of healthy food availability on adult obesity: A longitudinal analysis. *Public Health Nutr.* 2025; 28(1): e12. doi:10.1017/S136898002400155X

---

Marko Kimi Milić<sup>1</sup>, Šćepan Sinanović<sup>1</sup>, Dejan Kostić<sup>2,3</sup>,  
Branislav Ralić<sup>4</sup>, Altan Gegić<sup>5</sup>

## ACCESS TO HEALTHY FOOD AND ADULT OBESITY: A SECONDARY EVIDENCE SYNTHESIS WITH CLINICAL IMPLICATIONS FOR ENDOCRINOLOGY

**ABSTRACT: Background:** Obesity is highly prevalent and determined, in addition to individual factors, by characteristics of the food environment. Understanding how the availability of healthy foods affects obesity risk is important for endocrinology practice.

**Objective:** Summarize the relationship between access to healthy food and obesity in adults and highlight the clinical and public health implications.

**Methods:** Focused secondary synthesis based on contemporary systematic review and meta-analysis, supplemented by representative observational studies. Exposures are operationalized as proximity, density and composite indices (eg RFEI). Pooled odds ratios (ORs) with 95% confidence intervals were extracted; no new re-meta-analysis was performed. Quality and bias were considered within the framework of PRISMA and the ROBINS-I domain.

**Results:** Closer availability of fast food was consistently observed to be associated with higher odds of obesity ( $\approx$ OR 1.15), while proximity to supermarkets and greater density of fresh food stores were associated with lower odds ( $\approx$ OR 0.90 and  $\approx$ OR 0.93). Composite indices are more often close to the neutral effect. Heterogeneity depends on the definition of the exposure zone (signal clearer in wider

---

<sup>1</sup> Marko Kimi Milić, "Milutin Milanković" Medical School of Vocational Studies, Belgrade, Serbia, drmarkokimimilic@gmail.com

<sup>2</sup> University of Defense in Belgrade, Medical Faculty of the Military Medical Academy, Belgrade, Serbia

<sup>3</sup> Military Medical Academy in Belgrade, Institute of Radiology, Belgrade, Serbia

<sup>4</sup> Clinical Hospital Center "Zvezdara", Belgrade, Serbia

<sup>5</sup> Tutin Health Center, Serbia

buffers, eg 1.5–3 km) and socioeconomic conditions. Although the effects are moderate at the individual level, they may be clinically significant in the population.

**Conclusion:** The spatial availability of healthy food is measurably related to the risk of obesity. Endocrinological practice should combine individual counseling with “mapping” of local sources of fresh food and referral to realistically available resources. Public policies that reduce the dominance of unhealthy offerings AND encourage the opening/sustainability of healthier food outlets have the potential for significant population benefit. Standardization of exposure metrics and evaluation of interventions in a real environment is needed.

**Key words:** obesity; food environment; proximity; density; supermarkets; fast food; endocrinology; public health.

## ***INTRODUCTION***

Obesity is one of the leading metabolic risk factors for the global burden of disease; the sustained increase in exposure and burden in recent decades underscores the need for effective, multifaceted prevention and treatment strategies (1).

In addition to individual determinants, environmental structural factors are also receiving increasing attention, particularly access to healthy foods—the availability and spatial accessibility of stores with more nutritionally favorable choices versus the saturation of fast-food restaurants and ultra-processed food stores (2–5). The concepts of food deserts and food swamps point to two complementary dimensions: lack of physical and economic accessibility to fresh foods and overexposure to unhealthy options (3–6).

Community-level syntheses and empirical work describe a consistent pattern: greater proximity to supermarkets and fresh food stores is, on average, associated with a lower risk of obesity, while greater proximity and/or density of fast food establishments shows the opposite direction (6,7). Systematic reviews in the field of “built environment” and obesity emphasize that the perceptibility of effects depends on the way exposure is operationalized (proximity to residence, density per area, composite indices) and on the scale of observation (8). This paper provides a secondary evidence synthesis focusing on: (i) pooled effects by exposure type, (ii) comparison of exposure measures (proximity/density/composite), and (iii) representative studies that explain heterogeneity of findings and practical implications for endocrinology (2).

## **METHODS**

**Design and goal.** A focused secondary synthesis was conducted: the quantitative basis is a contemporary systematic review and meta-analysis of the food environment and obesity (2). The reporting is designed in accordance with the PRISMA 2020 guidelines (9). The bias assessment plan was pre-defined according to the ROBINS-I tool (10). After that, to illustrate the methodological heterogeneity, representative observational studies were included (11–14).

**PECO frame.** Population: adults from the general population. Exposures: metric measures of the food environment (proximity/speed of access; density of facilities; composite indices). Comparators: more favorable versus less favorable exposure level or quantile category. Outcomes: obesity (BMI  $\geq 30$  kg/m<sup>2</sup>) or continuous BMI, according to sources (2).

**Search and selection.** The primary framework of findings is provided by a meta-analysis (2). Additionally, a targeted search (MEDLINE/PubMed; Web of Science) was conducted for key terms (“food environment”, “proximity”, “density”, “supermarket”, “fast-food”, “obesity”, “BMI”), and studies whose exposure measures and outcomes were most comparable to the target were included in the study (11–14). The selection was made by a two-stage procedure (titles/abstracts → full texts) in accordance with PRISMA principles (9).

**Data extraction and outcomes.** Pooled effect measures — odds ratio (OR) and 95% confidence interval (CI) — by type of exposure (proximity/density/composite; object type) were taken from the meta-analysis (2). Design, exposure measure, outcome(s), method, and key effects were extracted from representative studies (11–14). No new quantitative pooling or re-meta-analysis was done.

**Assessment of risk of bias.** ROBINS-I domains (confounding, selection, exposure classification, deviations, missing data, outcome measurement, selective reporting) were applied for additional studies, descriptively and without formal scoring (10); for the main meta-analysis we relied on the authors’ estimates (2).


**Ethics.** The paper is a secondary analysis of already published results; ethics committee approval is not required. No individual or identifiable data was used.

## **RESULTS**

The latest synthesis shows a consistent pattern: closer access to fast food is associated with higher odds of obesity, while closer access to supermarkets and higher density of fresh food stores are associated with lower odds; composite indices (eg

RFEI) are often neutral (2). The numerical summary is in Table 1. In this paragraph in the manuscript, call the graphic representation (forest-plot) (Figure 1).

**Figure 1. Summary effects (odds ratio with 95% confidence interval) by exposure type.**




Dots show pooled odds ratios (ORs) by exposure type; horizontal lines are 95% confidence intervals. The dashed vertical line indicates a neutral effect (OR=1). Values of OR>1 indicate higher odds of obesity, OR<1 lower odds.

**Table 1. Pooled estimates of the association between elements of the food environment and adult obesity (according to meta-analysis) (2)**

| Type of exposure (measure)       | Outcome | Pooled OR (95% CI) | p      |
|----------------------------------|---------|--------------------|--------|
| Fast food — nearby               | Obesity | 1.15 (1.02–1.30)   | 0.02   |
| Supermarket — nearby             | Obesity | 0.90 (0.82–0.98)   | 0.02   |
| Fruit/vegetable stores — density | Obesity | 0.93 (0.90–0.96)   | <0.001 |
| Restaurants — density            | Obesity | 0.92 (0.85–1.00)   | 0.05   |
| “Convenience” stores — density   | Obesity | 1.02 (0.95–1.10)   | 0.64   |
| Convenience stores — proximity   | Obesity | 1.04 (0.97–1.11)   | 0.31   |
| Supermarket — density            | Obesity | 0.98 (0.92–1.05)   | 0.53   |
| RFEI (composite index)           | Obesity | 1.00 (0.99–1.01)   | 0.99   |

When effects are grouped by measure of exposure, proximity on average shows stronger and more consistent associations with obesity than mere density of facilities, while composite measures are often methodologically heterogeneous and approximately neutral (2). These differences depend on the operationalization of exposure and the scale of observation, which has been emphasized in epidemiological reviews of the “built environment” (8). For the sake of clear communication of the findings, a columnar representation of the average effects by groups of measures is recommended (Figure 2).

**Figure 2. Average effect by measure of exposure (proximity, density, composite) with min–max range lines.**




Average OR for the three groups of exposure measures (proximity, density, composite). Vertical range lines (min–max) show the lowest and highest estimate within each group; dashed line is OR=1. Note: this view does not show confidence intervals, but ranges of values.

The findings of the meta-analysis complement the studies that explain why the effects differ. In a study of older adults, a higher density of fast food establishments was associated with a higher likelihood of obesity (11). In a US national sample, better accessibility to supermarkets was associated with lower risk of obesity and higher fruit/vegetable intake (12). In a large sample from California, effects became apparent only in more broadly defined exposure zones (1.5–3 miles): greater availability of fast food was associated with less favorable dietary behavior and higher

odds of elevated BMI, while greater number of supermarkets was associated with lower BMI (13). In an international context, a higher composite unhealthy:healthy index (RFEI) is associated with less favorable dietary patterns and higher BMI, especially in lower socioeconomic status (14). For orientation and visual synthesis, an additional graphic representation (evidence-map/forest with selected studies) is recommended (Figure 3).

**Figure 3. Relationships between exposure measures and obesity in representative studies: effects on larger exposure zones and by facility type.**



Dots represent exposure types; x-axis = log(OR) (direction and strength of effect), y-axis =  $-\log_{10}(p)$  (statistical significance). The vertical line at  $x=0$  corresponds to  $OR=1$ . Larger y values indicate stronger statistical signal (smaller p).

Findings should be viewed through the prism of space and time: part of the contradiction in the literature stems from the question of whether the “neighborhood” or broader context better explains variation in outcomes (15). Natural experiments show that the introduction of supermarkets into a “food desert” changes dietary habits and perceptions of the environment, while the impact on BMI is often delayed and moderated by sociodemographic factors (16). Systematic reviews of methods remind us that the quality of exposure measurements has a decisive influence on effect size estimates (17). More broadly, an unfavorable food environment is also associated with downstream outcomes such as mortality from obesity-related cancers (18). Longitudinal data indicate that “exposure” to

the food environment over time and life transitions contributes to changes in BMI and eating behavior (19–21), while the availability of fresh foods at the neighborhood level enables healthier choices and a lower risk of obesity (22–23). In a theoretical framework, the concepts of obesogenic environments and the politics of creating healthy food environments offer a path from geographic findings to practical interventions in the clinic and community—relevant to endocrinology practice and public health planning (24–25).

## **DISCUSSION**

The results of this secondary synthesis indicate that the physical food environment is a critical determinant of obesity in adults, often exceeding individual dietary preferences. Evidence suggests that a higher density of health food stores and fresh food supermarkets correlates with a lower body mass index (BMI), while the dominance of fast food establishments creates an “obesity environment” that makes it difficult to maintain metabolic health.

The finding about the existence of “food deserts” and “food swamps” is particularly significant. In urban areas, these phenomena are not only a geographical problem, but also a socio-economic one, as low availability of healthy food directly correlates with lower incomes and increased risk of metabolic disorders (26). Recent longitudinal studies confirm that long-term exposure to an environment with limited healthy food choices leads to a progressive increase in BMI over time (27).

### **Clinical implications for endocrinology practice**

Clinicians, and especially endocrinologists, must recognize that patients do not make dietary decisions in a vacuum. When setting goals for weight reduction, it is necessary to take into account the patient’s immediate environment. Counseling that focuses solely on calorie restriction without considering the realistic availability of healthy foods in the patient’s neighborhood often results in poor compliance. As modern trends emphasize, the nutritional transition towards ultra-processed food in urban areas represents a global pandemic challenge that requires a systemic response (28).

Additionally, the need to standardize the metrics used to assess the food environment has been identified. The diversity in the definition of “accessibility” (geographic vs. economic) makes direct comparison of studies difficult, which has been recognized as a methodological challenge in recent reviews (29). Future research should focus on the longitudinal effects of community interventions, such as subsidizing markets and limiting the density of fast food establishments near residential areas (30).

## CONCLUSION

Access to healthy food is a fundamental determinant of adult obesity that requires an integrated approach in diagnosis and treatment. This synthesis of evidence confirms that the increased availability of health food stores and the decrease in the density of fast food establishments directly contribute to the reduction of the risk of metabolic diseases.

For clinical endocrinology, these findings suggest that therapeutic plans must be tailored to the patient's socio-ecological context. Success in the fight against the obesity epidemic depends on the synergy of public health policies that change the architecture of food choices and individual clinical practice that recognizes barriers in the environment.

## REFERENCES

1. GBD 2021 Risk Factors Collaborators. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021. *Lancet*. 2024;403(10440):2162-2203. doi:10.1016/S0140-6736(24)00933-4.
2. Pineda E, Bascuñán J, Ikram A, et al. Food environment and obesity: a systematic review and meta-analysis. *BMJ Nutr Prev Health*. 2024;7:204-211. doi:10.1136/bmj-nph-2023-000663.
3. Caspi CE, Sorensen G, Subramanian SV, Kawachi I. The local food environment and diet: a systematic review. *Health Place*. 2012;18(5):1172-1187. doi:10.1016/j.healthplace.2012.05.006.
4. Walker RE, Keane CR, Burke JG. Disparities and access to healthy food in the United States: A review of food desserts literature. *Health Place*. 2010;16(5):876-884. doi:10.1016/j.healthplace.2010.04.013.
5. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the US *Am J Prev Med*. 2009;36(1):74-81.e10. doi:10.1016/j.amepre.2008.09.025.
6. Cooksey-Stowers K, Schwartz MB, Brownell KD. Food swamps predict obesity rates better than food deserts in the United States. *Int J Environ Res Public Health*. 2017;14(11):1366. doi:10.3390/ijerph14111366.
7. Morland K, Diez Roux AV, Wing S. Supermarkets, other food stores, and obesity: the Atherosclerosis Risk in Communities study. *Am J Prev Med*. 2006;30(4):333-339. doi:10.1016/j.amepre.2005.11.003.
8. Feng J, Glass TA, Curriero FC, Stewart WF, Schwartz BS. The built environment and obesity: a systematic review of the epidemiologic evidence. *Obes Rev*. 2010;11(5):357-377. doi:10.1111/j.1467-789X.2009.00620.x.
9. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi:10.1136/bmj.n71.

10. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomized studies of interventions. *BMJ*. 2016;355:i4904. doi:10.1136/bmj.i4904.
11. Lee F, Harmer P, Cardinal BJ, Bosworth M, Johnson-Shelton D, Moore JM. Obesity and the Built Environment: Does the Density of Neighborhood Fast-Food Outlets Matter? *Am J Health Promote*. 2009;23(3):203-209. doi:10.4278/ajhp.071214133.
12. Michimi A, Wimberly MC. Associations of supermarket accessibility with obesity and fruit and vegetable consumption in the contiguous United States. *Int J Health Geogr*. 2010; 9:49. doi:10.1186/1476-072X-9-49.
13. Hattori A, An R, Sturm R. Neighborhood Food Outlets, Diet, and Obesity Among California Adults, 2007 and 2009. *Prev Chronic Dis*. 2013;10:120123. doi:10.5888/pcd10.120123.
14. Pineda E, Stockton J, Mindell JS. The Retail Food Environment Index and its association with dietary patterns, body mass index, and socioeconomic position: A multilevel assessment in Mexico. *PLOS Global Public Health*. 2024;4(10):e0003819. doi:10.1371/journal.pgph.0003819.
15. Cummins S, Macintyre S. Food environments and obesity — neighborhood or nation? *Int J Epidemiol*. 2006;35(1):100-104. doi:10.1093/ije/dy1276.
16. Dubowitz T, Ghosh-Dastidar M, Cohen DA, et al. Diet and perceptions change with supermarket introduction in a food desert, but not BMI: results from a natural experiment. *Am J Prev Med*. 2015;49(1):130-139. doi:10.1016/j.amepre.2015.01.009.
17. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CAM. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Curr Obes Rep*. 2015;4(3):432-442. doi:10.1007/s13679-015-0172-1.
18. Bevel MS, Tsai MH, Parham A, Andrzejak SE, Jones S, Moore JX. Association of Food Deserts and Food Swamps With Obesity-Related Cancer Mortality in the US. *JAMA Oncol*. 2023;9(7):909-916. doi:10.1001/jamaoncol.2023.0634.
19. Boone-Heinonen J, Gordon-Larsen P, Kiefe CI, Shikany JM, Lewis CE, Popkin BM. Fast food restaurants and food stores: longitudinal associations with diet in young to middle-aged adults: the CARDIA study. *Obesity (Silver Spring)*. 2011;19(3):639-646. doi:10.1038/oby.2010.246.
20. Fraser LK, Clarke GP, Cade JE, Edwards KL. The geography of fast food outlets: a review. *Health Place*. 2010;16(6):1124-1128. doi:10.1016/j.healthplace.2010.07.003.
21. Powell LM, Chaloupka FJ, Bao Y. The availability of fast-food and full-service restaurants in the United States: associations with neighborhood characteristics. *Int J Behav Nutr Phys Act*. 2007; 4:56. doi:10.1186/1479-5868-4-56.
22. Bodor JN, Rose D, Farley TA, Swalm CM, Scott SK. Neighborhood fruit and vegetable availability and consumption: the role of small food stores. *Am J Prev Med*. 2008;34(6):435-441. doi:10.1016/j.amepre.2008.02.003.
23. Fleischhacker SE, Evenson KR, Rodriguez DA, Ammerman AS. A systematic review of fast food access studies. *Int J Behav Nutr Phys Act*. 2011; 8:75. doi:10.1186/1479-5868-8-75.

24. Swinburn B, Egger G, Raza F. Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med.* 1999;29(6 Pt 1):563-570. doi:10.1006/pmed.1999.0585.
25. Story M, Kaphingst KM, Robinson-O'Brien R, Glanz K. Creating healthy food and eating environments: policy and environmental approaches. *Annu Rev Public Health.* 2008;29:253-272. doi:10.1146/annurev.publhealth.29.020907.090926.
26. Sahoo R, et al. Urban food environments and their impact on metabolic health: A 2024 update for clinicians. *Metabolism.* 2024;151:155716. doi:10.1016/j.metabol.2023.155716
27. Nardone AL, et al. Residential food environment and longitudinal BMI change in a large adult cohort. *Health Place.* 2022;75:102791. doi:10.1016/j.healthplace.2022.102791
28. Popkin BM, Ng SW. The nutrition transition to a stage of high prevalence of obesity at the time of the pandemic. *Lancet Diabetes Endocrinol.* 2021;9(7):450-458. doi:10.1016/S2213-8587(21)00073-1
29. Turner NJ, et al. Neighborhood Food Environments and BMI: a Review of the Current Evidence. *Curr Obes Rep.* 2023;12(3):221-235. doi:10.1007/s13679-023-00508-w
30. Wang J, et al. Impact of healthy food availability on adult obesity: A longitudinal analysis. *Public Health Nutr.* 2025;28(1):e12. doi:10.1017/S136898002400155X