Originalni naučni rad

**UTICAJ FIZIČKE AKTIVNOSTI NA STEPEN DEPRESIVNOSTI,
ANKSIOZNOSTI I STRESA STUDENATA
MEDICINSKOG FAKULTETA UNIVERZITETA U BEOGRADU¹**

UDK: 613.71-057.875(497.11)

159.944.4.072-057.875

616.89-008.441-057.875

616.89-008.454-057.875

DOI: 10.5937/snp2102057L

Uroš Lazarević²

Medicinski fakultet, Univerzitet u Beogradu, Srbija

Dragana Drljačić³

Visoka sportska i zdravstvena škola, Beograd, Srbija

Miloš Bojović

Visoka sportska i zdravstvena škola, Beograd, Srbija

Srdan Milosavljević

Visoka sportska i zdravstvena škola, Beograd, Srbija

Apstrakt: Problemi sa mentalnim zdravljem postaju sve zastupljeniji i među studentima. Shodno tome, cilj ovog istraživanja bio je da se ispita uticaj fizičke aktivnosti (FA) na ispoljavanje simptoma depresivnosti, anksioznosti i stresa (DAS) studenata medicine. Uzorak ispitanika činilo je 89 studenata (59 ženskog [Ž] i 30 muškog [M] pola), od prve do šeste godine studija Medicinskog fakulteta Univerziteta u Beogradu, starosti $22 \pm 3,3$ godina. Kao merni instrumenti primenjeni su kratka verzija Skale depresivnosti, anksioznosti i stresa (DASS-21) i kratka forma Međunarodnog upitnika o fizičkoj aktivnosti (IPAQ). Dobijeni rezultati ukazuju na to da se prosečni skorovi nalaze u kategoriji koja odgovara normalnom stepenu depresivnosti i anksioznosti i blagom stepenu stresa, pri čemu su za Ž zabeleženi veći skorovi na skali anksioznosti ($\chi^2 = 8,36, p = 0,004$). Između studenata različitih godina studija zabeležene su razlike na skali depresivnosti ($F = 2,515, p = 0,036$) i anksioznosti ($F = 3,594, p = 0,005$), ali ne i na skali stresa ($F = 1,55, p = 0,183$). Ukupna FA odgovarala je kategoriji umerene (medijana [interval] = 2613 [462-11931] MET-min/nedelja), pri čemu M značajno više upražnjavaju aktivnosti visokog intenziteta ($\chi^2 = 7,69, p = 0,006$). Istraživanjem je zabeležena i mala pozitivna korelacija između sedentarnih aktivnosti i simptoma stresa kod Ž ($r = 0,282, p = 0,030$), dok je kod M zabeležena mala negativna korelacija između umerene FA i simptoma depresije ($r = -0,279, p = 0,033$). Rezultati istraživanja ukazuju na to da je ukupna FA i stepen DAS ispitivanih studenata na zadovoljavajućem nivou.

Ključne reči: IPAQ, DASS-21, metabolički ekvivalent, neurotski poremećaji, poremećaji raspoloženja

¹ Rad primljen: 14.12.2021, korigovan: 24.12.2021, prihvaćen za objavljivanje: 24.12.2021.

² Uroš Lazarević je student specijalističkih studija na Medicinskom fakultetu Univerziteta u Beogradu.

³ ✉ dragana.drljadic@vss.edu.rs

UVOD

Fizička aktivnost je svaki pokret skeletnih mišića koji zahteva potrošnju energije (WHO, 2010), odnosno pokret tela proizveden od strane mišićno-koštanog sistema koji ima za posledicu potrošnju energije iznad praga koje telo troši u stanju mirovanja (Caspersen et al., 1985). Ona uključuje različite aktivnosti na dnevnom nivou (od obavljanja kućnih poslova, do rekreativnih aktivnosti i fizičkog vežbanja) (Pedišić, 2004; Stojanović et al., 2013). Poslednjih nekoliko decenija brojne naučne studije proučavale su vezu između fizičke aktivnosti i zdravlja, kao i efekte fizičkog vežbanja na pojedine organske sisteme (Kohl et al., 2013).

Pokazalo se da je uticaj fizičke aktivnosti na zdravlje dovoljno velik, da su istraživači čak i uz relativno grube tehnike merenja bili u stanju da dosledno dokumentuju jaku povezanost fizičkog vežbanja sa zdravstvenom koristi (Siscovick et al., 1985; Sylvia et al., 2014). Pokazana je veza između redovne fizičke aktivnosti sa smanjenjem incidencije određenih oboljenja, kao što su koronarne bolesti (Oguma & Shinoda-Tagawa, 2004), dijabetes (LaMonte et al., 2005), hipertenzija (Diaz & Shimbo, 2013), artritis (Shih et al., 2006), osteoporoza (Castrogiovanni et al., 2016), rak debelog creva (Wolin et al., 2009), depresija (Dinas et al., 2011), gojaznost (Chin et al., 2016) itd. Dokazano je da redovno upražnjavanje fizičke aktivnosti dovodi do poboljšanja određenih procesa u organizmu fizički aktivnih osoba, kao što su metabolizam ugljenih hidrata i masti, hemokoagulacioni procesi, funkcionisanje imunog sistema, poboljšanje kognitivnih funkcija i mentalnog zdravlja, uopšte (Kohl et al., 2013).

S obzirom na to da je jasan pozitivan uticaj fizičke aktivnosti u prevenciji i smanjenju pojave fizičkih oboljenja, svedočimo povećanju interesovanja za njen doprinos i u očuvanju i ublažavanju problema mentalnih oboljenja. Pozitivni efekti fizičke aktivnosti na mentalno zdravlje pokazani su u velikom broju studija (Erić, 2015; Goodwin, 2003). U jednoj od njih koja je uključivala preko 8000 ispitanika, dobijeni su nalazi koji ukazuju na to da redovna fizička aktivnost značajno smanjuje rizik od pojave depresije i anksioznih poremećaja (De Moor et al., 2006). Takođe, interventne studije pokazuju da čak i jednokratna fizička aktivnost dovodi do smanjenja simptoma depresije, anksioznosti i stresa odmah nakon završetka fizičke aktivnosti (Stonerock et al., 2015; Strickland & Smith, 2014).

Svetska zdravstvena organizacija procenjuje da su depresija, anksioznost i stres glavni uzročnici smanjenja kvaliteta življenja i skraćenja životnog veka u današnje vreme (Ostojić i sar., 2009). Usled povećanja incidence ozbiljnih mentalnih oboljenja, postoji bojazan i od povećanja broja osoba koje pate od hroničnih ili ponavljajućih, blagih do umerenih simptoma ovih oboljenja. Već sada u široj javnosti svedočimo povećanoj pojavi mentalnih problema koji se mogu okarakterisati kao blaga depresija, nisko samopoštovanje, visok stres i anksioznost i loše suočavanje sa teškoćama i problemima svakodnevice. Ovim pojavama doprinosi i smanjenje zastupljenosti fizičke aktivnosti u institucijama i promocije fizičke aktivnosti kao kulture življenja (Fox et al., 2000).

Problemi sa mentalnim zdravljem postaju sve zastupljeniji i među studentima. Istraživanja su pokazala da gotovo polovina univerzitetskih studenata pokazuje umerene nivoe mentalnih problema povezanih sa stresom, uključujući i anksioznost i depresiju (Regehr et al., 2013). Nakon što postanu akademski građani, studenti se suočavaju sa povećanim obimom akademskih obaveza i odgovornosti, a samim tim i sa brojnim psihološkim i psihosocijalnim izazovima koji su povezani sa njihovim ličnim razvojem i osamostaljivanjem. S tim u vezi, studije medicine spadaju u najizazovnije u celom svetu, pa i kod nas. Univerzitetski život studenata medicine tokom medicinskog usavršavanja podrazumeva veliku posvećenost i odgovornost u pogledu izvršavanja akademskih zadataka i brige o pacijentima i njihovim porodicama. Veliki broj sati proveden na predavanjima i kliničkoj praksi, okruženje koje često nije idealno za učenje, nedostatak sna kao i faktori koji ometaju svakodnevni lični život, uobičajeni su tokom ovog perioda (Moutinho et al., 2017). Svi ovi činioci utiču na mentalni i emocionalni status studenata (Latas et al., 2010) i uz nedostatak aktivnosti koje poboljšavaju kvalitet života, mogu doprineti razvoju fizičkih, mentalnih i emocionalnih problema.

Istraživanja su pokazala da studenti medicine spadaju u populaciju u kojoj su fenomeni predispitne anksioznosti posebno zastupljeni (Latas et al., 2010). Postoji dosta publikacija na temu predispitne anksioznosti, ali je literatura o uticaju fizičke aktivnosti na simptome depresivnosti, anksioznosti i stresa kod zdrave populacije studenata, prema saznanjima autora ovog rada, oskudna. Iz tog razloga cilj ovog rada bio je da se ispita uticaj fizičke aktivnosti na ispoljavanje simptoma depresivnosti, anksioznosti i stresa studenata medicinskog fakulteta.

METODE

Uzorak ispitanika

U istraživanju je učestvovalo 89 studenata oba pola ($\bar{X}$ = 59 i M = 30) od prve do šeste godine studija Medicinskog fakulteta Univerziteta u Beogradu (I godina, N = 10; II godina, N = 23; III godina, N = 9; IV godina, N = 23; V godina, N = 7 i VI godina, N = 17), prosečne starosti $22 \pm 3,3$, u opsegu od 19 do 35 godina. Putem semi-strukturisanog upitnika, najpre su prikupljeni podaci vezani za socio-demografske varijable (starost, pol, godina studija) i kliničke karakteristike ispitanika, koji su nadalje korišćeni kao kriterijumi za uključivanje ispitanika u istraživanje. Prisustvo bilo kakve psihijatrijske dijagnoze i/ili korišćenje psihofarmaka predstavljali su glavne kriterijume eliminacije iz daljeg toka istraživanja.

Istraživanje je sprovedeno u septembru 2019. godine, po tipu studije preseka i odobreno je od strane Etičkog komiteta Medicinskog fakulteta u Beogradu.

Merni instrumenti

Za merenje stepena depresivnosti, anksioznosti i stresa, primenjena je kratka verzija Skale depresivnosti, anksioznosti i stresa (DASS-21) sačinjena od 21 stavke (Lovibond & Lovibond, 1995). DASS-21 uključuje tri supskale, sa po sedam stavki (skala depresivnosti, npr. *Osećao sam da je život besmislen*; skala anksioznosti, npr. *Osećao sam da sam blizu panike*; skala stresa, npr. *Nerviralo me je kad me nešto prekida u onome što radim*). Skala depresivnosti procenjuje disforiju, beznađe, devalvaciju života, samoregulaciju, nedostatak interesovanja/uključenosti u događaje, anhedoniju i inertnost. Skala anksioznosti procenjuje reakcije autonomnog nervnog sistema, reakcije skeletnih mišića, situacijsku anksioznost i subjektivno iskustvo anksioznog afekta. Skala stresa osetljiva je na nivo hroničnog nespecifičnog uzbuđenja. Procenjuje poteškoće opuštanja, nervno uzbuđenje i lako uznemirenje i nestrpljenje.

Prilikom merenja ispitanici su imali zadatak da na četvorostepenoj skali Likertovog tipa (od 0 – nimalo, do 3 – uglavnom ili skoro uvek) ocene stepen u kom su iskusili svako od navedenih stanja u poslednjih sedam dana (Lovibond & Lovibond, 1995). Rezultati stepena depresivnosti, anksioznost i stresa (DAS) izračunavani su sabiranjem rezultata za relevantne stavke, pri čemu su se skorovi za svaku supskalu kretale u intervalu od 0 do 21. Kriterijumi za određivanje stepena DAS prikazani su u Tabeli 1.

Tabela 1. Kriterijumi za određivanje stepena depresivnosti, anksioznosti i stresa (Lovibond & Lovibond, 1995)

	depresivnost	anksioznost	stres
normalan osećaj	0-4	0-3	0-7
blag osećaj	5-6	4-5	8-9
umereni osećaj	7-10	6-7	10-12
jak osećaj	11-13	8-9	13-16
veoma jak osećaj	≥ 14	≥ 10	≥ 17

Legenda: U kolonama su dati intervali skorova na Skali depresivnosti, anksioznosti i stresa (DASS-21)

U svrhu ispitivanja nivoa fizičke aktivnosti (FA) korišćena je kratka forma Međunarodnog upitnika o fizičkoj aktivnosti (eng. *The International Physical Activity Questionnaire, IPAQ*) (IPAQ Research Committee, 2005), koja meri učestalost, vreme trajanja i intenzitet FA u poslednjih sedam dana. Ova forma IPAQ upitnika namenjena je za procenu FA povezanu sa zdravljem u populaciji starosti od 15 do 69 godina i procenjuje nivo FA kroz tri tipa aktivnosti: laka FA, umerena i intenzivna FA, kao i vreme provedeno u sedentarnim aktivnostima (sedenju/ležanju) (Hagströmer et al., 2006; Papathanasiou et al., 2009). Nivo FA procenjivan je kroz ukupnu energentsku potrošnju izraženu u metaboličkim ekvivalentima (eng. *Metabolic Equivalent of Task*). Jedan metabolički ekvivalent (MET) predstavlja bazalni nivo potrošnje kiseonika i utrošak kalorija koji je za njega vezan (oko 3,5 ml O₂/kg/min ili 1 Kcal/kg/h) i odgovara nivou metabolizma u mirovanju (Ainsworth et al., 2000). Izračunavanje MET rađeno je prema Smernicama za obradu i analizu podataka međunarodnog upitnika o FA (IPAQ Research Committee, 2005).

Vrednosti MET-min/nedelja < 600 smatrane su za nizak nivo energetske potrošnje i odgovarale su kategoriji lake FA. Vrednosti od 601 do 3000 MET-min/nedelja, odgovarale su umerenoj FA, a vrednosti > 3000 MET-min/nedelja, intenzivnoj FA (Craig et al., 2003). Nadalje, na osnovu zbirnog IPAQ skora ispitanici su klasifikovani u grupu aktivnih (MET-min/nedelji $\geq$ 3000) i grupu neaktivnih ispitanika (MET-min/nedelji < 3000).

Statistička analiza

Osnovni deskriptivni pokazatelji (frekvencije i procenti za atributivna obeležja, odnosno $SV \pm SD$, minimum i maksimum za numerička obeležja) izračunavani su za sve varijable. Normalnost distribucije numeričkih obeležja ispitivana je primenom Šapiro-Vilk (Shapiro-Wilk) testa, kojim je zabeleženo odsustvo normalne raspodele za sve praćene varijable. Shodno dobijenim rezultatima u svrhu ispitivanja razlika u stepenu DAS u odnosu na pol, razlika u tipovima FA u odnosu na pol, kao i razlika u stepenu DAS u odnosu na nivo FA između ispitanika istog pola, kao i fizički aktivnih i fizički neaktivnih studenata, sprimenjen je Kruškal-Valis (Kruskal-Wallis) test. S obzirom na to da se rezultati neparametrijske statistike nisu razlikovali od rezultata dobijenih parametrijskim testovima, za ispitivanje razlika u stepenu DAS između ispitanika različitih godina studija, kao i razlika u FA između ispitanika različitih godina studija, u radu su prikazani rezultati dobijeni jednofaktorskom analizom varijanse. U slučaju značajnosti glavnog faktora rađeni su naknadni Fišerovi LSD testovi. Spirmanov koeficijent korelacije ranga primenjen je u svrhu ispitivanja veze između nivoa FA i stepena DAS, testiranjem korelacije između pojedinačnih supskala IPAQ instrumenta i supskala DASS-21. Prema Koenovom kriterijumu vrednosti od 0,10 do 0,29 smatrane su za malu, od 0,30 do 0,49, za srednju i vrednosti od 0,50 do 1,00 za veliku korelaciju (Pallant, 2011). Statistički testovi računati su primenom SPSS 20.0 softvera (SPSS Inc, Chicago, IL), pri čemu je prag značajnosti statističkih analiza bio na nivou poverenja $p = 0,05$.

REZULTATI

Procenom stepena DAS na ukupnom uzorku dobijeni su rezultati koji spadaju u kategoriju normalnog osećaja na skali depresivnosti (medijana i interval, Med [interval] = 3 [0-20]) i anksioznosti (Med [interval] = 3 [0-17]) i blagog osećaja na skali stresa (Med [interval] = 8 [1-20]). Međutim, zabeležene su i visoke maksimalne vrednosti, za sve tri skale, koje spadaju u kategoriju veoma jakog osećaja.

Na Slici 1 prikazani su osnovni deskriptivni pokazatelji (Med i interval) stepena DAS i rezultati poređenja u odnosu na pol, pri čemu se može zapaziti da su sve tri vrste simptoma izraženije kod ispitanika ženskog pola. Međutim, značajne razlike zabeležene su samo kod simptoma anksioznosti ($\chi^2 = 8,36$, $p = 0,004$), ali ne i kod simptoma depresije ($\chi^2 = 0,94$, $p = 0,333$) i stresa ($\chi^2 = 2,54$, $p = 0,111$).

Slika 1. Osnovni deskriptivni pokazatelji (medijana i interval) stepena depresivnosti, anksioznosti i stresa i rezultati poređenja u odnosu na pol

Legenda: Pravougaoni dijagrami predstavljaju interkvartilni raspon, dok središnja linija predstavlja medijanu. Simbolom ** označene su razlike među polovima (M – muški i Ž – ženski) značajne na nivou $p < 0,01$

U Tabeli 2 prikazani su osnovni deskriptivni pokazatelji (Med i interval) stepena DAS, po godinama studija. Na skali depresivnosti najveći broj rezultata nalazi se u okviru normalnog osećaja (skor 0-4), osim za I i V godinu, gde se kreće u okviru umerenog osećaja depresije (7-10). Skorovi anksioznosti kreću se unutar normalnog osećaja (0-3) za III, IV i VI godinu, blagog za V, umerenog (6-7) za I i jakog osećaja anksioznosti za II godinu studija. Skorovi stresa kreću se od normalnog osećaja za III, IV i VI godinu (skor 0-7), blagog za II i V godinu (skor 8-9), do umerenog osećanja stresa (skor 10-11) za I godinu studija.

Tabela 2. Osnovni deskriptivni pokazatelji (medijana i interval) stepena depresivnosti, anksioznosti i stresa, po godinama studija (I-VI)

	I	II	III	IV	V	VI
Depresija	7 (0-20)	3 (0-17)	3 (0-9)	1 (0-13)	7 (3-8)	3 (0-16)
Anksioznost	7 (0-7)	8 (0-14)	2 (1-10)	2 (0-8)	5 (0-12)	3 (0-11)
Stres	10 (3-18)	9 (1-20)	7 (2-16)	6 (2-15)	9 (7-16)	7 (1-19)

Rezultati poređenja stepena DAS u odnosu na godinu studija prikazani su na Slici 2. Studenti I godine generalno su pokazali najveći stepen DAS u odnosu na studente ostalih godina studija, dok je najmanji stepen zabeležen kod studenata IV godine. Jednofaktorskom analizom varijanse ustanovljene su razlike između studenata različitih godina studija u stepenu depresivnosti ($F_{5,88} = 2,515, p = 0,036$) i anksioznosti ($F_{5,88} = 3,594, p = 0,005$), ali ne i u simptomima stresa ($F_{5,88} = 1,55, p = 0,183$). Kod studenata I godine simptomi depresije prisutni su značajno više nego kod studenata III (prosečna razlika [PR] = 4,33, $p = 0,042$) i studenata IV godine studija (PR = 4,66, $p = 0,009$), dok su kod studenata II godine simptomi depresije prisutni više nego kod studenata IV godine (PR = 3,61, $p = 0,009$). Studenti I godine pokazali su i veći stepen anksioznosti u odnosu na kolege sa IV (PR = 4,47, $p = 0,004$) i VI godine studija (PR = 3,42, $p = 0,036$). Veći stepen anksioznosti zabeležen je i kod studenata II godine u odnosu na studente IV (PR = 4,17, $p = 0,001$) i VI godine studija (PR = 3,12, $p = 0,018$).

Slika 2. Osnovni deskriptivni pokazatelji ($SV \pm SD$) stepena depresivnosti, anksioznosti i stresa i rezultati poređenja u odnosu na godinu studija

Legenda: DASS-21 Skor – vrednost skora na skali depresivnosti anksioznosti i stresa, simbolom * označene su razlike između grupa značajne na nivou $p < 0,05$, a simbolom ** razlike značajne na nivou $p < 0,01$

Procenom nivoa FA na nivou uzorka dobijene su vrednosti koje spadaju u kategoriju umerene ukupne FA (Med [interval]) = 2613 [462-11931] MET-min/nedelja, pri čemu sedentarne aktivnosti, Med [interval] = 408 [75-780] MET-min/nedelja; laka FA, Med [interval] = 1386 [132-4851] MET-min/nedelja; umerena, Med [interval] = 240 [0-3600] MET-min/nedelja i intenzivna, Med [interval] = 0 [0-6720] MET-min/nedelja.

Slika 3 prikazuje osnovne deskriptivne pokazatelje (Med i interval) različitih tipova FA u poslednjih sedam dana, i rezultate poređenja u odnosu na pol. Može se zapaziti da su ispitanici muškog pola pokazali veće skorove na IPAQ skali u odnosu na svoje koleginice, ali su te razlike značajne samo u slučaju intenzivne fizičke aktivnosti ($\chi^2 = 7,69, p = 0,006$), ali ne i u ostalim vrstama aktivnosti, kao i na zbirnom IPAQ skor.

Slika 3. Osnovni deskriptivni pokazatelji (medijana i interval) različitih tipova fizičke aktivnosti (FA), ukupne FA i rezultati poređenja u odnosu na pol

Legenda: MET-min/nedelja – metabolički ekvivalent, M – muški pol, Ž – ženski pol. Pravougaoni dijagrami predstavljaju interkvartilni raspon, dok središnja linija predstavlja medijanu. Simbol ** označava razlike značajne na nivou $p < 0,01$

U Tabeli 3 prikazani su osnovni deskriptivni pokazatelji (Med i interval) različitih tipova FA (sedentarna, laka, umerena i intenzivna) i ukupne FA prema godinama studija. Uočava se da su studenti III, V i VI godine ostvarili veliku ukupnu energetske potrošnje u toku jedne nedelje (MET-min/nedelja ≥ 3000), za razliku od studenata I, II i IV godine, čije su vrednosti MET-min/nedelja iznosile < 3000 i spadaju u kategoriju umerene energetske potrošnje. Najveću energetske potrošnje ostvarili su studenti III i V godine (MET-min/nedelja = 3732), dok su najmanje bili aktivni studenti I godine (MET-min/nedelja = 1527). Poređenjem rezultata u nivou FA, u odnosu na godinu studija nisu zabeležene razlike između grupa (F vrednosti su se kretale u rasponu od $F_{5,88} = 0,416$ do $F_{5,88} = 1,676$).

Tabela 3. Osnovni deskriptivni pokazatelji (medijana i interval) različitih tipova fizičke aktivnosti (FA) kao i ukupne FA u poslednjih sedam dana, prikazani u metaboličkim ekvivalentima (MET-min/nedelja), po godinama studija (I-VI)

	I	II	III
Sedentarna	435 (75-600)	420 (150-780)	360 (300-600)
Laka	1056 (330-4851)	1386 (132-4752)	1386 (132-4752)
Umerena	240 (0-3360)	0 (0-1800)	0 (0-3360)
Intenzivna	0 (0-6480)	0 (0-1920)	480 (0-4320)
Ukupna	1527 (462-11931)	1911 (693-6432)	3732 (1386-10878)
	IV	V	VI
Sedentarna	420 (90-720)	360 (180-720)	480 (120-540)
Laka	1386 (297-4851)	2720 (495-3465)	1155 (198-4158)
Umerena	240 (0-3360)	480 (0-3360)	480 (0-3600)
Intenzivna	320 (0-3840)	960 (0-4320)	960 (0-6720)
Ukupna	2892 (462-9891)	3732 (933-10452)	3108 (594-11466)

Upoređujući rezultate između fizički aktivnih i fizički neaktivnih ispitanika u stepenu DAS, uočen je trend da su sve tri vrste simptoma više izražene kod fizički neaktivnih studenata muškog pola, kao i kod fizičkih neaktivnih studentkinja (sa izuzetkom depresije). Međutim, značajne razlike koje bi potvrdile ovakav nalaz nisu zabeležene.

Naposletku, ispitujući povezanost između FA i simptoma DAS na ukupnom uzorku, nije zabeležena značajna korelacija između zbirnog IPAQ skora i simptoma depresije ($r = -0,049$, $p = 0,649$), anksioznosti ($r = -0,130$, $p = 0,224$) i stresa ($r = -0,018$, $p = 0,865$). Analizirajući rezultate prema polu, zabeležena je mala pozitivna korelacija između sedentarnih aktivnosti i simptoma stresa kod žena ($r = 0,282$, $p = 0,030$), dok je kod muškaraca zabeležena mala negativna korelacija između umerene fizičke aktivnosti i simptoma depresije ($r = -0,279$, $p = 0,033$).

DISKUSIJA

Efikasnost fizičke aktivnosti (FA) u redukciji mentalnih oboljenja bliska je efektima tradicionalnog tretmana – psihoterapije (Paluska & Schwenk, 2000). Upražnjavanje FA nužno ne zahteva finansijska ulaganja, a pojedinac može i samostalno da je realizuje uz odgovarajuće instrukcije. Mnogi drugi tretmani koji ne podrazumevaju upotrebu lekova (npr. kognitivna bihevioralna terapija), nisu lako dostupni širokoj populaciji (iz ekonomskih ili nekih drugih razloga) što FA daje još jednu prednost. Takođe, opšte je poznato da potrebe lečenja mentalnih oboljenja nikada ne mogu biti u potpunosti zadovoljene samo od strane zdravstvenih radnika.

Prepoznavanje značaja povezanosti FA sa zdravljem dovelo je do velikog interesovanja za procenu ponašanja povezanih sa FA. Ranija istraživanja koja su se bavila nivoom FA u studentskoj populaciji pokazala su da on zavisi od vrste fakulteta (Malčić, 2018). S obzirom na to da studije medicine postavljaju visoke zahteve pred studente po pitanju obimnosti gradiva i svakodnevnih obaveza, to se može odraziti na simptome depresije, anksioznosti i stresa (DAS) i kod studenata koji nemaju psihijatrijsku dijagnozu.

Ovom studijom zabeležen je normalan stepen depresivnosti i anksioznosti i blag stepen stresa studenata Medicinskog fakulteta Univerziteta u Beogradu, pri čemu su kod pojedinih studenata zabeležene visoke vrednosti DAS varijabli koje ukazuju na jak osećaj. Dobijeni rezultati delimično su u skladu sa studijom sprovedenom među studentima brazilskih univerziteta, koja je zabeležila veće skorove na DASS-21 skali, pri čemu su pojedine grupe studenata prijavile visok stepen anksioznosti i stresa (Moutinho et al., 2017). Dalja analiza je pokazala da studenti, u zavisnosti od godine studija, generalno pokazuju normalan do umeren osećaj depresije, normalan do jak osećaj anksioznosti i normalan do umeren osećaj stresa. Navedeni rezultati delimično odstupaju od rezultata poslednjih istraživačkih studija u svetu koje ukazuju na to da su studenti rizična populacija za razvoj mentalnih oboljenja. U Sjedinjenim Američkim Državama, sprovedena je studija u kojoj je učestvovalo 67000 studenata i čiji nalazi upućuju na to da su problemi sa mentalnim zdravljem uobičajeni u populaciji studenata (Liu et al., 2019). Naime, rezultati su pokazali da je jedan od četiri studenta lečen od nekog poremećaja mentalnog zdravlja, da je petina studenata razmišljala o suicidu, da je 9% studenata pokušalo samoubistvo, a skoro 20% iskusilo je samopovređivanje. Jedan od razloga zbog kojih se nalazi ovog istraživanja u stepenu DAS razlikuju od rezultata prethodno navedenih studija verovatno leži u različitim kriterijumima za uključivanje ispitanika u istraživanje, koji je u slučaju ove studije podrazumevao odsustvo psihijatrijske dijagnoze i uzimanja psihofarmaka, što je verovatno rezultiralo relativno niskim skorovima na DASS-21 skali.

Rezultati ovog istraživanja ukazuju na to da studentkinje, u odnosu na studente, pokazuju veće skorove u stepenu DAS, sa značajno izraženijim simptomima anksioznosti. Ovakav nalaz u skladu je sa ranijim istraživanjima iz oblasti anksioznosti kod studenata medicine. U studiji koja se bavila analizom predispitne anksioznosti studenata medicine, pokazano je da su studentkinje anksioznije (Latas et al., 2010). Neki autori ovu razliku objašnjavaju time da je ženski pol spremniji priznati svoju anksioznost za razliku od muškaraca, a moguće je i da kod pripadnica ženskog pola postoji generalno izraženija anksioznost nego kod pripadnika muškog pola.

Iako rezultati ove studije ukazuju na to da ispitanici nisu pod velikim stresnim opterećenjem i izraženim depresivnim i anksioznim tegobama, ako se posmatraju pokazatelji DAS između ispitanika različitih godina studija uočavaju se razlike u vrednostima navedenih pokazatelja. To je najuočljivije kod studenata prve godine, što se može objasniti velikim promenama i novim izazovima sa kojima se brukoši suočavaju. U literaturi se, između ostalog, kao glavne smetnje mentalnom zdravlju studenata navode nepravilni obrasci spavanja, fluktuacije u bliskim odnosima i osećaj pritiska koji je uzrokovan akademskim zahtevima (Kadison & DiGeronimo, 2004), što je verovatno najviše izraženo kod studentata prve godine studija.

Procena nivoa FA ukazuje na to da je ukupna FA ispitivanih studenata spada u kategoriju umerene, pri čemu studenti najviše upražnjavaju laku FA (hodanje/šetanje), a najmanje intenzivnu. Pri tom, razlike između studenata različitih godina studija nisu zabeležene. Kad je u pitanju pol, studenti muškog pola značajno više su upražnjavali intenzivne FA u odnosu na studentkinje.

Nadalje, u studiji koja je analizirala povezanost sedentanog životnog stila i psiholoških poremećaja (Hamer et al., 2014) pokazano je da je umerena FA povezana sa manjim izgledom od depresije kod žena, dok je snažna do umerena FA bila povezana sa većim rizikom od depresije kod muškaraca. Slični nalazi proistekli su i iz studije Vanga i sar. (Wang et al., 2019) u kojoj je povezanost između depresije i FA u velikoj meri zavisila od intenziteta FA i pola. Niža učestalost, kraće trajanje i umerena FA bila je povezana sa manjim rizikom od depresije kod žena. S druge strane, kod muškaraca koji su primenjivali FA veće učestalost, trajanja i intenziteta, rizik od depresije bio je veći. S tim u vezi, nalazi ovog istraživanja delom su u skladu sa nalazima dobijenim u prethodnim studijama. Naime, procenjujući povezanost nivoa FA i stepena DAS, zabeležena je samo mala pozitivna korelacija između sedenja i simptoma stresa kod studentkinja, kao i mala negativna korelacija između umerene fizičke aktivnosti i simptoma depresije.

ZAKLJUČAK

Na osnovu nalaza ovog istraživanja može se zaključiti da je nivo fizičke aktivnosti, kao i stepen depresivnosti, anksioznosti i stresa ispitivanih studenata Medicinskog fakulteta Univerziteta u Beogradu na zadovoljavajućem nivou. Dobijeni rezultati delimično su u skladu sa prethodnim studijama, te bi za potvrdu ovakvih nalaza u nekim budućim istraživanjima trebalo obezbediti veći i homogeniji uzorak ispitanika, uz praćenje nivoa FA i pokazatelja DAS u dužem vremenskom periodu, a sve sa ciljem razvoja različitih sistema podrške u svrhu prevencije i smanjenja pojave mentalnih problema i teškoća.

LITERATURA

1. Ainsworth, B. E., Haskell, W. L., Whitt, M. C., Irwin, M. L., Swartz, A. M., Strath, S. J., O'Brien, W. L., Bassett, D. R., Schmitz, J. K. H., Emplainscourt, P. O., Jacobs, D. R., & Leon, A. S. (2000). Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise*, 32(9), 498-504.
2. Caspersen, C., Powell, K., Christenson G. (1985). Physical Activity, Exercise, and Physical Fitness: Definitions and Distinctions for Health-Related Research. *Pub Health Rep.* 100(2), 126–31.
3. Castrogiovanni, P., Trovato, F. M., Szychlinska, M. A., Nsir, H., Imbesi, R., & Musumeci, G. (2016). The importance of physical activity in osteoporosis. From the molecular pathways to the clinical evidence. *Histol Histopathol*, 31, 1183-1194.
4. Chin, S. H., Kahathuduwa, C. N., & Binks, M. (2016). Physical activity and obesity: what we know and what we need to know. *Obesity Reviews*, 17(12), 1226-1244.
5. Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine and science in sports and exercise*, 35(8), 1381–1395. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>.
6. De Moor, M. H. M., Beem, A. L., Stubbe, J. H., Boomsma, D. I., & De Geus, E. J. C. (2006). Regular exercise, anxiety, depression and personality: a population-based study. *Preventive Medicine*, 42(4), 273–279. <https://doi.org/10.1016/J.YPMED.2005.12.002>.
7. Diaz, K. M., & Shimbo, D. (2013). Physical activity and the prevention of hypertension. *Current hypertension reports*, 15(6), 659-668.
8. Dinas, P. C., Koutedakis, Y., & Flouris, A. D. (2011). Effects of exercise and physical activity on depression. *Irish journal of medical science*, 180(2), 319-325.
9. Erić L., (2015). *Enciklopedija straha*. Beograd: Službeni glasnik.

10. Fox, K. R., Boutcher, H. S., Guy, E., Biddle, F., & Biddle, J. H. S. (2000). The case for exercise in the promotion of mental health and psychological well-being. In: *Physical Activity and Psychological Well-Being*. London: Routledge; p. 1–10.
11. Goodwin, R. D. (2003). Association between physical activity and mental disorders among adults in the United States. *Preventive Medicine*, 36(6), 698–703. [https://doi.org/10.1016/S0091-7435\(03\)00042-2](https://doi.org/10.1016/S0091-7435(03)00042-2).
12. Hagströmer, M., Oja, P., & Sjöström, M. (2006). The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutrition*, 9(6), 755–762. <https://doi.org/10.1079/PHN2005898>.
13. Hamer, M., Coombs, N., & Stamatakis, E. (2014). Associations between objectively assessed and self-reported sedentary time with mental health in adults: an analysis of data from the Health Survey for England. *BMJ Open*, 4(3), e004580. <https://doi.org/10.1136/BMJOPEN-2013-004580>.
14. IPAQ Research Committee. (2005) *Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ). Short and long forms*. Stockholm, Karolinska Institute.
15. Kadison, R., & DiGeronimo, T. F. (2004). *College of the overwhelmed: The campus mental health crisis and what to do about it*. San Francisco, CA, US: Jossey-Bass.
16. Kohl, H. W. III, & Cook, H. D. (2013). Physical activity, fitness, and physical education: Effects on academic performance. In *Educating the student body: Taking physical activity and physical education to school*. (pp. 161–196). Washington DC: National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK201501/>.
17. Latas, M., Pantić, M., & Obradović, D. (2010). Analysis of test anxiety in medical students. *Medicinski Pregled*, 63(11–12), 863–866. <https://doi.org/10.2298/MPNS1012863L>.
18. LaMonte, M. J., Blair, S. N., & Church, T. S. (2005). Physical activity and diabetes prevention. *Journal of applied physiology*, 99(3), 1205–1213.
19. Liu, C. H., Stevens, C., Wong, S. H. M., Yasui, M., & Chen, J. A. (2019). The prevalence and predictors of mental health diagnoses and suicide among U.S. college students: Implications for addressing disparities in service use. *Depression and Anxiety*, 36(1), 8–17. <https://doi.org/10.1002/DA.22830>.
20. Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour research and therapy*, 33(3), 335–343. [https://doi.org/10.1016/0005-7967\(94\)00075-U](https://doi.org/10.1016/0005-7967(94)00075-U).
21. Malčić, B. (2018). Fizička aktivnost studenata Univerziteta u Novom Sadu – realnost i perspektive. *SPORT - Nauka i Praksa*, 8(1), 13–26.
22. Moutinho, I. L. D., De Castro Pecci Maddalena, N., Roland, R. K., Lucchetti, A. L. G., Tibiriçá, S. H. C., Da Silva Ezequiel, O., & Lucchetti, G. (2017). Depression, stress and anxiety in medical students: A cross-sectional comparison between students from different semesters. *Revista Da Associacao Medica Brasileira (1992)*, 63(1), 21–28. <https://doi.org/10.1590/1806-9282.63.01.21>
23. Oguma, Y., & Shinoda-Tagawa, T. (2004). Physical activity decreases cardiovascular disease risk in women: review and meta-analysis. *American journal of preventive medicine*, 26(5), 407–418.
24. Ostojić, S. M., Stojanović, M., Veljović, D., Stojanović, M. D., Međedović, B., Ahmetović Z. (2009). Fizička aktivnost i zdravlje: definicija problema, savremena zapažanja i preporuke. *TIMS Acta*, 3, 1–13.
25. Pallant, J. (Ed.) (2011). *SPSS: priručnik za preživljavanje*. Beograd: Mikro knjiga.
26. Paluska, S. A., & Schwenk, T. L. (2000). Physical activity and mental health: current concepts. *Sports Medicine (Auckland, N.Z.)*, 29(3), 167–180. <https://doi.org/10.2165/00007256-200029030-00003>.
27. Papathanasiou, G., Georgoudis, G., Papandreou, M., Spyropoulos, P., Georgakopoulos, D., Kalfakakou, V., & Evangelou, A. (2009). Reliability measures of the short International Physical Activity Questionnaire (IPAQ) in Greek young adults. *Hellenic journal of cardiology : HJC = Hellenike kardiologike epitheorese*, 50(4), 283–294.
28. Pedišić, Ž., Greblo, Z., Phongsavan, P., Milton, K., & Bauman, A.E. (2015). Are Total, Intensity- and Domain-Specific Physical Activity Levels 64 Associated with Life Satisfaction among University Students. *PLoS One*, 10(2).
29. Regehr, C., Glancy, D., & Pitts, A. (2013). Interventions to reduce stress in university students: a review and meta-analysis. *Journal of affective disorders*, 148(1), 1–11. <https://doi.org/10.1016/j.jad.2012.11.026>.
30. Shih, M., Hootman, J. M., Kruger, J., & Helmick, C. G. (2006). Physical activity in men and women with arthritis: National Health Interview Survey, 2002. *American journal of preventive medicine*, 30(5), 385–393.

31. Siscovick, D. S., LaPorte, R. E., & Newman, J. M. (1985). The disease-specific benefits and risks of physical activity and exercise. *Public health reports (Washington, D.C. : 1974)*, 100(2), 180–188.
32. Stojanović, M. D., Ostojić, S. M., & Milošević, Z. S. (2013). Fizička (ne)aktivnost - definicija, učestalost i ekonomski aspekti. *Teme*, 2, 857-866.
33. Stonerock, G. L., Hoffman, B. M., Smith, P. J., & Blumenthal, J. A. (2015). Exercise as Treatment for Anxiety: Systematic Review and Analysis. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 49(4), 542–556. <https://doi.org/10.1007/S12160-014-9685-9>.
34. Strickland, J. C., & Smith, M. A. (2014). The anxiolytic effects of resistance exercise. *Frontiers in Psychology*, 5. <https://doi.org/10.3389/FPSYG.2014.00753>.
35. Sylvia, L. G., Bernstein, E. E., Hubbard, J. L., Keating, L., & Anderson, E. J. (2014). Practical guide to measuring physical activity. *Journal of the Academy of Nutrition and Dietetics*, 114(2), 199–208. <https://doi.org/10.1016/j.jand.2013.09.018>.
36. The International Physical Activity Questionnaire. (2005). Available at: <http://www.ipaq.ki.se/>.
37. Wang, R., Bishwajit, G., Zhou, Y., Wu, X., Feng, D., Tang, S., Chen, Z., Shaw, I., Wu, T., Song, H., Fu, Q., & Feng, Z. (2019). Intensity, frequency, duration, and volume of physical activity and its association with risk of depression in middle- and older-aged Chinese: Evidence from the China Health and Retirement Longitudinal Study, 2015. *PloS One*, 14(8). <https://doi.org/10.1371/JOURNAL.PONE.0221430>.
38. WHO. (2010). Global recommendations on physical activity for health. *Geneva World Health Organ*, 60.
39. Wolin, K. Y., Yan, Y., Colditz, G. A., & Lee, I. M. (2009). Physical activity and colon cancer prevention: a meta-analysis. *British journal of cancer*, 100(4), 611-616.

Original scientific paper

INFLUENCE OF PHYSICAL ACTIVITY ON DEGREE OF DEPRESSION, ANXIETY AND STRESS IN STUDENTS OF UNIVERSITY OF BELGRADE - FACULTY OF MEDICINE¹

UDC: 613.71-057.875(497.11)

159.944.4.072-057.875

616.89-008.441-057.875

616.89-008.454-057.875

DOI: 10.5937/snp2102057L

Uroš Lazarević²

Medical Faculty, University of Belgrade, Serbia

Dragana Drljačić³

College of Sports and Health, Belgrade, Serbia

Miloš Bojović

College of Sports and Health, Belgrade, Serbia

Srdan Milosavljević

College of Sports and Health, Belgrade, Serbia

Abstract: Mental health problems have become increasingly present among students. Therefore, the aim of this study was to examine the influence of physical activity (PA) on the manifestation of the symptoms of depression, anxiety and stress (DAS) in medical students. The sample of examinees included 89 students (59 female [F] and 30 male [M] students), from the first to the sixth year of studies at the Faculty of Medicine, University of Belgrade, aged 22 ± 3.3 . The measuring instruments were a short version of the Depression, Anxiety and Stress Scale (DASS-21) and a short form of the International Physical Activity Questionnaire (IPAQ). The obtained results indicate that average scores fall into the category of normal degree of depression and anxiety, and a moderate degree of stress, whereby F show higher scores on the anxiety scale ($\chi^2 = 8.36$, $p = 0.004$). Different scores were noted among students of different years of study on the depression scale ($F = 2.515$, $p = 0.036$), as well the anxiety scale ($F = 3.594$, $p = 0.005$), but not on the stress scale ($F = 1.55$, $p = 0.183$). Total PA was assessed as moderate (median [interval] = 2613 [462-11931] MET-min/week), whereby M significantly more frequently engage in high intensity activities ($\chi^2 = 7.69$, $p = 0.006$). The research also recorded a small positive correlation between sedentary activities and stress symptoms in F ($r = 0.282$, $p = 0.030$), while small negative correlation was noted in M between moderate PA and depression symptoms ($r = -0.279$, $p = 0.033$). The research results indicate that total PA and the degree of DAS in examined students is at a satisfactory level.

Keywords: *IPAQ, DASS-21, metabolic equivalent, neurotic disorders, mood*

¹ Paper received: 14 December 2021, edited: 24 December 2021, accepted for publication: 24 December 2021.

² Uroš Lazarević attends specialist studies at the Faculty of Medicine, University of Belgrade.

³ ✉ dragana.drljajic@vss.edu.rs

INTRODUCTION

Physical activity is any skeletal muscle movement that requires energy consumption (WHO, 2010), i.e. body movement produced by the musculoskeletal system that results in energy consumption above the threshold that the body consumes at rest (Caspersen et al., 1985). It includes various activities on a daily basis (from housework to recreational and physical exercise) (Pedišić, 2004; Stojanović et al., 2013). In the last few decades, numerous scientific studies have examined the relationship between physical activity and health, as well as the effects of physical exercise on certain organ systems (Kohl et al., 2013).

The extent of the impact of physical activity on health has turned out to be large enough that researchers, even with relatively rough measurement techniques, have been able to consistently document the strong association of physical exercise with health benefits (Siscovick et al., 1985; Sylvia et al., 2014). It has been proved that there is an evident link between regular physical activity and a reduction in the incidence of certain diseases, such as coronary heart disease (Oguma & Shinoda-Tagawa, 2004), diabetes (LaMonte et al., 2005), hypertension (Diaz & Shimbo, 2013), arthritis (Shih et al., 2006), osteoporosis (Castrogiovanni et al., 2016), colon cancer (Wolin et al., 2009), depression (Dinas et al., 2011), obesity (Chin et al., 2016) etc. There is also evidence that regular physical activity leads to the improvement of certain bodily processes of physically active people, such as carbohydrate and fat metabolism, hemocoagulation processes, immune system function, improving cognitive function and mental health in general (Kohl et al., 2013).

Considering that there is a clear positive impact of physical activity in the prevention and reduction of physical illness, we are witnessing increased interest in its contribution and in preserving and alleviating the problems of mental illnesses. The positive effects of physical activity on mental health have been shown in a large number of studies (Erić, 2015; Goodwin, 2003). In one of them, which included over 8,000 respondents, findings were obtained that indicate that regular physical activity significantly reduces the risk of depression and anxiety disorders (De Moor et al., 2006). Also, intervention studies show that even one-time physical activity leads to a reduction in symptoms of depression, anxiety and stress immediately after the end of physical activity (Stonerock et al., 2015; Strickland & Smith, 2014).

The World Health Organization estimates that depression, anxiety and stress are the main causes of reduced quality of life and shortened life expectancy today (Ostojić et al., 2009). Due to the increasing incidence of serious mental illnesses, there is a fear of an increase in the number of people suffering from chronic or recurrent, mild to moderate symptoms of these illnesses. We are already witnessing an increased occurrence of mental problems in the general public, which can be characterized as mild depression, low self-esteem, high stress and anxiety, and poor coping with the difficulties and problems of everyday life. The reduction of the presence of physical activity in institutions and the promotion of physical activity as a culture of living also contribute to these phenomena (Fox et al., 2000).

Mental health problems are becoming more common among students as well. Research has shown that almost half of university students show moderate levels of stress-related mental health problems, including anxiety and depression (Regehr et al., 2013). After becoming academic citizens, students face an increased scope of academic obligations and responsibilities, and thus a number of psychological and psychosocial challenges associated with their personal development and independence. In this regard, the studies of medicine are among the most challenging in the world, including in our country. The university life of medical students during medical training implies great commitment and responsibility in terms of performing academic tasks and caring for patients and their families. A large number of hours spent in lectures and clinical practice, an environment that is often not ideal for learning, lack of sleep as well as factors that interfere with daily personal life, are common during this period (Moutinho et al., 2017). All of these factors affect the mental and emotional status of students (Latas et al., 2010) and, in the absence of activities that improve quality of life, can contribute to the development of physical, mental, and emotional problems.

Research has shown that medical students belong to a population in which the phenomena of pre-examination anxiety are particularly prevalent (Latas et al., 2010). There are many publications on the topic of pre-examination anxiety, but the literature on the impact of physical activity on the symptoms of depression, anxiety and stress in a healthy student population, according to the authors of this paper, is scarce. For this reason, the aim of this study was to examine the impact of physical activity on the manifestation of symptoms of depression, anxiety and stress of students of medicine.

METHOD

Examinee sample

The research included 89 male and female students (F= 59 and M = 30) attending the first to the sixth year of studies of the Faculty of Medicine, University of Belgrade (1st year, N = 10; 2nd year, N = 23; 3rd year, N = 9; 4th year, N = 23; 5th year, N = 7 and 6th year, N = 17), of average age 22 ± 3.3 years, covering ages from 19 to 35. Relevant data concerning socio-demographic variables (age, sex, year of study) and clinical characteristics of the examinees were collected using a semi-structured questionnaire, and then used as criteria for the inclusion of examinees into the research. The presence of any kind of psychiatric diagnosis and/or the use of psychopharmaceuticals represented the main criteria for elimination from further research.

The research was conducted in September 2019, as a cross-sectional study, and it was approved by the Ethical Committee of the Faculty of Medicine in Belgrade.

Measuring instruments

To measure the degree of depression, anxiety, and stress, a short version of the Depression, Anxiety, and Stress Scale (DASS-21) consisting of 21 items was used (Lovibond & Lovibond, 1995). DASS-21 includes three subscales, with seven items each (depression scale, e.g. *I felt life was meaningless*; anxiety scale, e.g. *I felt close to panic*; stress scale, e.g. *I was annoyed when something interrupted me in what I do*). The Depression Scale assesses dysphoria, hopelessness, devaluation of life, self-regulation, and lack of interest / involvement in events, anhedonia and inertia. The anxiety scale assesses autonomic nervous system reactions, skeletal muscle reactions, situational anxiety, and the subjective experience of anxiety affect. The stress scale is sensitive to levels of chronic nonspecific arousal. It assesses the difficulties in relaxation, nervous excitement and easily triggered anxiety and impatience.

During the measurements, the examinees had the task to assess the degree to which they experienced each of these conditions in the last seven days on a four-point Likert-type scale (from 0 - not at all, to 3 - mostly or almost always) (Lovibond & Lovibond, 1995). Depression, anxiety, and stress (DAS) scores were calculated by summing the scores for the relevant items, with scores for each subscale ranging from 0 to 21. The criteria for determining DAS levels are shown in Table 1.

Table 1. Assessment criteria for levels of depression, anxiety and stress (Lovibond & Lovibond, 1995)

Feeling	depression	anxiety	stress
normal	0-4	0-3	0-7
mild	5-6	4-5	8-9
moderate	7-10	6-7	10-12
strong	11-13	8-9	13-16
very strong	≥ 14	≥ 10	≥ 17

Legend: Columns show score intervals on the Depression, Anxiety and Stress Scale (DASS-21)

A short form of the International Physical Activity Questionnaire (IPAQ) was used to examine the level of physical activity (PA) (IPAQ Research Committee, 2005), which measures the frequency, duration and intensity of PA in the last seven days. This form of the IPAQ questionnaire is intended for health-related PA assessment in the population aged 15 to 69 and assesses PA levels through three types of activities: light PA, moderate and intense PA, and time spent in sedentary activities (sitting / lying down) (Hagströmer et al., 2006; Papathanasiou et al., 2009). The level of PA was estimated through the total energy consumption expressed in Metabolic Equivalents of Task. One metabolic equivalent (MET) represents the basal level of oxygen consumption and caloric expenditure associated with it (about 3.5 ml O₂ / kg / min or 1 Kcal / kg / h) and corresponds to the level of resting metabolism (Ainsworth et al., 2000). The calculation of MET was done according to the Guidelines for Data Processing and Analysis of the International PA Questionnaire (IPAQ Research Committee, 2005). MET-min / week values < 600 were considered low energy consumption and corresponded to the category of light PA. Values from 601 to 3000 MET-min / week

corresponded to moderate PA, and values > 3000 MET-min / week to intensive PA (Craig et al., 2003). Furthermore, based on the cumulative IPAQ score, the examinees were classified into the group of active subjects (MET-min / week ≥ 3000) and the group of inactive subjects (MET-min / week < 3000).

Statistical analysis

Basic descriptive indicators (frequencies and percentages for attribute features, i.e. $SV \pm SD$, minimum and maximum for numerical features) were calculated for all variables. The normality of the distribution of numerical features was examined using the Shapiro-Wilk test, which showed the absence of a normal distribution for all monitored variables. In line with the results obtained for the purpose of examining differences in the degree of DAS in relation to sex, differences in types of PA in relation to sex, as well as differences in the degree of DAS in relation to the level of PA between respondents of the same sex and physically active and physically inactive students, the Kruskal-Wallis test was applied. Since the results of nonparametric statistics did not differ from the results obtained by parametric tests, to examine differences in the degree of DAS between respondents of different years of study, as well as differences in PA between respondents of different years of study, the paper presents results obtained by one-factor analysis of variance. In case of significance of the main factor, subsequent Fischer LSD tests were performed. Spearman's rank correlation coefficient was applied to examine the relationship between PA levels and DAS levels, by testing the correlation between individual IPAQ instrument subscales and DASS-21 subscales. According to Cohen's criteria, values from 0.10 to 0.29 were considered small, from 0.30 to 0.49 medium, and values from 0.50 to 1.00 were considered to indicate high correlation (Pallant, 2011). Statistical tests were calculated using SPSS 20.0 software (SPSS Inc, Chicago, IL), where the threshold of significance of statistical analyses was at the level of confidence $p = 0.05$.

RESULTS

Assessing the degree of DAS on the total sample yielded results that fall into the category of normal feeling on the scale of depression (median and interval, Med [interval] = 3 [0-20]) and anxiety (Med [interval] = 3 [0-17]) and a mild feeling on the stress scale (Med [interval] = 8 [1-20]). However, high maximum values were also recorded, for all three scales, which fall into the category of very strong feeling.

Picture 1 shows the basic descriptive indicators (Med and interval) of the degree of DAS and the results of comparisons in relation to gender, where it can be observed that all three types of symptoms are more pronounced in female subjects. However, significant differences were observed only in the symptoms of anxiety ($\chi^2 = 8.36$, $p = 0.004$), but not in the symptoms of depression ($\chi^2 = 0.94$, $p = 0.333$) and stress ($\chi^2 = 2.54$, $p = 0.111$).

Picture 1. Basic descriptive indicators (median and interval) of the degree of depression, anxiety and stress and the results of comparisons in relation to sex

Legend: Rectangular diagrams represent the interquartile range, while the centre line represents the median. The symbol ** indicates differences between the sexes (M-male and F-female) significant at the level of $p < 0.01$

Table 2 shows the basic descriptive indicators (Med and interval) of DAS degree, by years of study. On the scale of depression, the largest number of results is within the normal feeling of depression (score 0-4), except for the 1st and the 5th year, where it is within the moderate feeling of depression (7-10). Anxiety scores range from normal feeling (0-3) for the 3rd, 4th and 6th year, mild for the 5th, moderate (6-7) for the 1st and strong feeling of anxiety for the 2nd year of study. Stress scores range from normal feelings for the 3rd, 4th and 6th year (score 0-7), mild for the 2nd and 5th (score 8-9), to moderate feelings of stress (score 10-11) for the 1st year of study.

Table 2. Basic descriptive indicators (median and interval) of the degree of depression, anxiety and stress, by years of study (I-VI)

	I	II	III	IV	V	VI
Depression	7 (0-20)	3 (0-17)	3 (0-9)	1 (0-13)	7 (3-8)	3 (0-16)
Anxiety	7 (0-7)	8 (0-14)	2 (1-10)	2 (0-8)	5 (0-12)	3 (0-11)
Stress	10 (3-18)	9 (1-20)	7 (2-16)	6 (2-15)	9 (7-16)	7 (1-19)

The results of comparing the degree of DAS in relation to the year of study are shown in Picture 2. First-year students generally showed the highest degree of DAS in relation to students of other years of study, while the lowest degree was recorded in fourth-year students. One-factor analysis of variance revealed differences between students of different years of study in the degree of depression ($F_{5,88} = 2.515$, $p = 0.036$) and anxiety ($F_{5,88} = 3.594$, $p = 0.005$), but not in symptoms of stress ($F_{5,88} = 1.55$, $p = 0.183$). Depression symptoms are significantly more present in first-year students than in third-year students (average difference [AD] = 4.33, $p = 0.042$) and in fourth-year students (AD = 4.66, $p = 0.009$), while in students of the 2nd year, the symptoms of depression were more present than in the fourth year students (AD = 3.61, $p = 0.009$). First-year students also showed a higher degree of anxiety compared to colleagues attending the 4th (AD = 4.47, $p = 0.004$) and 6th year of study (AD = 3.42, $p = 0.036$). A higher degree of anxiety was also recorded among second-year students compared to fourth-year students (AD = 4.17, $p = 0.001$) and sixth-year students (AD = 3.12, $p = 0.018$).

Picture 2. Basic descriptive indicators ($SV \pm SD$) of the degree of depression, anxiety and stress and comparison results in relation to the year of study

Legend: DASS-21 Score - the value of the score on the scale of depression, anxiety and stress, the symbol * indicates differences between groups significant at the level of $p < 0.05$, and the symbol ** differences significant at the level of $p < 0.01$

Estimation of PA levels at the sample level yielded values that fall into the category of moderate total PA (Med [interval]) = 2613 ([462-11931] MET-min / week), with sedentary activities, Med [interval] = 408 [75- 780]

MET-min / week; light PA, Med [interval]) = 1386 [132-4851] MET-min / week; moderate, Med [interval]) = 240 [0-3600] MET-min / week and intense, Med [interval]) = 0 [0-6720] MET-min / week.

Picture 3 shows the basic descriptive indicators (Med and interval) of different types of PA in the last seven days, and the results of comparisons in relation to gender. It can be noticed that male respondents scored higher on the IPAQ scale compared to their female counterparts, but these differences are significant only in the case of intense physical activity ($x_2 = 7.69$, $p = 0.006$), but not in other types of activities, as well as on the cumulative IPAQ score.

Picture 3. Basic descriptive indicators (median and interval) of different types of physical activity (PA), total PA and results of comparisons in relation to gender

Legend: MET-min / week - metabolic equivalent, M - male, F - female. Rectangular diagrams represent the interquartile range, while the centre line represents the median. The symbol ** indicates differences that are significant at the level of $p < 0.01$

Table 3 shows the basic descriptive indicators (Med and interval) of different types of PA (sedentary, light, moderate and intensive) and total PA according to years of study. It can be noticed that students of the 3rd, 5th and 6th year of study achieved a large total energy consumption during one week ($\text{MET-min / week} \geq 3000$), in contrast to students of the 1st, 2nd and 4th year, whose values of MET-min / week were < 3000 and belong to the category of moderate energy consumption. The highest energy consumption was achieved by students of the 3rd and 5th year ($\text{MET-min / week} = 3732$), while the least active were first-year students ($\text{MET-min / week} = 1527$). Comparing the results at the PA level, in relation to the year of study, no differences were observed between the groups (F values ranged from $F_{5.88} = 0.416$ to $F_{5.88} = 1.676$).

Table 3. Basic descriptive indicators (median and interval) of different types of physical activity (PA) as well as total PA in the last seven days, presented in metabolic equivalents (MET-min / week), by years of study (1st-6th)

	1st	2nd	3rd
Sedentary	435 (75-600)	420 (150-780)	360 (300-600)
Light	1056 (330-4851)	1386 (132-4752)	1386 (132-4752)
Moderate	240 (0-3360)	0 (0-1800)	0 (0-3360)
Intense	0 (0-6480)	0 (0-1920)	480 (0-4320)
Total	1527 (462-11931)	1911 (693-6432)	3732 (1386-10878)
	4th	5th	6th
Sedentary	420 (90-720)	360 (180-720)	480 (120-540)
Light	1386 (297-4851)	2720 (495-3465)	1155 (198-4158)
Moderate	240 (0-3360)	480 (0-3360)	480 (0-3600)
Intense	320 (0-3840)	960 (0-4320)	960 (0-6720)
Total	2892 (462-9891)	3732 (933-10452)	3108 (594-11466)

Comparing the results between physically active and physically inactive subjects in the DAS degree, a trend was observed that all three types of symptoms were more pronounced in physically inactive male students, as well as in physically inactive female students (with the exception of depression). However, significant differences that would confirm this finding were not noted.

Finally, examining the association between PA and DAS symptoms in the total sample, no significant correlation was observed between the cumulative IPAQ score and symptoms of depression ($r = -0.049$, $p = 0.649$), anxiety ($r = -0.130$, $p = 0.224$) and stress ($r = -0.018$, $p = 0.865$). Analysing the results by gender, there was a small positive correlation between sedentary activities and stress symptoms in women ($r = 0.282$, $p = 0.030$), while in men there was a small negative correlation between moderate physical activity and symptoms of depression ($r = -0.279$, $p = 0.033$).

DISCUSSION

The effectiveness of physical activity (PA) in the reduction of mental illness is close to the effects of traditional treatment - psychotherapy (Paluska & Schwenk, 2000). Exercising PA does not necessarily require financial investments, and an individual can practice it independently with appropriate instructions. Many other non-drug treatments (e.g., cognitive behavioural therapy) are not readily available to the general population (for economic or other reasons), which is another advantage of PA. It is also common knowledge that treatment needs for mental illness can never be fully met by health professionals alone.

Recognizing the importance of the connection of PA with health has led to great interest in assessing PA-related behaviours. Earlier research that studied the level of PA in the student population showed that it depends on the type of faculty (Malčić, 2018). Given that medical studies place high demands on students in terms of volume of material and daily obligations, this can be reflected in the symptoms of depression, anxiety and stress (DAS) even in students who do not have a psychiatric diagnosis.

This study showed a normal degree of depression and anxiety and a mild degree of stress in students of the Faculty of Medicine, University of Belgrade, with some students experiencing high values of DAS variables that indicate a strong feeling. The results obtained are partly in line with a study conducted among students of Brazilian universities, which recorded higher scores on the DASS-21 scale, with some groups of students reporting high levels of anxiety and stress (Moutinho et al., 2017). Further analysis showed that students, depending on the year of study, generally show a normal to moderate feeling of depression, a normal to strong feeling of anxiety and a normal to moderate feeling of stress. These results deviate in part from the results of recent research studies in the world, which indicate that students are a risk population for the development of mental illness. In the United States, a study was conducted on 67,000 students, and its findings indicate that mental health problems are common in the student population (Liu et al., 2019). Namely, the results showed that one in four students was treated for a mental health disorder, that a fifth of students were thinking about suicide, that 9% of students had attempted suicide, and almost 20% had experienced self-harm. One of the reasons why the findings of this DAS study differ from the results of the previous studies probably lies in the different criteria for inclusion of respondents in the study, which in the case of this study implied the absence of psychiatric diagnosis and taking psychopharmaceuticals, which probably resulted in relatively low scores on the DASS-21 scale.

The results of this research indicate that female students, compared to male students, show higher scores in the degree of DAS, with significantly more pronounced symptoms of anxiety. This finding is in line with previous research in the field of anxiety in students of medicine. In a study that analysed the pre-examination anxiety of medical students, it was shown that female students are more anxious (Latas et al., 2010). Some authors explain this difference by saying that female respondents are more willing to admit their anxiety than males, and it is possible that females are generally more anxious than males.

Although the results of this study indicate that the respondents are not under high stress and severe depressive and anxiety problems, if the DAS indicators are observed between the respondents of different years of study, there are differences in the values of these indicators. This is most noticeable with first-year students, which can be explained by the big changes and new challenges that they face. In the literature, among other things, irregular sleep patterns, fluctuations in close relationships and feelings of pressure caused by academic demands are cited as the main obstacles to students' mental health (Kadison & DiGeronimo, 2004), which is probably most pronounced in first-year students.

The assessment of the level of PA indicates that the total PA of the examined students belongs to the category of moderate, with students practicing light PA (walking) the most, and intense the least. At the same time, differences between students of different years of study were not noticed. When it comes to differences between sexes, male students practiced intensive PAs significantly more than female students.

Furthermore, a study that analysed the association between sedentary lifestyle and psychological disorders (Hamer et al., 2014) showed that moderate PA was associated with a lower chance of depression in women, while intense to moderate PA was associated with a higher risk of depression in men. Similar findings emerged from the study by Wang et al. (2019) in which the association between depression and PA largely depended on the intensity of PA and respondents' sex. Lower incidence, shorter duration, and moderate PA were associated with a lower risk of depression in women. On the other hand, in men who used PA with a higher frequency, duration and intensity, the risk of depression was higher. In this regard, the findings of this study are partly consistent with the findings of previous studies. Namely, assessing the relationship between PA levels and DAS levels, only a small positive correlation was found between sedentary lifestyle and stress symptoms in female students, as well as a small negative correlation between moderate physical activity and depression symptoms.

CONCLUSION

Based on the findings of this study, it can be concluded that the level of physical activity, as well as the degree of depression, anxiety and stress of the examined students of the Faculty of Medicine, University of Belgrade is at a satisfactory level. The obtained results are partly in line with previous studies, and to confirm such findings in some future research should provide a larger and more homogeneous sample of respondents, with monitoring of PA levels and DAS indicators over time, all with the aim of developing different support systems for prevention and reduction of mental problems and difficulties.

REFERENCES

1. Ainsworth, B. E., Haskell, W. L., Whitt, M. C., Irwin, M. L., Swartz, A. M., Strath, S. J., O'Brien, W. L., Bassett, D. R., Schmitz, J. K. H., Emplaincourt, P. O., Jacobs, D. R., & Leon, A. S. (2000). Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise*, 32(9), 498-504.
2. Caspersen, C., Powell, K., Christenson G. (1985). Physical Activity, Exercise, and Physical Fitness: Definitions and Distinctions for Health-Related Research. *Pub Health Rep.* 100(2), 126–31.
3. Castrogiovanni, P., Trovato, F. M., Szychlińska, M. A., Nsir, H., Imbesi, R., & Musumeci, G. (2016). The importance of physical activity in osteoporosis. From the molecular pathways to the clinical evidence. *Histol Histopathol*, 31, 1183-1194.
4. Chin, S. H., Kahathuduwa, C. N., & Binks, M. (2016). Physical activity and obesity: what we know and what we need to know. *Obesity Reviews*, 17(12), 1226-1244.
5. Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine and science in sports and exercise*, 35(8), 1381–1395. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>.
6. De Moor, M. H. M., Beem, A. L., Stubbe, J. H., Boomsma, D. I., & De Geus, E. J. C. (2006). Regular exercise, anxiety, depression and personality: a population-based study. *Preventive Medicine*, 42(4), 273–279. <https://doi.org/10.1016/J.YPMED.2005.12.002>.
7. Diaz, K. M., & Shimbo, D. (2013). Physical activity and the prevention of hypertension. *Current hypertension reports*, 15(6), 659-668.
8. Dinas, P. C., Koutedakis, Y., & Flouris, A. D. (2011). Effects of exercise and physical activity on depression. *Irish journal of medical science*, 180(2), 319-325.
9. Erić L., (2015). *Enciklopedija straha*. Beograd: Službeni glasnik.

10. Fox, K. R., Boutcher, H. S., Guy, E., Biddle, F., & Biddle, J. H. S. (2000). The case for exercise in the promotion of mental health and psychological well-being. In: *Physical Activity and Psychological Well-Being*. London: Routledge; p. 1–10.
11. Goodwin, R. D. (2003). Association between physical activity and mental disorders among adults in the United States. *Preventive Medicine*, 36(6), 698–703. [https://doi.org/10.1016/S0091-7435\(03\)00042-2](https://doi.org/10.1016/S0091-7435(03)00042-2).
12. Hagströmer, M., Oja, P., & Sjöström, M. (2006). The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutrition*, 9(6), 755–762. <https://doi.org/10.1079/PHN2005898>.
13. Hamer, M., Coombs, N., & Stamatakis, E. (2014). Associations between objectively assessed and self-reported sedentary time with mental health in adults: an analysis of data from the Health Survey for England. *BMJ Open*, 4(3), e004580. <https://doi.org/10.1136/BMJOPEN-2013-004580>.
14. IPAQ Research Committee. (2005) *Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ). Short and long forms*. Stockholm, Karolinska Institute.
15. Kadison, R., & DiGeronimo, T. F. (2004). *College of the overwhelmed: The campus mental health crisis and what to do about it*. San Francisco, CA, US: Jossey-Bass.
16. Kohl, H. W. III, & Cook, H. D. (2013). Physical activity, fitness, and physical education: Effects on academic performance. In *Educating the student body: Taking physical activity and physical education to school*. (pp. 161–196). Washington DC: National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK201501/>.
17. Latas, M., Pantić, M., & Obradović, D. (2010). Analysis of test anxiety in medical students. *Medicinski Pregled*, 63(11–12), 863–866. <https://doi.org/10.2298/MPNS1012863L>.
18. LaMonte, M. J., Blair, S. N., & Church, T. S. (2005). Physical activity and diabetes prevention. *Journal of applied physiology*, 99(3), 1205–1213.
19. Liu, C. H., Stevens, C., Wong, S. H. M., Yasui, M., & Chen, J. A. (2019). The prevalence and predictors of mental health diagnoses and suicide among U.S. college students: Implications for addressing disparities in service use. *Depression and Anxiety*, 36(1), 8–17. <https://doi.org/10.1002/DA.22830>.
20. Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour research and therapy*, 33(3), 335–343. [https://doi.org/10.1016/0005-7967\(94\)00075-U](https://doi.org/10.1016/0005-7967(94)00075-U).
21. Malčić, B. (2018). Fizička aktivnost studenata Univerziteta u Novom Sadu – realnost i perspektive. *SPORT - Nauka i Praksa*, 8(1), 13–26.
22. Moutinho, I. L. D., De Castro Pecci Maddalena, N., Roland, R. K., Lucchetti, A. L. G., Tibiriçá, S. H. C., Da Silva Ezequiel, O., & Lucchetti, G. (2017). Depression, stress and anxiety in medical students: A cross-sectional comparison between students from different semesters. *Revista Da Associacao Medica Brasileira (1992)*, 63(1), 21–28. <https://doi.org/10.1590/1806-9282.63.01.21>
23. Oguma, Y., & Shinoda-Tagawa, T. (2004). Physical activity decreases cardiovascular disease risk in women: review and meta-analysis. *American journal of preventive medicine*, 26(5), 407–418.
24. Ostojić, S. M., Stojanović, M., Veljović, D., Stojanović, M. D., Međedović, B., Ahmetović Z. (2009). Fizička aktivnost i zdravlje: definicija problema, savremena zapažanja i preporuke. *TIMS Acta*, 3, 1–13.
25. Pallant, J. (Ed.) (2011). *SPSS: priručnik za preživljavanje*. Beograd: Mikro knjiga.
26. Paluska, S. A., & Schwenk, T. L. (2000). Physical activity and mental health: current concepts. *Sports Medicine (Auckland, N.Z.)*, 29(3), 167–180. <https://doi.org/10.2165/00007256-200029030-00003>.
27. Papathanasiou, G., Georgoudis, G., Papandreou, M., Spyropoulos, P., Georgakopoulos, D., Kalfakakou, V., & Evangelou, A. (2009). Reliability measures of the short International Physical Activity Questionnaire (IPAQ) in Greek young adults. *Hellenic journal of cardiology : HJC = Hellenike kardiologike epitheorese*, 50(4), 283–294.
28. Pedišić, Ž., Greblo, Z., Phongsavan, P., Milton, K., & Bauman, A.E. (2015). Are Total, Intensity- and Domain-Specific Physical Activity Levels 64 Associated with Life Satisfaction among University Students. *PLoS One*, 10(2).
29. Regehr, C., Glancy, D., & Pitts, A. (2013). Interventions to reduce stress in university students: a review and meta-analysis. *Journal of affective disorders*, 148(1), 1–11. <https://doi.org/10.1016/j.jad.2012.11.026>.
30. Shih, M., Hootman, J. M., Kruger, J., & Helmick, C. G. (2006). Physical activity in men and women with arthritis: National Health Interview Survey, 2002. *American journal of preventive medicine*, 30(5), 385–393.

31. Siscovick, D. S., LaPorte, R. E., & Newman, J. M. (1985). The disease-specific benefits and risks of physical activity and exercise. *Public health reports (Washington, D.C. : 1974)*, 100(2), 180–188.
32. Stojanović, M. D., Ostojić, S. M., & Milošević, Z. S. (2013). Fizička (ne)aktivnost - definicija, učestalost i ekonomski aspekti. *Teme*, 2, 857-866.
33. Stonerock, G. L., Hoffman, B. M., Smith, P. J., & Blumenthal, J. A. (2015). Exercise as Treatment for Anxiety: Systematic Review and Analysis. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 49(4), 542–556. <https://doi.org/10.1007/S12160-014-9685-9>.
34. Strickland, J. C., & Smith, M. A. (2014). The anxiolytic effects of resistance exercise. *Frontiers in Psychology*, 5. <https://doi.org/10.3389/FPSYG.2014.00753>.
35. Sylvia, L. G., Bernstein, E. E., Hubbard, J. L., Keating, L., & Anderson, E. J. (2014). Practical guide to measuring physical activity. *Journal of the Academy of Nutrition and Dietetics*, 114(2), 199–208. <https://doi.org/10.1016/j.jand.2013.09.018>.
36. The International Physical Activity Questionnaire. (2005). Available at: <http://www.ipaq.ki.se/>.
37. Wang, R., Bishwajit, G., Zhou, Y., Wu, X., Feng, D., Tang, S., Chen, Z., Shaw, I., Wu, T., Song, H., Fu, Q., & Feng, Z. (2019). Intensity, frequency, duration, and volume of physical activity and its association with risk of depression in middle- and older-aged Chinese: Evidence from the China Health and Retirement Longitudinal Study, 2015. *PloS One*, 14(8). <https://doi.org/10.1371/JOURNAL.PONE.0221430>.
38. WHO. (2010). Global recommendations on physical activity for health. *Geneva World Health Organ*, 60.
39. Wolin, K. Y., Yan, Y., Colditz, G. A., & Lee, I. M. (2009). Physical activity and colon cancer prevention: a meta-analysis. *British journal of cancer*, 100(4), 611-616.