Molecular Genetic Analysis of Cereal β -Amylase Genes Using Exon-Primed Intron-Crossing (EPIC) PCR

Olga Stratula · James Cockram · Ruslan Kalendar

received: 25 September 2014, accepted: 12 January 2015

published online: 6 February 2015

© 2014 IFVC

doi:10.5937/ratpov51-6808

Summary: The proteins encoded by cereal β -amylase genes *Bamy1* and *Bamy2* genes play an important role in seedling germination and in the brewing process. Here, we use exon-primed intron-crossing (EPIC) to analyse *Bamy1* and *Bamy2* genetic diversity among 38 accessions belonging to six *Poaceae* tribes. DNA sequence alignment of multiple *Poaceae* species β -amylase sequences allowed design of EPIC primers that simultaneously amplify *Bamy1* and *Bamy2* in all the cereal species investigated. The genetic variation observed in the samples investigated is analysed and discussed, and illustrates the effectiveness of this approach for intra- and interspecific analysis in plant species.

Keywords: β -amylase, barley, genetic diversity, intron-length polymorphism, *Poaceae*, rye, wheat

Abbreviation: Exon-Primed Intron-Crossing (EPIC)

Introduction

Due to the growing volume of data provided by genome sequencing projects, and the presence of regulatory regions within introns, the study of intronic polymorphism is emerging as an important aspect of genetic research in agricultural crops (Braglia et al. 2010). Compared to exonic regions, introns are more variable due to reduced selective pressure, i.e. the rate of mutation accumulation in these regions is comparatively high (Ludwig 2002, Morello & Breviario 2008, Yang et al. 2007). Introns often contain a variety of functional elements, including enhancers, silencers, regulators of alternative splicing, trans-splicing elements and other regulatory elements (some of which can be found in the expanded

introns). Furthermore, intron characteristics such as length variability, position within the gene, dependency on the length of the exons, can also modulate gene properties such as expression, transcription, splicing and microRNA lifetime (Braglia et al. 2010, Ludwig 2002, Rose & Beliakoff 2000). In plants, investigation into intron function has highlighted their role in gene expression, and specifically in tissue- or stage-specific expression (Morello & Breviario 2008). Analysis of plant intron sequences has been applied in numerous studies (Chetelat et al. 1995, Fridman et al. 2000, Holland et al. 2001, Hongtrakul V 1998), with a well-studied example in cereal species being the role of putative *cis*-elements in intron 1 of vernalization response (*Vrn*) genes. Deletions spanning this region are thought to result in vernalization non-responsiveness (Cockram et al. 2007), and underlie the creation of globally important spring-sown cereal cultivars.

The polymerase chain reaction (PCR) based technique, 'exon-primed intron-crossing' (EPIC-PCR) (Palumbi & Baker 1994) has gained favour in plant and animal studies, and relies on design of primers selected to anneal to highly-conserved regions of the exons. For example, it has been used to study conserved regions within eukaryotic 18S and 28S ribosomal genes and prokaryotic 16S and 23S ribosomal genes, for amplification of variable intergenic regions known as internal transcribed spacers (ITS), containing 5.8S ribosomal gene (Gardes 1993).

O. Stratula
Plant Breeding and Genetics Institute - National Center of Seed and Cultivar Investigation
Department of Genomics and Biotechnology, 3 Ovidiopskaya Road, 65036 Odessa, Ukraine

J. Cockram
National Institute of Agricultural Botany (NIAB), Huntingdon Road, Cambridge, CB3 0LE Cambridgeshire, United Kingdom

R. Kalendar*
LUKE Plant Genomics Laboratory, Institute of Biotechnology, University of Helsinki, Biocentre 3, P.O. Box 65, Viikinkaari 1, 00014 Helsinki, Finland
e-mail: ruslan.kalendar@helsinki.fi
Laboratory of Plant Genomics and Bioinformatics RSE "National Center for Biotechnology", 13/1 Sh. Valikhanov Str, 010000 Astana, Kazakhstan
e-mail: ruslan.kalendar@biocenter.kz

EPIC has a number of experimental advantages, including, (i) primers complementary to conserved regions within exons can be used across a wide taxonomic range, (ii) homologous amplified sequences can be easily defined by comparing their exonic and intronic regions depending on genetic distance between the taxons, and (iii) exonic and intronic fragments can help in the simultaneous studies of genetic variety at intraspecific and interspecific levels (Bierne et al. 2000, Ishikawa et al. 2007, Lessa 1992, Li et al. 2010). A further advantage of EPIC is that it avoids the need to develop DNA markers on a species-by-species basis, which can be costly and time consuming.

In cereal species (tribe: *Triticeae*), β -amylase hydrolyses starch by splitting α -1,4-glycosidic bonds, resulting in the formation of the high molecular weight dextrin and maltose - a disaccharide, that easily diffuses and can be used by the growing embryo (James et al. 2003). It is also an important component of the mash, a process used in brewing by which grain and water are combined and heated allowing enzymatic degradation of starch into sugars for fermentation (Priest 1986). Numerous species within the *Triticeae* can be used for brewing, including

barley, rye, triticale, wheat, oat, and millet (Briggs 1998). Only members of the *Triticeae* are reported to have two different forms of β -amylase (Mason-Gamer 2005), differing significantly in their patterns of gene expression: *Bamy1* is specific to the endosperm (endospermal β -amylase), while *Bamy2* is expressed in all tissues (ubiquitous β -amylase). These genes are paralogs, and both consist of 7 exons and 6 introns.

In this study, EPIC-PCR was used to investigate *Bamy1* and *Bamy2* intronic genetic variation within representative species belonging to the *Poaceae*. Due to their importance within cereal species, molecular genetic characterisation of cereal *Bamy1* and *Bamy2* will provide new information on the range of inter- and intra-specific genetic variation present. This data will help inform how *Triticeae* germplasm resources may be used as important sources of qualitative character enhancement, allowing enhancement of β -amylase activity in cereals.

Materials and Methods

Germplasm and DNA extraction

The 38 accessions investigated, belonging to six *Poaceae* tribes, are listed in Table 1.

Table 1. *Poaceae* germplasm used in this study

Plant material	Source	Cultivars/Lines
Accessions used for interspecific analyses		
38 species of the family <i>Poaceae</i>	Botanical Institute of the V.L. Koma-rova, RAS; Plant Breeding and Genetics Institute, Odessa	<i>Hordeum vulgare</i> (Djau Kabutak), <i>H. vulgare</i> (Pallidum 107), <i>H. vulgare</i> (Odesskiy 17), <i>H. vulgare</i> (Odesskiy 31), <i>H. spontaneum</i> , <i>H. murinum</i> , <i>H. marinum</i> , <i>H. brachyantherum</i> , <i>H. leporinum</i> , <i>Triticum durum</i> (AABB), <i>Aegilops speltoides</i> , <i>Agropyron cristatum</i> , <i>Amblyopyrum muticum</i> , <i>Comopyrum comosum</i> , <i>Crithodium monococcum</i> , <i>Crithopsis delileana</i> , <i>Dasyphyrum villosum</i> , <i>Eremopyrum distans</i> , <i>Henrardia persica</i> , <i>Heteranthelium piliferum</i> , <i>Lophopyrum elongatum</i> , <i>Peridictyon sanctum</i> , <i>Pseudoroegneria spicata</i> , <i>Secale strictum</i> , <i>Taeniatherum caput-medusae</i> , <i>Thinopyrum bessarabicum</i> , <i>Psathyrostachys fragillis</i> , <i>fragilis</i> , <i>Psathyrostachys fragillis</i> , <i>villosus</i> , <i>Festucopsis serpentinii</i> , <i>Elymus repens</i> , <i>Phleum pretense</i> , <i>Zingieria biebersteiniana</i> , <i>Z. trichopoda</i> , <i>Colpodium versicolor</i> , <i>Spartina alterniflora</i> , <i>Bromus sterilis</i> , <i>Avena sativa</i> , <i>Brachypodium distachyon</i>
Accessions used for intraspecific analyses		
25 cultivars of spring triticale (x <i>Triticosecale</i> Witt.)	Genbank Slovakia	Arc en Ciel, Fronteiro, Tentudia, Sierra de Almaraz, Sierra de Arroyo, Sierra de Lobos, Alter, Camarma, Cume, Curtido, Vrodi, Thisbi, Vrito, Legalo, Senatrit, Trimour, Dublet, Wanad, Gabo, Logo, Matejko, Nobi, Noe, Somtri, Sierra de Villueras
36 cultivars of winter triticale		Benetto, NE 422T, UCRTCL 1, Greneder, Plains, Terreland 22, Mungis, Leontino, Constant, Trizeps, Tribeca, Dusi, Alekto, Cosinus, Kandar, Largus, Magistral, Noe, Pletomax, Tatra, Massimo, Nutriseeds 1-18, Bienvenu, Trismart, Trimmer, Wilfried, Pizarro, Aprim, Innoval, Amarillo 105, Blenio, Kinerit, UCRTCL3, Algosio, UCRTCL2, Flavius
Five cultivars of rye (<i>Secale cereale</i> L.)	MTT Agrifood Research, Finland	Riähi, Akusti, Iissavaara, Loppi, Hirvessalmi;
<i>Aegilops speltoides</i> syn. <i>Sitopsis speltoides</i> (Tausch)Ä.Löve	The Institute of Evolution, Haifa	Seeds are selected randomly from the population received
<i>Triticum turgidum</i> subsp. <i>Dicoccoides</i>		

Genomic DNA was extracted from 5-day-old etiolated seedling or green leaves using CTAB buffer following established protocols (<http://primerdigital.com/dna.html>). DNA concentration was estimated by spectrophotometric analysis; DNA quality was determined by agarose gel electrophoresis.

PCR-amplification

PCR primers were designed using genomic β -amylase sequences retrieved from the NCBI database (<http://www.ncbi.nlm.nih.gov/nucleotide/>). DNA multiple sequence alignments were performed using the program, Multalin (Corpet 1988), while primer design was performed using FastPCR (Kalendar et al. 2011). After comparison of the amplification effectiveness of primer pair options, pairs 3162 and 3816 were selected: 3162 (5'-TCCAAGTCTACGTCATGCTCC-3', which anneals to *Bamy1* exon 1 position 1389-1409 bp (FJ161080), and *Bamy2* exon 1 position 54-74 bp (DQ889983) and 3816 (5'-GCTGCTGCTGCTTTGAAGTCTGCT-3', which anneals to *Bamy1* exon 1 position 3660-3683 bp (FJ161080), and *Bamy2* exon 1 position 1386-1409 bp (DQ889983).

PCRs were performed in 25 μ l reaction mixtures, containing 25 ng genomic DNA, 1x DreamTaq buffer, 0.3 μ M each primer, 0.2 mM each dNTP, 1 U DreamTaq DNA polymerase (Thermo Scientific). PCR cycling profiles were as follows: initial denaturation 95°C - 3 min, subsequently 32 cycles: 15 sec at 95°C, 60 sec at 65°C and 2 min at 72 °C, final elongation - 5 min at 72°C. Electrophoresis of the amplified products was performed using 1.3% ethidium bromide stained agarose gels and amplicons visualized using a FLA-5100 imaging system (Fuji Photo Film GmbH, Europe).

Results and Discussion

Design of EPIC primers for cereal β -amylase genes

Primer design was performed according to the complete nucleic acid sequences of the *Bamy1* and *Bamy2* genes of the barley, and from genomic gene fragments and cDNA of *Bamy1* and *Bamy2* of barley, soft wheat, rye, sorghum and maize, identified in the NCBI database (<http://www.ncbi.nlm.nih.gov/nucleotide/>). Altogether, 45 nucleic acid sequences were identified and used for subsequent analyses, GenBank accessions: AB048949, AB306504, AF012343, AF061203, AF061204, AF300799, AF300800, AF414081, AF414082, AF470353, AJ301645, AK365629, AK375028, AY111979, AY454398, AY835429, AY835430, D21349, D49999, D63574, DQ889983, EF175466, EF175467, EF175468, EF175469, EF175470, EF175471, EF175472, EF175473, EU589327, EU589328, FJ161078, FJ161079, FJ161080, FJ936153, FN179393, FN179394, GU017481, NM_001112026, NM_001188314, X52321, X56785, X98504, XM_003562919, Z11772.

DNA multiple sequence alignments of the β -amylase genes identified regions of DNA

conservation within exons, as well as variability within intronic regions (Supplemental File). The sequence alignment was used in conjunction with the program FastPCR (Kalendar et al. 2011) to design EPIC-primers which target the most conserved protein-coding regions of *Bamy1* and *Bamy2*. It should be noted that the exonic regions chosen for primer annealing show high levels of conservation not only for cereals, but also for more distant plant species, thus demonstrating the flexibility of EPIC for plant molecular studies.

Interspecific Poaceae *Bamy1* and *Bamy2* genetic variation

To study interspecific genetic variability in *Bamy1* and *Bamy2*, 38 *Poaceae* species were analysed (Tab. 2). Detection of β -amylase genetic polymorphism was performed using EPIC primers 3162 and 3816 (complementary to exons 1 and 4 in barley, respectively), allowing simultaneous study of *Bamy1* and *Bamy2* amplicons from multiple species. Analysis of EPIC PCR products amplified from all species found a wide range of genetic variation (Fig. 1). The expected amplicon size of barley PCR amplicons fragments from *H. vulgare* was 2295 or 2196 bp for *Bamy1* (due to a 126 bp MITE insertion in intron 3 (Erkkilä 1999), and 1365 bp for *Bamy2*. Amplification products of these sizes were identified in *Hordeum vulgare* (Fig. 1, samples 1-4), as well as for *H. spontaneum*, *H. murinum* and *T. durum*. Amplicons from the remaining species differed from these, ranging between 1700 and 2500 bp for *Bamy1*, and between 1300 and 1500 bp for *Bamy2* (Fig. 1). Species belonging to the tribes *Aveneae*, *Poeae* and *Brachypodieae* (*Phleum pratense*, *Zingiberia biebersteiniana*, *Colpodium versicolor*, *Spartina alterniflora*, *Avena sativa*, *Brachypodium distachyon*; samples 31-34, 36-37) contain just one β -amylase gene, *Bamy2*. Similarly, for *Bromus sterilis* (which belongs to the Bromaeae tribe), searches of public DNA databases found sequences for *Bamy1* alone (HE565904, HE565905). However, the EPIC analysis undertaken here found *B. sterilis* to possess both *Bamy1* and *Bamy2* (Fig 1, sample 35), indicating that the presence of endospermal β -amylase genes is not exclusive to the Triticeae tribe.

In total, ten different *Bamy1* alleles (from one to three amplicons per sample) and four *Bamy2* alleles (one amplicon per sample) were identified. EPIC-PCR analyses revealed both inter- and intraspecific variation in the samples analysed. Accordingly, the EPIC approach is appropriate for rapid analysis of genetic variety within the *Poaceae* family.

Triticeae β -amylase genetic variation

EPIC analyses of β -amylase intraspecific genetic variation for four members of the Triticeae tribe were undertaken: *Triticum dicoccoides*, *Aegilops speltoides*, rye and triticale.

Table 2. Analysis of *Bamy1* and *Bamy2* genes in species of the *Poaceae* family

no	Species	Tribe	<i>Bamy1</i>	<i>Bamy2</i>
1	<i>Hordeum vulgare</i>	<i>Triticeae</i>	+	+
2	<i>Hordeum spontaneum</i>	<i>Triticeae</i>	+	+
3	<i>Hordeum murinum</i>	<i>Triticeae</i>	+	+
4	<i>Hordeum marinum</i>	<i>Triticeae</i>	+	+
5	<i>Hordeum brachyantherum</i>	<i>Triticeae</i>	+	+
6	<i>Hordeum erectifolium</i>	<i>Triticeae</i>	+	+
7	<i>Triticum durum</i>	<i>Triticeae</i>	+	+
8	<i>Aegilops speltoides</i>	<i>Triticeae</i>	+	+
9	<i>Agropyron cristatum</i>	<i>Triticeae</i>	+	+
10	<i>Amblyopyrum muticum</i>	<i>Triticeae</i>	+	+
11	<i>Comopyrum comosum</i>	<i>Triticeae</i>	+	+
12	<i>Crithodium monococcum</i>	<i>Triticeae</i>	+	+
13	<i>Crithopsis delileana</i>	<i>Triticeae</i>	+	+
14	<i>Dasyphyrum villosum</i>	<i>Triticeae</i>	+	+
15	<i>Eremopyrum distans</i>	<i>Triticeae</i>	+	+
16	<i>Henrardia persica</i>	<i>Triticeae</i>	+	+
17	<i>Heteranthelium piliferum</i>	<i>Triticeae</i>	+	+
18	<i>Lophopyrum elongatum</i>	<i>Triticeae</i>	+	+
19	<i>Peridictyon sanctum</i>	<i>Triticeae</i>	+	+
20	<i>Pseudoroegneria spicata</i>	<i>Triticeae</i>	+	+
21	<i>Secale strictum</i>	<i>Triticeae</i>	+	+
22	<i>Taeniatherum caput-medusae</i>	<i>Triticeae</i>	+	+
23	<i>Thinopyrum bessarabicum</i>	<i>Triticeae</i>	+	+
24	<i>Psathyrostachys fragilis, villosus</i>	<i>Triticeae</i>	+	+
25	<i>Psathyrostachys stoloniformis</i>	<i>Triticeae</i>	+	+
26	<i>Festucopsis serpentinii</i>	<i>Triticeae</i>	+	+
27	<i>Elymus repens</i>	<i>Triticeae</i>	+	+
28	<i>Bromus sterilis</i>	<i>Bromeae</i>	+	+
29	<i>Avena sativa</i> (cv. Vali)	<i>Aveneae</i>	-	+
30	<i>Zingeria biebersteiniana</i>	<i>Aveneae</i>	-	+
31	<i>Zingeria trichopoda</i>	<i>Aveneae</i>	-	+
32	<i>Colpodium versicolor</i>	<i>Poeae</i>	-	+
33	<i>Pbleum pratense</i>	<i>Poeae</i>	-	+
34	<i>Spartina alterniflora</i>	<i>Zoysieae</i>	-	+
35	<i>Brachypodium distachyon</i>	<i>Brachypodieae</i>	-	+

Figure 1. Electrophoregram products amplified from *BamY1* and *BamY2* in species of the family, *Poaceae*. M - DNA ladder (SM1173, Thermo Scientific);

1-4 *Hordeum vulgare*: 1 - cv. Djau Kabutak; 2 - cv. Pallidum 107; 3 - cv. Odesskiy 17; 4 - cv. Odesskiy 31; 5 - *H. spontaneum*; 6 - *H. murinum*; 7 - *H. marinum*; 8 - *H. brachyantherum*; 9 - *H. leporinum*; 10 - *Triticum durum* (AABB); 11 - *Aegilops speltoides*; 12 - *Agropyron cristatum*; 13 - *Amblyopyrum muticum*; 14 - *Comopyrum comosum*; 15 - *Critobodium monococcum*; 16 - *Critobopsis delileana*; 17 - *Dasyphyrum villosum*; 18 - *Eremopyrum distans*; 19 - *Henrardia persica*; 20 - *Heteranthelium piliferum*; 21 - *Lophopyrum elongatum*; 22 - *Peridictyon sanctum*; 23 - *Pseudoroegneria spicata*; 24 - *Secale strictum*; 25 - *Taeniatherum caput-medusae*; 26 - *Thinopyrum bessarabicum*; 27 - *Psathyrostachys fragilis*; 28 - *Psathyrostachys fragilis*, *villosus*; 29 - *Festucopsis serpentini*; 30 - *Elymus repens*; 31 - *Phleum pratense*; 32 - *Zingiber biebersteiniana*; 33 - *Colpodium versicolor*; 34 - *Spartina alterniflora*; 35 - *Bromus sterilis*; 36 - *Avena sativa*; 37 - *Brachypodium distachyon*.

These species were selected for the following reasons: *T. dicoccoides* (emmer or spelt wheat) is the tetraploid parent of the cultivated hexaploid wheat *T. aestivum* ($2n = 6x = 42$, AABBDD), while *A. speltoides* ($2n = 2x = 14$, BB or SS) carries valuable agronomic attributes exploited within the artificial hybridization with bread wheat. As well as being a valuable crop in its own right, rye is used to form a hybrid with wheat to produce triticale, a crop which combines the valuable characteristics of these genera (Lukaszewski 2006, Yang et al. 2011).

Three EPIC amplicons were found to be common for all four species investigated: products of 2295 bp (as expected for *BamY1*), ~1900 bp (included in the allelomorphic spectrum of *BamY1*) and 1356 bp (*BamY2*) (Fig. 2, 3, 4). It should be noted that EPIC amplicons of size 2295 and 1356 bp were also present in the

allelomorphic spectrum of *Hordeum* (samples 1 - 6) and *T. durum* (10) (Fig. 1). Analysis of *BamY1* polymorphism in triticale and rye showed that their level of common polymorphism is higher than those of the cultivars of barley (Fig. 1), in which only two *BamY1* alleles were found. Comparison of triticale and rye genotypes showed that the majority of the triticale allelomorphic spectrum was present in rye (including fragments of 2295 and 1356 bp). In addition, the allelomorphic spectrums of winter and spring triticale contained amplicons of ~2000 bp (Fig. 2 – samples no. 9, 17, 18, 22, 29 and 34). These were absent in rye, and as they were identified here in wheat, it is likely they originate from the wheat parent (Fig. 4 A). The 1850 bp amplicon found in the triticale samples was absent from rye and wheat, indicating this allele was not captured in the donor species samples screened in this study.

Figure 2. *BamY1* and *BamY2* genetic variation identified by EPIC in winter cultivars of triticale and rye. M - DNA ladder (SM1173); 1-36 cultivars of winter triticale: 1 - Benetto; 2 - NE 422T; 3 - UCRTCL1; 4 - Greneder; 5 - Plains; 6 - Terreland 22 7 - Mungis; 8 - Leontino; 9 - Constant; 10 - Trizeps; 11 - Tribeca; 12 - Dusi; 13 - Alekto; 14 - Cosinus; 15 - Kandar; 16 - Largus; 17 - Magistral; 18 - Noe; 19 - Pletomax; 20 - Tatra; 21 - Massimo; 22 - Nutriseeds 1 - 18; 23 - Bienvenu; 24 - Trismart; 25 - Trimmer; 26 - Wilfried; 27 - Pizarro; 28 - Aprim; 29 - Innoval; 30 - Amanillo 105; 31 - Blenio; 32 - Kinerit; 33 - UCRTCL3; 34 - Algosio; 35 - UCRTCL2; 36 - Flavius; 37-41 cultivars of rye: 37 - Rihi; 38 - Akusti; 39 - Iissavaara; 40 - Loppi; 41 - Hirvessalmi.

Up to eight *Bamy1* polymorphic PCR products (maximum per accession = 4) and three *Bamy2* variants (maximum per accession = 2) were detected in the samples of winter triticale analysed. For spring triticale, EPIC primers for *Bamy1* and *Bamy2* identified a total of six (maximum per accession = 3) and two amplicons (maximum per accession = 1), respectively. For triticale *Bamy1*, the frequency of the 2295 bp fragments varied with the seasonal growth habit (SGH) classification of the accessions studied: it is found in 86% of the winter cultivars, while it is present in just 48% of spring lines. The *Bamy1* 2000 bp fragment varied in frequency

between SGH classes, being present in 52% of spring triticale, but found in just 13% of winter lines. Similarly, the 1900 bp *Bamy1* amplicon is found in 56% of spring triticale cultivars, and in 27% of winter lines (Fig. 2 and 3). EPIC PCR products of ~1700 bp (found in cultivars Plains, Constant and Flavius) and ~1750 bp (cultivar Tatra) were found only in winter triticale. The most common *Bamy2* variant identified in the triticale cultivars analysed was the 1356 bp fragment. For rye, six *Bamy1* (from one to three amplicons per cultivar) and two *Bamy2* (from one to two per cultivar) products were found.

Figure 3. *Bamy1* and *Bamy2* genetic variation identified by EPIC in spring cultivars of triticale and rye.

M - DNA ladder (SM1173);

1-25 cultivars of spring triticale: 1 - Arc en Ciel; 2 - Fronteiro; 3 - Tentudia; 4 - Sierra de Almaraz; 5 - Sierra de Arroyo; 6 - Sierra de Lobos; 7 - Alter; 8 - Camarma; 9 - Cume; 10 - Curtido; 11 - Vrodi; 12 - Thisbi; 13 - Vrito; 14 - Legalo; 15 - Senatrit; 16 - Trimour; 17 - Dublet; 18 - Wanad; 19 - Gabo; 20 - Logo; 21 - Matejko; 22 - Nob; 23 - Noe; 24 - Somtri; 25 - Sierra de Villueras; 26-30 cultivars of rye: 26 - Rihi; 27 - Akusti; 28 - Iissavaara; 29 - Loppi; 30 - Hirvessalmi.

EPIC analysis of β -amylase genes from Emmer wheat revealed that *T. dicoccoides* lines commonly possess two *Bamy1* amplicons: the first (~1900 bp) was found to be present in all *T. dicoccoides* investigated, while amplicons of ~2000 bp and 2296 bp were observed in 79% and 16 % of samples, respectively (Fig. 4A). Among the 42 *Aegilops speltoides* accessions investigated, three and two amplicons were found for *Bamy1* and *Bamy2*, respectively. In addition to amplicons of 2295 and 1900 bp (found also in *T.*

turgidum, triticale, rye and barley), products of ~2600 bp (not found in any of the other species investigated) were also identified in *A. speltoides*, where it was found in 63% of accessions (Fig. 4B). Comparison of the molecular profiles of *T. dicoccoides* and *Aegilops speltoides* highlighted the presence of common *Bamy1* amplicon sizes between these species, of 2295 and 1900 bp (Fig. 4B). The 1900 bp amplification fragment was found in 16% of *A. speltoides* accessions and in 100% of *T. dicoccoides* lines (Fig. 4A-B).

Figure 4. *Bamy1* and *Bamy2* genetic variation identified by EPIC in emmer wheat and *Aegilops speltoides* accessions. M - DNA ladder (SM1173);

A – wild wheat lines *T. turgidum* subsp. *dicoccoides*;

B – *T. dicoccoides* (1-7) and *A. speltoides* (8-38).

Conclusion

Conservation of exonic regions allowed EPIC to be used to analyse intra- and interspecific polymorphism within β -amylase genes from multiple cereal species. Here, the ability to use EPIC for the analysis of a wide taxonomic range within the *Poaceae* was demonstrated. Thus, genomic data obtained from one or more organisms can be effectively used to rapidly design EPIC-markers for additional related species. This work demonstrates how EPIC can be used to determine inter- and intra-specific genetic variation within plant taxonomic groups, and could represent a useful tool for marker-assisted selection of novel β -amylase alleles within breeding programmes.

References

- Bierné, N., Lehnert, S.A., Bedier, E., Bonhomme, F., & Moore, S.S. (2000). Screening for intron-length polymorphisms in penaeid shrimps using exon-primed intron-crossing (EPIC)-PCR. *Molecular Ecology* 9: 233-235. doi:10.1046/j.1365-294x.2000.00842.x
- Braglia, L., Manca, A., Mastromauro, F., & Breviario, D. (2010). cTBP: A Successful Intron Length Polymorphism (ILP)-Based Genotyping Method Targeted to Well Defined Experimental Needs. *Diversity* 2: 572-585. doi:10.3390/d2040572
- Briggs, D.E. (1998). *Malts and Malting*. Blackie Academic and Professional, New York.
- Chetelat, R.T., Deverna, J.W., & Bennett, A.B. (1995). Introgression into tomato (*Lycopersicon esculentum*) of the *L. chmielewskii* sucrose accumulator gene (sucr) controlling fruit sugar composition. *Theoretical and Applied Genetics* 91: 327-333. doi:10.1007/BF00220895
- Cockram, J., Mackay, I.J., & O'Sullivan, D.M. (2007). The role of double-stranded break repair in the creation of phenotypic diversity at cereal VRN1 loci. *Genetics* 177: 2535-2539. doi:10.1534/genetics.107.074765
- Corpet, F. (1988). Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Research* 16: 10881-10890. doi:10.1093/nar/16.22.10881
- Erkkilä, M.J. (1999). Intron III-Specific Markers for Screening of β -amylase Alleles in Barley Cultivars. *Plant Molecular Biology Reporter* 17 (2): 139-147. doi:10.1023/A:1007595821379
- Fridman, E., Pleban, T., & Zamir, D. (2000). A recombination hotspot delimits a wild-species quantitative trait locus for tomato sugar content to 484 bp within an invertase gene. *Proceedings of the National Academy of Sciences* 97: 4718-4723. doi:10.1073/pnas.97.9.4718
- Gardes, M.B.T. (1993). ITS primers with enhanced specificity for basidiomycetes—application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2:113-118
- Holland, J.B., Helland, S.J., Sharopova, N., & Rhyne, D.C. (2001). Polymorphism of PCR-based markers targeting exons, introns, promoter regions, and SSRs in maize and introns and repeat sequences in oat. *Genome* 44: 1065-1076.
- Hongtrakul, V. S.M., & Knapp, S.J. (1998). DFLP, SSCP, and SSR markers for delta 9-stearoyl-acyl carrier protein desaturases strongly expressed in developing seeds of sunflower: intron lengths are polymorphic among elite inbred lines. *Molecular Breeding* 4: 195-203.
- Ishikawa, G., Yonemaru, J., Saito, M., & Nakamura, T. (2007). PCR-based landmark unique gene (PLUG) markers effectively assign homoeologous wheat genes to A, B and D genomes. *BMC Genomics* 8: 135. doi:10.1186/1471-2164-8-135
- James, M.G., Denyer, K., & Myers, A.M. (2003). Starch synthesis in the cereal endosperm. *Current Opinion in Plant Biology* 6: 215-222. doi:10.1016/s1369-5266(03)00042-6
- Kalendar, R., Lee, D., & Schulman, A.H. (2011). Java web tools for PCR, *in silico* PCR, and oligonucleotide assembly and analysis. *Genomics* 98: 137-144. doi:10.1016/j.ygeno.2011.04.009
- Lessa, E.P. (1992). Rapid surveying of DNA sequence variation in natural populations. *Molecular Biology and Evolution* 9: 323-330.
- Li, C., Riethoven, J.J., & Ma, L. (2010). Exon-primed intron-crossing (EPIC) markers for non-model teleost fishes. *BMC Evolutionary Biology* 10: 90. doi:10.1186/1471-2148-10-90
- Ludwig, M. (2002). Functional evolution of noncoding DNA. *Current Opinion in Genetics & Development* 12: 634-639. doi:10.1016/s0959-437x(02)00355-6
- Lukaszewski, A.J. (2006). Cytogenetically Engineered Rye Chromosomes 1R to Improve Bread-making Quality of Hexaploid Triticale. *Crop Science* 46: 2183. doi:10.2135/cropsci2006.03.0135
- Mason-Gamer, R.J. (2005). The β -amylase genes of grasses and a phylogenetic analysis of the Triticaceae (Poaceae). *American Journal of Botany* 92: 1045-1058. doi:10.3732/ajb.92.6.1045
- Morello, L., & Breviario, D. (2008). Plant spliceosomal introns: not only cut and paste. *Current Genomics* 9: 227-238. doi:10.2174/138920208784533629
- Palumbi, S.R., & Baker, C.S. (1994). Contrasting population structure from nuclear intron sequences and mtDNA of humpback whales. *Molecular Biology and Evolution* 11: 426-435.
- Priest, F.G. (1986). Toward better beer. In: J. S. Hough (Ed), *The Biotechnology of Malting and Brewing*. Cambridge University Press, 1985, pp. 188. BioEssays 4:139-139 doi:10.1002/bies.950040313
- Rose, A.B., & Beliakoff, J.A. (2000). Intron-mediated enhancement of gene expression independent of unique intron sequences and splicing. *Plant Physiology* 122: 535-542 doi:10.1104/pp.122.2.535
- Yang, E.N., Yang, Z.J., Zhang, J.F., Zou, Y.C., & Ren, Z.L. (2011). Molecular cytogenetic characterization of a new leaf rolling triticale. *Genetics and Molecular Research: GMR* 10: 2953-2961 doi:10.4238/2011. November 29.6
- Yang, L., Jin, G., Zhao, X., Zheng, Y., Xu, Z., & Wu, W. (2007). PIP: a database of potential intron polymorphism markers. *Bioinformatics* 23: 2174-2177. doi:10.1093/bioinformatics/btm296

Molekularna genetička analiza gena za β -amilazu korišćenjem Exon-Primed Intron-Crossing (EPIC) PCR

Olga Stratula · James Cockram · Ruslan Kalendar

Sažetak: Proteini koje kodiraju geni za β -amilazu kod strnih žita *Bamy1* i *Bamy2*, igraju veoma značajnu ulogu u klijanju semena i procesu proizvodnje piva. U ovom radu, korišćeni su exon-primed intron-crossing (EPIC) markeri za analizu genetičke divergentnosti *Bamy1* i *Bamy2* gena između 38 genotipova, klasifikovanih u 6 tribusa familije *Poaceae*. Podudarnost DNK sekvenci za β -amilazu kod različitih *Poaceae* vrsta omogućila je dizajniranje EPIC prajmera koji istovremeno amplifikuju *Bamy1* i *Bamy2* sekvence u svim ispitivanim vrstama žitarica. Utvrđena genetička varijabilnost u ispitivanim uzorcima je analizirana i diskutovana i ilustruje efikasnost ovog pristupa za intra- i interspecies analize različitih biljnih vrsta.

Gljučne reči: β -amilaza, genetska raznovrsnost, ječam, *Poaceae*, polimorfizam dužine introna, pšenica, raž

Figure S1 Multiple alignments of β -amylase genes sequences. Exons are marked in red.

FJ161080	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
AF061204	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
FJ936153	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
FJ161078	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
AJ301645	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
FJ161079	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
D49999	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
EJ589327	-----
EJ589328	-----
DQ889983	-----
AK375028	-----
AF012343	-----
FJ179394	-----
AK365629	-----
EJ11772	-----
X98504	-----
AY835429	-----
AY835430	-----
XM_003562919	-----
NM_001112026	-----
GU017481	-----
D63574	AAATGTTTAACC--TTTTACCTCAAATTTGCAGGTAGCATTAG--ATATACCTAAGTACATAAATAAATTTGTCTTGATTTTTTTCATTTGAATTTTTTGGCCCCG--AAGCATATTCTTCGGGAGCCAAATTGACATTCGGTCATGATGTGGCTTGGATCCCAAGTTAGTTATACAGATAAGGATATAT
AF470353	-----
	810 820 830 840 850 860 870 880 890 900 910 920 930 940 950 960 970 980 990 1000
AB048949	-----
AF300799	-----
AF300800	-----
AF414081	-----
AF414082	-----
D21349	-----
X52321	-----
FJ179393	-----
AB306504	-----
EF175466	-----
EF175470	-----
EF175468	-----
EF175467	-----
EF175473	-----
EF175469	-----
EF175471	-----
EF175472	-----
AF061203	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
FJ161080	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
AF061204	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
FJ936153	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
FJ161078	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
AJ301645	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
FJ161079	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
D49999	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
EJ589327	-----
EJ589328	-----
DQ889983	-----
AK375028	-----
AF012343	-----
FJ179394	-----
AK365629	-----
EJ11772	-----
X98504	-----
AY835429	-----
AY835430	-----
XM_003562919	-----
NM_001112026	-----
GU017481	-----
D63574	CTTACCTCAACGGAATCTAGGTTACACAAGCTTAAACACTCATGCAATAGTGCCATCTAGACTCTCTTAGAAGGCAATGGTTTACACACCAATTTGGGTTGTACAAATCCCAACACTCTTCTCAATCATAACTTTAAGAAGTCAGAGATTGATCTTAAAGTTCTGATGTTATCTGTTTAAAG--AATAGCTTGGGAACA
AF470353	-----
	1010 1020 1030 1040 1050 1060 1070 1080 1090 1100 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200
AB048949	-----
AF300799	-----
AF300800	-----
AF414081	-----
AF414082	-----
D21349	-----
X52321	-----
FJ179393	-----
AB306504	-----
EF175466	-----
EF175470	-----
EF175468	-----
EF175467	-----
EF175473	-----
EF175469	-----
EF175471	-----
EF175472	-----
AF061203	TATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
FJ161080	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
AF061204	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
FJ936153	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
FJ161078	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
AJ301645	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
FJ161079	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
D49999	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
EJ589327	-----
EJ589328	-----
DQ889983	-----
AK375028	-----
AF012343	-----
FJ179394	-----
AK365629	-----
EJ11772	-----
X98504	-----
AY835429	-----
AY835430	-----
XM_003562919	-----
NM_001112026	-----
GU017481	-----
D63574	CATGTACAACTTAAGTCAGGATGTGGCGCTTTCACCTTGTATAGGGTGGCGTTTGGTTGAGAGTGGAGAGCGTGTATGGGTCACAGCTTGGTATATAATAGATAATGATCCCACTCTCTCCCAATCAGAGCTCAATCTTGAAGTTTGGCTGTGTGGTGAAGAAACCAAAATAGTTACAAAAGCGCCCTTTCT
AF470353	-----
	1210 1220 1230 1240 1250 1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360 1370 1380 1390 1400
AB048949	-----
AF300799	-----
AF300800	-----
AF414081	-----
AF414082	-----
D21349	-----
X52321	-----
FJ179393	-----
AB306504	-----
EF175466	-----
EF175470	-----
EF175468	-----
EF175467	-----
EF175473	-----
EF175469	-----
EF175471	-----
EF175472	-----
AF061203	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
FJ161080	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
AF061204	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
FJ936153	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
FJ161078	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
AJ301645	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
FJ161079	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
D49999	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
EJ589327	-----
EJ589328	-----
DQ889983	-----
AK375028	-----
AF012343	-----
FJ179394	-----
AK365629	-----
EJ11772	-----
X98504	-----
AY835429	-----
AY835430	-----
XM_003562919	-----
NM_001112026	-----
GU017481	-----
D63574	GAGTGACAAGCCTATTTCTGACATATCTG--ATTGGAAAGCTAGGTTGGCGCTTGGCTCACAATCTAGGATACATCTTTTCATAGTATAAATAGACCTTTTATTAAGTCCTCGCATATCCAAACAAACCTTTGAAGTTGTAGAGCATATCATATAGCAGCATCCCAATGGAGTTGAACGTGAAGGCCAATCA
AF470353	-----

3162 5-TCCAAGTCTACGTCATGCTCC->

.....

=====

[illegible]

[illegible]

[illegible]

A0048949
 AF300799
 AF300800
 AF314081
 AF314082
 D21349
 X25231
 FN179393
 FN179394
 FN175466
 FN175467
 FN175468
 FN175469
 FN175470
 FN175471
 FN175472
 FN175473
 FN175474
 FN175475
 FN175476
 FN175477
 FN175478
 FN175479
 FN175480
 FN175481
 FN175482
 FN175483
 FN175484
 FN175485
 FN175486
 FN175487
 FN175488
 FN175489
 FN175490
 FN175491
 FN175492
 FN175493
 FN175494
 FN175495
 FN175496
 FN175497
 FN175498
 FN175499
 FN175500
 FN175501
 FN175502
 FN175503
 FN175504
 FN175505
 FN175506
 FN175507
 FN175508
 FN175509
 FN175510
 FN175511
 FN175512
 FN175513
 FN175514
 FN175515
 FN175516
 FN175517
 FN175518
 FN175519
 FN175520
 FN175521
 FN175522
 FN175523
 FN175524
 FN175525
 FN175526
 FN175527
 FN175528
 FN175529
 FN175530
 FN175531
 FN175532
 FN175533
 FN175534
 FN175535
 FN175536
 FN175537
 FN175538
 FN175539
 FN175540
 FN175541
 FN175542
 FN175543
 FN175544
 FN175545
 FN175546
 FN175547
 FN175548
 FN175549
 FN175550
 FN175551
 FN175552
 FN175553
 FN175554
 FN175555
 FN175556
 FN175557
 FN175558
 FN175559
 FN175560
 FN175561
 FN175562
 FN175563
 FN175564
 FN175565
 FN175566
 FN175567
 FN175568
 FN175569
 FN175570
 FN175571
 FN175572
 FN175573
 FN175574
 FN175575
 FN175576
 FN175577
 FN175578
 FN175579
 FN175580
 FN175581
 FN175582
 FN175583
 FN175584
 FN175585
 FN175586
 FN175587
 FN175588
 FN175589
 FN175590
 FN175591
 FN175592
 FN175593
 FN175594
 FN175595
 FN175596
 FN175597
 FN175598
 FN175599
 FN175600
 FN175601
 FN175602
 FN175603
 FN175604
 FN175605
 FN175606
 FN175607
 FN175608
 FN175609
 FN175610
 FN175611
 FN175612
 FN175613
 FN175614
 FN175615
 FN175616
 FN175617
 FN175618
 FN175619
 FN175620
 FN175621
 FN175622
 FN175623
 FN175624
 FN175625
 FN175626
 FN175627
 FN175628
 FN175629
 FN175630
 FN175631
 FN175632
 FN175633
 FN175634
 FN175635
 FN175636
 FN175637
 FN175638
 FN175639
 FN175640
 FN175641
 FN175642
 FN175643
 FN175644
 FN175645
 FN175646
 FN175647
 FN175648
 FN175649
 FN175650
 FN175651
 FN175652
 FN175653
 FN175654
 FN175655
 FN175656
 FN175657
 FN175658
 FN175659
 FN175660
 FN175661
 FN175662
 FN175663
 FN175664
 FN175665
 FN175666
 FN175667
 FN175668
 FN175669
 FN175670
 FN175671
 FN175672
 FN175673
 FN175674
 FN175675
 FN175676
 FN175677
 FN175678
 FN175679
 FN175680
 FN175681
 FN175682
 FN175683
 FN175684
 FN175685
 FN175686
 FN175687
 FN175688
 FN175689
 FN175690
 FN175691
 FN175692
 FN175693
 FN175694
 FN175695
 FN175696
 FN175697
 FN175698
 FN175699
 FN175700
 FN175701
 FN175702
 FN175703
 FN175704
 FN175705
 FN175706
 FN175707
 FN175708
 FN175709
 FN175710
 FN175711
 FN175712
 FN175713
 FN175714
 FN175715
 FN175716
 FN175717
 FN175718
 FN175719
 FN175720
 FN175721
 FN175722
 FN175723
 FN175724
 FN175725
 FN175726
 FN175727
 FN175728
 FN175729
 FN175730
 FN175731
 FN175732
 FN175733
 FN175734
 FN175735
 FN175736
 FN175737
 FN175738
 FN175739
 FN175740
 FN175741
 FN175742
 FN175743
 FN175744
 FN175745
 FN175746
 FN175747
 FN175748
 FN175749
 FN175750
 FN175751
 FN175752
 FN175753
 FN175754
 FN175755
 FN175756
 FN175757
 FN175758
 FN175759
 FN175760
 FN175761
 FN175762
 FN175763
 FN175764
 FN175765
 FN175766
 FN175767
 FN175768
 FN175769
 FN175770
 FN175771
 FN175772
 FN175773
 FN175774
 FN175775
 FN175776
 FN175777
 FN175778
 FN175779
 FN175780
 FN175781
 FN175782
 FN175783
 FN175784
 FN175785
 FN175786
 FN175787
 FN175788
 FN175789
 FN175790
 FN175791
 FN175792
 FN175793
 FN175794
 FN175795
 FN175796
 FN175797
 FN175798
 FN175799
 FN175800
 FN175801
 FN175802
 FN175803
 FN175804
 FN175805
 FN175806
 FN175807
 FN175808
 FN175809
 FN175810
 FN175811
 FN175812
 FN175813
 FN175814
 FN175815
 FN175816
 FN175817
 FN175818
 FN175819
 FN175820
 FN175821
 FN175822
 FN175823
 FN175824
 FN175825
 FN175826
 FN175827
 FN175828
 FN

[illegible][illegible]

Ratar. Povrt. 51:3 (2014) 175-189

[illegible]